

References

Alster, T.S. (1994):

Improvement of erythematous and hypertrophic scars by the 585 nm pulsed dye laser. *Ann Plast Surg*; 32:186– 190.

Alster, T.S. and Tanzi, E.L. (2003):

Hypertrophic scars and keloids: etiology and management. *Am J Clin Dermatol*; 4(4):235- 243.

Altavilla, D.; Saitta, A. and Cucinotta, D. (2001):

Inhibition of lipid peroxidation restores impaired vascular endothelial growth factor expression and stimulates wound healing and angiogenesis in the genetically diabetic mouse. *Diabetes*; 50(3):667 –674.

Amadeu, T.; Braune, A.; Mandarim-de-Lacerda, C.; Porto, L.C.; Desmouliere, A. and Costa, A. (2003):

Vascularization pattern in hypertrophic scars and keloids: a stereological analysis. *Pathol Res Pract*; 199(7):469-473.

Amadeu, T.P.; Braune, A.S.; Porto, L.C.; Desmouliere, A. and Costa, A.M. (2004):

Fibrillin-1 and elastin are differentially expressed in hypertrophic scars and keloids. *Wound Repair Regen*; Mar-Apr; 12(2):169-174.

Baumann, L.S., and Spencer, J. (1999):

The effects of topical vitamin E on the cosmetic appearance of scars. *Dermatol Surg*; 25(4):311 – 315.

Bayat, A.; Arscott, G.; Ollier, W.E.; Ferguson, M.W. and Mc Grouther, D.A. (2004):

Description of site-specific morphology of keloid phenotypes in an Afrocaribbean population. Br J Plast Surg. Mar; 57(2):122-133.

Bennett, R.G. (2002):

Anatomy and Physiology of the Skin. In: Facial Plastic and Reconstructive Surgery; 2nd edition; (ed.) by ***Papel, I.D.*** New York, Stuttgart: Thieme. PP: 3-15.

Berman, B. and Bielely, H.C. (1995):

Keloids. J Am Acad Dermatol; 33:117– 123.

Berman, B. and Bielely, H.C. (1996):

Adjunct therapies to surgical management of keloids. Dermatol Surg; 22:126–130.

Berman, B. and Flores, F. (1997):

Recurrence rates of excised keloids treated with postoperative triamcinolone acetonide injections or interferon alfa-2b injections. J Am Acad Dermatol; 37:755– 777.

Berman, B. and Villa, A. (2003):

Imiquimod 5% cream for keloid management. Dermatol Surg; 29(10):1050-1051.

Buck, M.; Houglum, K. and Chojkier, M. (1996):

Tumor necrosis factor-alpha inhibits collagen alpha a (I) gene expression and wound healing in a murine model of cachexia. Am J Pathol; 149:195- 204.

Bullard, K.M.; Lund, L. and Mudgett, J.S. (1999):

Impaired wound contraction in stromelysin-1 deficient mice. *Ann Surg*; 230:260–265.

Burns, J.L.; Mancoll, J.S. and Phillips, L.G. (2003):

Impairments to wound healing. *Clin Plastic Surg* 30(1): 47– 56.

Capon, A. and Mordon, S. (2003):

Can thermal lasers promote skin wound healing? *Am J Clin Dermatol*; 4(1):1-12.

Carroll, L.A. and Koch, R.J. (2003):

Heparin stimulates production of bFGF and TGF-beta 1 by human normal, keloid, and fetal dermal fibroblasts. *Med Sci Monit*; 9(3): 97-108.

Chang, P.; Laubenthal, K.N. and Lewis, R.W. (1995):

Prospective, randomized study of the efficacy of pressure garment therapy in patients with burns. *J Burn Care Rehabil*; 16:473– 480.

Cheng, E.T.; Pollard, J.D. and Koch, R.J. (2003):

Effect of blended CO₂ and Erbium: YAG laser irradiation on normal and keloid fibroblasts: a serum-free study. *J Clin Laser Med Surg*; 21(6):337-343.

Chin, G.S.; Liu, W. and Peled, Z. (2001):

Differential expression of transforming growth factor-beta receptors I and II and activation of Smad 3 in keloid fibroblasts. *Plast Reconstr Surg*; 108(2):423–429.

Chou, T.D.; Chen, S.L. and Lee, T.W. (2001):

Reconstruction of burn scar of the upper extremities with artificial skin. *Plast Reconstr Surg*; 108(2):378– 384.

Cines, D.B.; Pollack, E.S. and Buck, C.A. (1998):

Endothelial cells in physiology and in the pathophysiology of vascular disorders. *Blood*; 91:3527– 3561.

Cohen, I.K.; Diegelman, R.F. and Dome, R.Y. (1999):

Wound care and wound healing. In: *Principles of Surgery*; 7th edition; (ed.) by ***Schwartz, S.I.; Shires, G.T.; and Spencer, F.C.*** New York: McGraw-Hill. PP: 263-295.

Coleman, D.J. (2004):

Wounds, tissue repair and scars. In: *Bailey and Love's Short Practice of Surgery*; 24th edition; (ed.) by ***Russel, R.C.G.; Williams, N.S. and Bulstrode, C.J.K.*** London: Arnold. PP: 84- 95.

Connors, D.; Gies, D. and Lin, H. (2000):

Increase in wound breaking strength in rats in the presence of positively charged dextran beads correlates with an increase in endogenous transforming growth factor-beta1 and its receptor TGF-beta RI in close proximity to the wound. *Wound Repair Regen*; 8(4):292 –303.

Copcu, E.; Sivrioglu, N. and Oztan, Y. (2004):

Combination of surgery and intralesional verapamil injection in the treatment of the keloid. *J Burn Care Rehabil*; 25(1):1-7.

Cosman, B. and Wolff, M. (1974):

Bilateral earlobe keloids. *Plast Reconstr Surg*; 53:540– 543.

Daian, T.; Ohtsuru, A.; Rogounovitch, T.; Ishihara, H. and Hirano, A. (2003):

Insulin-like growth factor-I enhances transforming growth factor-beta-induced extracellular matrix protein production through the P38/activating transcription factor-2 signaling pathway in keloid fibroblasts. *J Invest Dermatol*; 120(6):956-962.

Dalkowski, A.; Fimmel, S.; Beutler, C. and Zouboulis, C.C. (2003):

Cryotherapy modifies synthetic activity and differentiation of keloidal fibroblasts in vitro. *Exp Dermatol*. 2003 Oct; 12(5):673-681.

Darzi, M.A.; Chowdri, N.A. and Kaul, S.K. (1992):

Evaluation of various methods of treating keloids and hypertrophic scars: a 10-year follow up study. *Br J Plast Surg*; 45:374– 379.

Doong, H.; Myerov, A. and Lee, R.C. (1993):

Calcium channel blockers depolymerize fibroblast actin to induce collagenase synthesis. *Surg Forum*; 44:738– 740.

Ehrlich, H.P.; Desmouliere, A. and Diegelmann, R.F. (1994):

Morphological and immunochemical differences between keloids and hypertrophic scars. *Am J Pathol*; 145:105–113.

Eishi, K.; Bae, S.J.; Ogawa, F.; Hamasaki, Y.; Shimizu, K. and Katayama, I. (2003):

Silicone gel sheets relieve pain and pruritus with clinical improvement of keloid: possible target of mast cells. J Dermatolog Treat; 14(4):248-252.

Fauci, A.S. and Haynes, B.F. (1994):

Cellular and molecular basis of immunity. In: Harrison's Principles of Internal Medicine; 13th edition; (ed.) by ***Issclacher, K.; Braunwald, E. and Wilson, J.*** New York: McGraw-Hill. PP: 1552.

Fisher, E.J. and Frodel, J.L.; (2002):

Wound healing. In: In: Facial Plastic and Reconstructive Surgery; 2nd edition; (ed.) by ***Papel, I.D.*** New York, Stuttgart: Thieme. PP: 15-25.

Fulton, J.E. (1995):

Silicone gel sheeting for the prevention and management of evolving hypertrophic and keloid scars. Dermatol Surg; 21:947– 951.

Funayama, E.; Chodon, T.; Oyama, A. and Sugihara, T. (2003):

Keratinocytes promote proliferation and inhibit apoptosis of the underlying fibroblasts: an important role in the pathogenesis of keloid. J Invest Dermatol; 121(6):1326- 1331.

Gaboriau, H.P. and Murakami, C.S. (2001):

Skin anatomy and flab physiology. Otolaryngologic Clinic of North America, Vol. 34 (3), 555-569.

Galeano, M.; Torre, V. and Deodato, B. (2001):

Rexofelast, a hydrophilic vitamin E-like antioxidant, stimulates wound healing in genetically diabetic mice. *Surgery* 2001; 129(4):467–477.

Garner, W.L. (1998):

Epidermal regulation of dermal fibroblast activity. *Plast Reconstr Surg*; 102(1):135–139.

Gira, A.K.; Brown, L.F.; Washington, C.V.; Cohen, C. and Arbiser, J.L. (2004):

Keloids demonstrate high-level epidermal expression of vascular endothelial growth factor. *J Am Acad Dermatol*; 50(6):850-853.

Giugliano, G.; Pasquali, D.; Notaro, A.; Brongo, S.; Nicoletti, G. and D'Andrea, F. (2003):

Verapamil inhibits interleukin-6 and vascular endothelial growth factor production in primary cultures of keloid fibroblasts. *Br J Plast Surg*; 56(8):804-809.

Glat, P.M. and Longaker, M.T. (1997):

Wound healing. In: Grabb and Smith's Plastic Surgery; 5th edition; (ed.) by **Aston, S.J.; Besley, R.W. and Thorn, C.H.** Philadelphia: Lippincott-Raven. PP: 3.

Granstein, R.D.; Rook, A. and Flotte, T.J. (1990):

Controlled trial of intralesional recombinant interferon- γ in the treatment of keloidal scarring. *Arch Dermatol*; 126:1295– 1302.

Han, H. and Mustoe, T.A. (2000):

Structure and function of the skin. In: Plastic Surgery: Indications, Operations and Outcome; 1st edition; (ed.) by ***Ashour, B.M. and Eriksson, E.*** St. Louis, London, Philadelphia: Mosby. PP: 65- 79.

Hanasono, M.M.; Kita, M.; Mikulec, A.A.; Lonergan, D. and Koch, R.J. (2003):

Autocrine growth factor production by fetal, keloid, and normal dermal Fibroblasts. Arch Facial Plast Surg; 5(1):26-30.

Hanasono, M.M.; Lum, J.; Carroll, L.A.; Mikulec, A.A. and Koch, R.J. (2004):

The effect of silicone gel on basic fibroblast growth factor levels in fibroblast cell culture. Arch Facial Plast Surg; 6(2):88-93.

Haroon, Z.A.; Raleigh, J.A. and Greenberg, C.S. (2000):

Early wound healing exhibits cytokine surge without evidence of hypoxia. Ann Surg; 231:137– 147.

Har-Shai, Y.; Amar, M. and Sabo, E. (2003):

Intralesional cryotherapy for enhancing the involution of hypertrophic scars and keloids. Plast Reconstr Surg; 111(6):1841-1852.

Hasegawa, T.; Nakao, A.; Sumiyoshi, K.; Tsuboi, R. and Ogawa, H. (2003):

IFN-gamma fails to antagonize fibrotic effect of TGF-beta on keloid-derived dermal fibroblast. J Dermatol Sci; 32(1):19-24.

Haws, M.J.; Kucan, J.O. and Roth, A.C. (1996):

The effects of chronic ketorolac tromethamine (toradol) on wound healing. *Ann Plast Surg*; 37(2):147–151.

Hunt, J.A.; Moisidis, E. and Haertsch, P. (2004):

Initial experience of Integra in the treatment of post-burn anterior cervical neck contracture. *Br J Plast Surg*; 53(8):652–658.

Iocono, J.A.; Colleran, K.R. and Remick, D.G. (2000):

Interleukin-8 levels and activity in delayed-healing human thermal wounds. *Wound Repair Regen*; 8(3): 216–225.

Jacob, S.E.; Berman, B.; Nassiri, M. and Vincek, V. (2003):

Topical application of imiquimod 5% cream to keloids alters expression genes associated with apoptosis. *Br J Dermatol*; 149 Suppl 66:62-55.

Jeschke, M.G.; Herndon, D.N. and Ebener, C. (2001):

Nutritional intervention high in vitamins, protein, amino acids, and omega3 fatty acids improves protein metabolism during the hypermetabolic state after thermal injury. *Arch Surg*; 136(11):1301–1306.

Jorgensen, L.N.; Kallehave, F. and Christensen, E. (1998):

Less collagen production in smokers. *Surgery*; 123(4):450–455.

Kamamoto, F.; Paggiaro, A.O.; Rodas, A.; Herson, M.R. and Mathor, M.B. (2003):

A wound contraction experimental model for studying keloids and wound-healing modulators. *Artif Organs*; 27(8):701-705.

Kelly, A.P. (2004):

Medical and surgical therapies for keloids. *Dermatol Ther*; 17(2):212-218.

Kim, D.Y.; Kim, E.S.; Kim, K.S.; Lee, S.Y. and Cho BH. (2004):

A surgical approach for earlobe keloid: keloid fillet flap. *Plast Reconstr Surg*; 113(6):1668-1674.

Kossi, J.; Vaha-Kreula, M.; Peltonen, J.; Risteli, J. and Laato, M. (2004):

Hexose sugars differentially alter collagen gene expression and synthesis in fibroblasts derived from granulation tissue, hypertrophic scar and keloid. *Arch Dermatol Res*; 295(12):521-526.

Kumar, S.; Singh, R.J.; Reed, A.M. and Lteif, A.N. (2004):

Cushing's syndrome after intra-articular and intradermal administration of triamcinolone acetonide in three pediatric patients. *Pediatrics*; 113(6):1820-1824.

Kuo, Y.R.; Jeng, S.F.; Wang, F.S.; Chen, T.H.; Huang, H.C.; Chang, P.R. and Yang, K.D. (2004):

Flashlamp pulsed dye laser (PDL) suppression of keloid proliferation through down-regulation of TGF-beta1 expression and extracellular matrix expression. *Lasers Surg Med*; 34(2):104-108.

Kwiatkowski, T.C.; Hanley, E.N. and Ramp, W.K. (1996):

Cigarette smoking and its orthopedic consequences. *Am J Ortho*; 25(9):590– 597.

Larrabee, W.F.; East, C.A.; Jaffe, H.S. (1990):

Intralesional interferon gamma treatment for keloids and hypertrophic scars. Arch Otolaryngol Head Neck Surg 1990; 116:1159– 1161.

Lawrence, W.T. (1998):

Physiology of the acute wound. Clin Plas Surg; 25(3):321- 340.

Leake, D.; Doerr, T.D. and Scott, G. (2003):

Expression of urokinase-type plasminogen activator and its receptor in keloids. Arch Otolaryngol Head Neck Surg; 129(12):1334-1338.

Lee, T.Y., Chin, G.S., and Kim, W.J. (1999):

Expression of transforming growth factor beta 1, 2, and 3 proteins in keloids. Ann Plast Surg; 43(2):179– 184.

Lee, J.Y.; Yang, C.C.; Chao, S.C. and Wong, T.W. (2004):

Histopathological Differential Diagnosis of Keloid and Hypertrophic Scar. Am J Dermatopathol; 26(5):379-384.

Lim, I.J.; Phan, T.T. and Song, C. (2001):

Investigation of the influence of keloid-derived keratinocytes on fibroblast growth and proliferation in vitro. Plast Reconstr Surg;107(3):797– 808.

Linares, H.A.; Larson, D.L. and Galstaun, B.A. (1993):

Historical notes on the use of pressure in the treatment of hypertrophic scars or keloids. Burns; 19: 17–21.

Liu, W.; Wang, D.R. and Cao, Y.L. (2004):

TGF-beta: a fibrotic factor in wound scarring and a potential target for anti-scarring gene therapy. *Curr Gene Ther*; 4(1):123-136.

Ma, H.; Xu, R.; Cheng, H.; Kuo, H.S.; During, M. and Fang, R.H. (2004):

Gene transfer into human keloid tissue with adeno-associated virus vector. *J Trauma*; 54(3):569-573.

Malaker, K.; Vijayraghavan, K.; Hodson, I. and Al Yafi, T. (2004):

Retrospective analysis of treatment of unresectable keloids with primary radiation over 25 years. *Clin Oncol*; 16(4):290-298.

Malaker, K.; Zaidi, M. and Franka, M.R. (2004) [2]:

Treatment of earlobe keloids using the cobalt 60 teletherapy unit. *Ann Plast Surg*; 52(6):602-604.

Mancoll, J.S., and Phillips, L.G. (2003):

Delayed wound healing. In: *Plastic Surgery: Indications, Operations and Outcome*; 1st edition; (ed.) by ***Ashour, B.M. and Eriksson, E.*** St. Louis, London, Philadelphia: Mosby. PP: 65- 79

Marneros, A.G.; Norris, J.E., Watanabe, S.; Reichenberger, E. and Olsen, B.R. (2004):

Genome scans provide evidence for keloid susceptibility loci on chromosomes 2q23 and 7p11. *J Invest Dermatol*; 122(5):1126-1132.

Mast, B.A., and Cohen, I.K. (2000):

Normal wound healing. In: In: Plastic Surgery: Indications, Operations and Outcome; 1st edition; (ed.) by ***Ashour, B.M. and Eriksson, E.*** St. Louis, London, Philadelphia: Mosby. PP: 37- 53

Monaco, J.L. and Lawrence, W.T. (2003):

Acute wound healing: An overview. Clin Plastic Surg; 30(1): 1– 12.

Muzaffar, A.R.; Rafols, F.; Masson, J.; Ezaki, M. and Carter, P. (2004):

Keloid formation after syndactyly reconstruction: associated conditions, prevalence, and preliminary report of a treatment method. J Hand Surg [Am]; 29(2):201-208.

Nanda, S. and Reddy, B. (2004):

Intralesional 5-fluorouracil as a treatment modality of keloids. Dermatol Surg; 30(1):54-6; discussion 56-57.

Nedelec, B.; Shankowsky, H. and Scott, P. (2001):

Myofibroblasts and apoptosis in human hypertrophic scars: the effect of interferon- α 2b. Surgery; 130(5): 798– 808.

Niessen FB, Andriessen MP, and Schalkwijk J. (2001):

Keratinocyte-derived growth factors play a role in the formation of hypertrophic scars. J Pathol; 194(2):207– 216.

Ogawa, R.; Mitsuhashi, K.; Hyakusoku, H. and Miyashita, J. (2003):

Postoperative electron-beam irradiation therapy for keloids and hypertrophic scars: retrospective study of 147 cases followed for more than 18 months. Plast Reconstr Surg; 111(2):547-53; discussion 554-555.

Osswald, S.S.; Elston, D.M. and Vogel, P.S. (2003):

Giant right plantar keloid treated with excision and tissue-engineered allograft. J Am Acad Dermatol; 48(1):131-134.

Parsons, S.L.; Watson, S.A. and Brown, P.D. (1997):

Matrix metalloproteinases. Br J Surg; 84:160– 166.

Perkins, K.; Davey, R.B. and Wallis, K.A. (1983):

Silicone gel: a new treatment for burn scars and contractures. Burns; 9:201–204.

Phan, T.T.; See, P.; Tran, E.; Nguyen, T.T.; Chan, S.Y.; Lee, S.T. and Huynh, H. (2003):

Suppression of insulin-like growth factor signalling pathway and collagen expression in keloid-derived fibroblasts by quercetin: its therapeutic potential use in the treatment and/or prevention of keloids. Br J Dermatol; 148(3):544-552.

Poochareon, V.N. and Berman, B. (2003):

New therapies for the management of keloids. J Craniofac Surg; 14(5):654-657.

Quirinia, A. and Viidik, A. (1997):

Diclofenac and indomethacin influence the healing of normal and ischemic incisional wounds in skin. Scand J Plast Recon Surg Hand Surg; 31(3):213– 219.

Ragoowansi, R.; Cornes, P.G.; Moss, A.L. and Glees, J.P. (2003):

Treatment of keloids by surgical excision and immediate postoperative single-fraction radiotherapy. Plast Reconstr Surg; 111(6):1853-1859.

Rahban, S.R. and Garner, W.L. (2003):

Fibroproliferative scars. Clin Plastic Surg 30: 77– 89

Riley, G.P.; Cox, M. and Harrall, R.L. (2001):

Inhibition of tendon cell proliferation and matrix glycosaminoglycan synthesis by non-steroidal anti-inflammatory drugs in vitro. J Hand Surg (Br); 26(3):224–228.

Robson, M.C. (2003):

Fibroproliferative scarring. Surg Clin N Am, 83: 557-569.

Rohrich, R.J.; Coberly, D.M. and Krueger, J.K. (2002):

Planning elective operations on patients who smoke: survey of North American plastic surgeons. Plast Recon Surg; 109(1):350– 355.

Sayah, D.N.; Soo, C. and Shaw, W.W. (1999):

Downregulation of apoptosis-related genes in keloid tissues. J Surg Res; 87(2):209- 216.

Senel, O.; Cetinkale, O. and Ozbay, G. (1997):

Oxygen free radicals impair wound healing in ischemic rat skin. Ann Plast Surg; 39(5):516– 523.

Sharquie, K.E. and Al-Dhalimi, M.A. (2003):

Keloid in Iraqi patients: a clinicohistopathologic study. Dermatol Surg; 29(8):847- 851.

Sherris, D.A.; Labbabee Jr, W.F. and Murakami, C.S. (1995):

Management of scar contractures, hypertrophic scars, and keloids. *Otolaryngol Clin North Am*; 28:1057–1068.

Shi, H.P.; Efron, D.T. and Most, D. (2002):

Supplemental dietary arginine enhances wound healing in normal but not inducible nitric oxide synthase knockout mice. *Surgery*; 128(2):374– 378.

Simman, R.; Alani, H. and Williams, F. (2003):

Effect of mitomycin C on keloid fibroblasts: an in vitro study. *Ann Plast Surg*; 50(1):71-76.

Sinnatamby, C.S. (1999):

Introduction to regional anatomy. In *Last's anatomy: regional and applied*. 10th edition. London: Churchill Livingstone. p 1- 35.

Stadelmann, W.K.; Digenis, A.G. and Tobin, G.R. (1998):

Impediments to wound healing. *Am J Surg*; (Suppl 2A): 39–47.

Tenaud, I.; Sainte-Marie, I. and Jumbou, O. (1999):

In vitro modulation of keratinocyte wound healing integrins by zinc, copper and manganese. *Br J Dermatol*; 140(1):26– 34.

Tredget, E.E.; Nedelec, B., and Scott, P.G. (1997):

Hypertrophic scars, keloids, and contractures. The cellular and molecular basis for therapy. *Surg Clin North Am*; 77(3):701– 730.

Tredget, E.E.; Shankowsky, H.A. and Pannu, R. (1998):

Transforming growth factor- β in thermally injured patients with hypertrophic Scars: effect of interferon α -2b. *Plast Reconstr surg*, 102: 1317-1330.

Tredget, E.E.; Wang, R. and Shen, Q. (2000):

Transforming growth factor-beta mRNA and protein in hypertrophic scar tissues and fibroblasts: antagonism by IFN-alpha and IFN-gamma in vitro and in vivo. *J Interferon Cytokine Res*; 20(2):143– 151

Uchida, G.; Yoshimura, K.; Kitano, Y.; Okazaki, M. and Harii, K. (2003):

Tretinoin reverses upregulation of matrix metalloproteinase-13 in human keloid-derived fibroblasts. *Exp Dermatol*; 12 Suppl 2:35-42.

Wendling, J.; Marchand, A.; Mauviel, A. and Verrecchia, F. (2003):

5-fluorouracil blocks transforming growth factor-beta-induced alpha 2 type I collagen gene (COL1A2) expression in human fibroblasts via c-Jun NH2-terminal kinase/activator protein-1 activation. *Mol Pharmacol*; 64(3):707-713.

Wenk, J.; Foitzik, A. and Achterberg, V. (2001):

Selective pickup of increased iron by deferoxamine-coupled cellulose abrogates the iron-driven induction of matrix-degrading metalloproteinase 1 and lipid peroxidation in human dermal fibroblasts in vitro: a new dressing concept. *J Invest Dermatol*; 116(6):833– 839.

Witte, M.B., and Barbul, A. (1997):

Wound healing: general principles of wound healing. *Surg Clin N Am*; 77:509– 528.