

REFERENCES

«References»

Aguwa,C.N,(1986):

Endocrine diseases . In : Therapeutic basis of clinical pharmacy
in the tropics. P 200-203.

Alexandrova, M. and Sloff, M.S. (1980):

oxytocin receptors and parturition. In. control of oxytocin
receptor concentration in the rat myometrium at term.
Endocrinology, 106:730-5.

Al Gamal,A.M.A.(1999):

Effect of fetal membranes and calcium channels on uterine
contractility in normal and excited conditions.
Zagazig, Benha University ; Physiology department, thesis
MD(7).

Ann,W.; Kristing,E. and Casey,M.L.(1992):

Contractile effects of prostaglandins, oxytocin and endothelein 1
in human myometrium in vitro: Refractoriness of myometrial
tissue of pregnant women to prostaglandins E₂ and F_{2α} .
J Clin Endocrinol and Metabol 75:1027-1032.

Anwer,K. and Sanborn,B.M.(1989):

Changes in intracellular free calcium in isolated myometrial cells
; role of extracellulr and intracellular calcium and possible
involvement of guanine nucleotide-sensitive proteins .
Endocrinolog ; 124:17.

Atkinson, M.A. and Maclaren, N.K. (1994):

Mechanisms of disease : The pathogenesis of insulin dependent diabetes mellitus .

N.Engl.J Med., 331:1428 .

Aviado, D.M. (1980) :

Ergot Alkaloids used during the postpartum period. In: Pharmaceutical principles of medical practice .Ed .Williams & Wilkins ,Blatimore 703-717.

Barker, L.; Egler, J.M.; Klein, S.H.; et al . (1981):

Meticulous control of diabetes during organogenesis prevents congenital lumbosacral defects in rats .

Diabetes., 30:955 .

Baron, A.D.; Johnson, A.; Fineberg, N.; Brechtel, G. and

Steinberg, H. (1994):

Skeletal muscle blood flow independently modulates insulin mediated glucose uptake .

J Clin Invest 266: E248:E253 .

Bennett, A.; Murray, J. and Wyllie, J. (1988):

PGs and gastric secretion in man. Br J Pharma ,32:339 .

Bergstöm, S. (1967): Prostaglandins :

members of a new hormonal system : these physiologically very potent compounds of ubiquitous occurrence are formed essential fatty acid . Science 157:382-391.

Bergstörn,S. and Samuelsson,B. (1962):

Isolation of PGE₁ from human seminal plasma .
J Biolol Chem .237:PC :3005 .

Bergstörn,S. and Sjovall,J.(1957):

The isolation of prostaglandin .
Acta Chemical Scandinavica,11,1086.

Bergstörn,S.; Carlson,L.A. and Weeks,J.R.(1968):

The prostaglandins : a family of biologically active lipids .
Pharmacological Rev., 20:1-48 .

Bergstörn,S.; Danielsson,H. and Samuelsson,B. (1964):

The enzymatic formation of PGE₂ from arachidonic acid .
Biochemica et Biophysica Acta .90; 207-210.

Bergstörn,S.; Ryhage,R.; Samuelsson,B. and Sjovall,J.(1962):

The structure of PGE₁ and F_{2α} .
Acta Chem Scand .16:501 .

Berne and Christian;(2005):

Type 1 diabetes is associated with increased cyclooxygenase and
cytokine mediated inflammation .
Diabetes Care.

Berry,J.L.; and Gabbe,S.G.(1986):

Diabetes mellitus in pregnancy .In : High risk pregnancy
;Knuppel,R.A.and Drukker,J.E. P.400-401.

Boreth,N.; Dey,N. and Lincoln,T.(1997):

cGMP and GMP dependent protein kinase regulates vascular smooth muscle cell phenotype .

J vas Res. 34:245-259 .

Brownlee,M. and Cerami,A.(1981):

The biochemistry of the complications of diabetes .

Annu Rev Biochem,50: 385-432.

Bygdeman,M.(1984):

The use of prostaglandins and their analogues for abortion .

Clin Obstet Gynecol 11:573-584.

Carrier and Martin,J.(2002):

Plasma F₂ isoprostanes :Direct evidence of increased free radical damage during acute hyperglycemia in type 2 diabetes .

Diabetes care.

Carsten, M.F. (1979):

Calcium assumption by human uterine microsomal preparations: effects of progesterone and oxytocin.

Am. J. Obstet. Gynecol. 133:598.

Carsten,M.E. and Miller,J.D. (1985):

Biochem Biophys Res Commu .130,1027-1031. .[PubMed]cited from: Nitric oxide in the contractile action of bradykinin, oxytocin and PGF_{2α} in the estrogenized rat uterus (1997).

Chaud,M.; Franchi,A.M.; McCann,S.M. and Gimeno,M.F.(1997):

Nitric oxide in the contractile action of bradykinin, oxytocin and
PGF_{2α} in the esterogenized rat uterus .

Proc Natl Acad Sci USA : 94(20):11049-11054 .

Chung,C.S. and Myrianthopoulos,N.C.(1975):

Effect of maternal diabetes on congenital malformations. In :
Bergsma, D.,ed. Factors affecting risks of congenital
malformations. New York: Stratton Intercontinental Medical
Book;23-35.

Chwalisz,K. and Garfield,R.(1994):

Role of progesterone during pregnancy models of parturition and
preeclampsia.

Geburtsch Perinatol ,198,170-180.

Claman,P.; Carpenter,R.J. and Reiter,A.(1984):

Uterine rupture with the use of vaginal PGE₂ for induction of
labor .

Am J obstet Gynecol .150:889.

Crankshaw,D.J.(1987):

The sensitivity of longitudinal and circular muscle layer of rats
myometrium to oxytocin in vitro .

Can J Physiol Pharmacol:65:778-7.

Crossland,J.(1980):

Lewis's pharmacology ,5th edition , Chuchill Livingstone ,
London, N.Y.,P.373,426,507.

Csapo, A.; Erdos,T.; Mattos, C.R.De.; Gramss, E. and Moscovitz,C.(1965):

Stretch-induced uterine growth, protein synthesis and function.
Nature (London), 207:1378-1379.

Cullen, M.B. and Harkness,R.D. (1960):

The effect of hormones on the physical properties and collagen
content of the rats uterine cervix.
J. Physiol. (Lond.) 152:419-436.

Cunningham, F.G.; McDonald, P.C. and Grant,N.F. (1989):

Maternal adaptation of pregnancy and diabetes mellitus
complicating pregnancy. In Williams Obstetrics. Appleton and
Lang. (ed.).

Cunningham,F.G.; MacDonald,P.G.; Gant,N.F.and Gray,D.v.(1997):

Normal labor and delivery . In: Williams Obstetrics.20th (ed) vol
1. Appleton and Lang P:227.

Dale,H.H.(1906):

On some physical actions of ergot.
J physiol ., London., 34,163.

Dawood, M.Y. (1977):

Neurohypophyseal hormones. In: Fuchs, F.; Klopper, A. eds.
Endocrinology of pregnancy. Philadelphia Harper & Raw,
204:28.