

References

References

Acarregui, M. J. K., Ramirez, K. R. and Brown, R. K. : Vascular endothelial growth factor (VEGF) induces airway epithelial cell proliferation and surfactant protein gene expression in human fetal lung in vitro. *Pediatr. Res.* 43: 44, 1998.

Adini, A., Kornaga, T. and Firoozbakht, F. : Placental growth factor is a survival factor for tumour endothelial cells and macrophages. *Cancer Res.* 62:2749,2002.

Akeson, AL., Greenberg, J.M. and Cameron, J.E. : Temporal and spatial regulation of VEGF-A controls vascular patterning in the embryonic lung. *Dev Biol.* 264: 443, 2003.

Anil, N., Praveen, K. and Sourabh, D. : Surfactant Therapy for Hyaline Membrane Disease *Indian Pediatrics.* 38:640,2001.

Askie, L.M. : The effect of differing oxygen saturation targeting ranges on long term growth and development of extremely preterm, oxygen dependent infants. *Pediatr Res .* 51:378A,2002.

Attar. M.A. and Donn. S.M.: Mechanisms of ventilator-induced lung injury in premature infants. *Semin. Neonatol.* 7(5):353,2002.

Azzopardi, D. : Pilot study of treatment with whole body hypothermia for neonatal encephalopathy. *Pediatrics .* 106:684,2000.

Bancalari, E. : Mechanical ventilation. *Effective Care of the Newborn Infant.* Oxford University Press. 200, 1992.

Bates. D.O., Cui, and T.G. Doughty, J.M. : VEGF165b, an inhibitory splice variant of vascular endothelial growth factor, is down-regulated in renal cell carcinoma. *Cancer. Res.* 62:4123,2002.

Becker, P.M., Verin, A.D. and Booth, M.A. : Differential regulation of diverse physiological responses to VEGF in pulmonary endothelial cells. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 281:L1500,2001.

Becker, P.M., Waltenberger, J. and Yachechko, R. : Neuropilin-1 regulates vascular endothelial growth factor-mediated endothelial permeability. *Circ. Res.* 96:1257,2005.

Bernatchez, P.N., Winstead, M.V. and Dennis, E.A : VEGF stimulation of endothelial cell PAF synthesis is mediated by group V 14 kDa secretory phospholipase A2. *Br. J. Pharmacol.* 134, 197,2001 .

Bevilacqua, G., Parmigiani, S. and Robertson, B. : Prophylaxis of respiratory distress syndrome by treatment with modified porcine surfactant at birth: a multicentre prospective randomized trial. *J. Perinat. Med.* 24:1,1996.

Bhatt, A.J., Pryhuber, G.S. and Huyck. H. : Disrupted pulmonary vasculature and decreased vascular endothelial growth factor, Flt-1, and TIE-2 in human infants dying with bronchopulmonary dysplasia. *Am. J. Crit. Care. Med.* 164: 1971, 2001.

Bhuta. T. and Henderson-Smart. DJ.: Rescue high frequency oscillatory ventilation versus conventional ventilation for pulmonary dysfunction in preterm infants. *Syst. Rev.* (2): 000438,2000.

Bhutta. Z.A. and Yusuf. K. : Profile and outcome of the respiratory distress syndrome among newborns in Karachi, risk factors for mortality. *J. Trop. Pediatr.* 43(3):143,1997.

Bland, R. and Coalson, J.J. : Chronic lung disease in early infancy. New York, Marcel. Dekker. Inc. 2000.

Bloom, R. : A consideration of neonatal resuscitation. *Pediatr. Clin. North. Am.* 51:669,2004.

Bober, K. and Swietliński, J. : Diagnostic utility of ultrasonography for respiratory distress syndrome in neonates. *Med. Sci. Monit.* 12(10): 440, 2006.

Braems, G.A., Han, V.K. and Challis, J.R. : Gestational age-dependent changes in the levels of mRNAs encoding cortisol biosynthetic enzymes and IGF-II in the adrenal gland of fetal sheep during prolonged hypoxemia. *J. Endocrinol.* 159:257,1996

Brus, F., Van Oeveren, W. and Okken, A. : Number and activation of circulating polymorphonuclear leukocytes and platelets are associated with neonatal respiratory distress syndrome severity . *Pediatrics.* 99(5):672,1997.

Carlo, W.A. : Minimal ventilation to prevent bronchopulmonary dysplasia in extremely-low-birth-weight infants. *J. Pediatr.* 141:3705,2002.

Carmeliet, P., Ng, Y.S. and Nuyens, D. : Impaired myocardial angiogenesis and ischemic cardiomyopathy in mice lacking the vascular endothelial growth factor isoforms VEGF164 and VEGF188. *Nat. Med.* 5: 495, 1999.

Caukwell, S. and Murphy, D.J. : The effect of mode of delivery and gestational age on neonatal outcome of the non-cephalic- presenting second twin. *Am. J. Obstet. Gynecol.* 187:1356,2002.

Chatburn, R.L. : A new system for understanding mechanical ventilators. *Respir. Care.* 36:1123,1991.

Cho, K., Matsuda, T. and Okajima, S. : Prediction of respiratory distress syndrome by the level of pulmonary surfactant protein A in cord blood sera. *Biol. Neonate.* 77:83,2000.

Choe, K.H., Taraseviciene-Stewart, L. and Scerbavicius, R. : Methylprednisolone causes matrix metalloproteinase-dependent emphysema in adult rats. *Am. J. Respir. Crit. Care. Med.* 167: 1516,2004.

Clerch, L.B., Baras, A.S. and Massaro, G.D. : DNA microarray analysis of neonatal mouse lung connects regulation of KDR with dexamethasone-induced inhibition of alveolar formation. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 286: L411,2003.

Compernelle, V., Brusselmans, K. and Acker, T. P. : Loss of HIF-2 α and inhibition of VEGF impair fetal lung maturation, whereas treatment with VEGF prevents fatal respiratory distress in premature mice. *Nat. Med.* 8: 702, 2002.

Corne, J., Chupp, G. and Lee, C.G. : IL-13 stimulates vascular endothelial cell growth factor and protects against hyperoxic acute lung injury. *J. Clin. Invest.* 106:783,2000.

Cossar, D., Bell, J. and Lang, M. : Development of human fetal lung in organ culture compared with in utero ontogeny. *In Vitro Cellular and Developmental Biology.* 29A, 319,1993.

Crouch, E. and Wright, J.R. : Surfactant proteins a and d and pulmonary host defense. *Annu. Rev. Physiol.* 63:521,2001.

D'Angio, C. , Ryan, R. M. and Maniscalco, W.M. : Vascular endothelial growth factor in pulmonary lavage fluid from premature infants : Effects of age and postnatal dexamethasone. *Biol. Neonate.* 76:273, 1999.

Demissie, K., Ananth, C.V. and Martin, J. : Fetal and neonatal mortality among twin gestations in the United States: the role of intrapair birth weight discordance. *Obstet. Gynecol.* 100:474,1998.

Donn, S.M. : Newer techniques of mechanical ventilation. *Semin. Neonatol.* 7:401,2002.

Donn, S.M. : Rapid detection of neonatal intracranial hemorrhage by transillumination. *Pediatrics.* 64:843, 1979.

Donovan, E.F., Ehrenkranz, R.A. and Shankaran, S. : Outcomes of very low birth weight twins cared for in the National Institute of Child Health and Human Development Neonatal Research Network's intensive care units. *Am. J. Obstet. Gynecol.* 179:742 ,2001.

Downes, J. : Respiratory distress syndrome of newborn infants. I. New clinical scoring system with acid-base and blood-gas correlation. *Clin. Pediatr.* 9:325, 1970.

Edelman, J.L., Lutz, D. and Castro, M.R. : Corticosteroids inhibit VEGF induced vascular leakage in a rabbit model of blood-retinal and bloodaqueous barrier breakdown. *Exp. Eye. Res.* 80: 249 ,2005.

Engle, W.A. : Surfactant-replacement therapy for respiratory distress in the preterm and term neonate. *Pediatrics.* 121:32,2008.

Ferrara, N., and Henzel, W.J. : Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cell. *Biochem. Biophys. Res. Commun.* 161: 851 ,1989.

Fiori, H.H., Linderholm, B. and Fiori, R.M : Computerized image analysis of bubbles in gastric aspirate for prediction of respiratory distress syndrome. *Acta. Paediatr.* 90:1402 ,2000.

Foster, R.R., Hole, R. and Anderson, K. : Functional evidence that vascular endothelial growth factor may act as an autocrine factor on human podocytes. *Am. J. Physiol. Renal. Physiol.* 284:F1263 ,2003.

Friedman, S.A., Schiff, E. and Lubarsky, S.L. : Expectant management of severe preeclampsia remote from term. *Clin. Obstet. Gynecol.* 42:470,1999.

Furman, B., Shoham-Vardi, I. and Bashiri, A. : Preterm premature rupture of membranes is not an independent risk factor for neonatal morbidity. *J. Matern. Fetal. Med.* 10:107,2000.

Gebb, S.A. and Shannon, J.M. : Tissue interactions mediate early events in pulmonary vasculogenesis. *Dev. Dyn.* 217: 15 ,2000.

Gerber, H.P., Dixit, V. and Ferrara, N. : Vascular endothelial growth factor induces expression of the antiapoptotic proteins Bcl-2 and A1 in vascular endothelial cells. *J. Biol. Chem.* 273:13313 ,1999.

Gerber, H.P., Hillan, K.J. and Ryan, A.M. : VEGF is required for growth and survival in neonatal mice. *Development.* 126: 1149 ,2001.

Gissler, M., Vuori, E.R. and Ritvanen, A. : Finnish Reproduction Statistics 2000. National Research and Development Centre for Welfare and Health. 1–136,2002.

Goldenberg, R.L., Hauth, J.C. and Andrews, W.W. : Intrauterine infection and preterm delivery. *N. Engl. J. Med.* 342:1500,2000.

Greenough, A., Morley, G.J. and Robertson, N.R.C. : Acute respiratory disease in the newborn. Respiratory distress syndrome in : *Textbook of Neonatology.* Edinburgh . 385-486,1996.

Greenough, A. and Donn, S.M. : Matching ventilatory support strategies to respiratory pathophysiology. *Clin. Perinatol.* Mar;34(1):35, 2007.

Gregory, G.A. : Treatment of the idiopathic respiratory distress syndrome with continuous positive airway pressure. N. Engl. J. Med. 284:1333,1971.

Grier, D.G. and Halliday, H.L. : Effects of glucocorticoids on fetal and neonatal lung development. Treat . REspir. Med. 3(5):295 ,2002.

Grover, T.R., Parker, T.A. and Zenge, J.P. : Intrauterine hypertension decreases lung VEGF expression and VEGF inhibition causes pulmonary hypertension in the ovine foetus. Am. J. Physiol. Lung. Cell. Mol. Physiol. 284:L508 ,2003.

Hacking, D., Watkins, A. and Fraser, S. : Respiratory distress syndrome and birth order in premature twins. Arch. Dis. Child. Fetal. Neonatal .Ed. 84:F117 ,2002.

Halliday, H.L. : Hyaline membrane syndrome disease. Campbell. AGM. and McIntoch. N, 133, 1992.

Hallman, M. : Cytokines, pulmonary surfactant and consequences of intrauterine infection. Biol. Neonate. 1:2 ,1999.

Hallman, M. : Exogenous human surfactant for treatment of severe respiratory distress syndrome: A randomized prospective clinical trial. Pediatr . 106:963,1985.

Hansbrough, J.R., Fine, S.M. and Gordon, J I. : A transgenic mouse model for studying the lineage relationships and differentiation program of type II pneumocytes at various stages of lung development. Journal of Biological Chemistry. 268,9762 ,1993.

Hansen Tand Corbet, A. : Disorders of the transition , Hyaline membrane disease. Avery,s Diseases of the Newborn. Philadelphia. pp609,1998.

Hanson, L. : The transfer of immunity from mother to child. Ann. N. Y. Acad. Sci. 987:199,2003.

Hawgood, S. and Clements, J.A. : Pulmonary surfactant and its apoproteins. J. Clin. Invest. 86:1 ,1990.

Healy, A.M., Morgenthau, L. and Zhu, X. : VEGF is deposited in the subepithelial matrix at the leading edge of branching airways and stimulates neovascularization in the murine embryonic lung. Dev. Dyn. 219: 341 ,2000.

Helbich, T.H., Popow, C. and Dobner, M. : New-born infants with severe hyaline membrane disease. Radiological evaluation during high frequency oscillatory versus conventional ventilation. Eur. J. Radiol. Oct . 28(3):243 ,1998.

Hiratsuka, S., Minowa, O. and Kuno, J. : Flt-1 lacking the tyrosine kinase domain is sufficient for normal development and angiogenesis in mice. Proc. Natl. Acad. Sci. USA . 95:9349 ,2002.

Hislop, A. and Reid, L. : Development of the acinus in the human lung. Thorax. 29, 90 ,1974.

Hitti, J., Tarczy-Hornoch, P. and Murphy, J. : Amniotic fluid infection, cytokines, and adverse outcome among infants at 34 weeks' gestation or less. Obstet. Gynecol. 98:1080 ,2001.

Horbar, J.D., Badger, G.J. and Carpenter, J.H. : Trends in mortality and morbidity for very low birth weight infants . Pediatrics. 110:143 ,2002.

Hosford, G.E. and Olson, D.M. : Effects of hyperoxia on VEGF, its receptors, and HIF-2_α in the newborn rat lung. Am. J. Physiol. Lung. Cell. Mol. Physiol. 285: L161 ,2003.

Ingemarrson, I. : Gender aspects of preterm birth. BJOG . 110(suppl 20):34 , 2003.

Jaarsma, A.S. : Activation of the inflammatory reaction within minutes after birth in ventilated preterm lambs with neonatal respiratory distress syndrome. Biol. Neonate. 86:1,2004.

Jakkula, M., Le Cras, T.D. and Gebb, S. : Inhibition of angiogenesis decreases alveolarization in the developing rat lung. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 279: L600 ,2000.

Jakkula, M., Le Cras, T.D. and Gebb, S. : Inhibition of angiogenesis decreases alveolarization in the developing rat lung. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 279: L600 ,2002.

Jenkinson. S.G. : Oxygen toxicity. *New. Horiz.* (4):504,1993.

Jobe, A. : Phospholipid metabolism and turnover. Fetal and neonatal physiology. Philadelphia . 986,1992.

Kaneko, K., Shimizu, H. and Arakawa, H. : Pulmonary surfactant protein A in sera for assessing neonatal lung maturation. *Early. Hum. Dev.* 62:11 ,2001.

Kaner, R.J., Ladetto, J.V. and Singh, R. : Lung overexpression of the vascular endothelial growth factor gene induces pulmonary oedema. *Am. J. Respir. Cell . Mol . Biol .* 22:657 ,2000.

Kaner, R.J. and Crystal, R.G. : Compartmentalization of vascular endothelial growth factor to the epithelial surface of the human lung. *Mol. Med.* 7:240, 2001.

Karmaliotis, D., Kosmidou, I. and Ingenito, EP. : Angiogenic growth factors in the pathophysiology of a murine model of acute lung injury. *Am. J. Physiol. Lung. Cell. Mol. Physiol .* 283:L585 ,2002.

Khoury, M.J., Marks, J.S. and McCarthy, B.J. : Factors affecting the sex differential in neonatal mortality . the role of respiratory distress syndrome. *Am. J. Obstet. Gynecol.* 151:777 ,2004.

King, J. and Flenady, V. : Prophylactic antibiotics for inhibiting preterm labour with intact membranes. *Cochrane. Database. Syst. Rev.* 2002.

Kirkpatrick, B.V. and Mueller, D.C. : Respiratory distress syndrome. Disorders of Respiratory Tract in Children . Philadelphia . pp332,1998.

Kitsukawa, T., Shimizu, M. and Sanbo, M. : Neuropilin-semaphorin III/D-mediated chemorepulsive signals play a crucial role in peripheral nerve projection in mice. Neuron . 19:995 ,1997.

Kleigman, R.M. : The fetes and neonatal infants, the high risk infant, prematurity and intrauterine growth retardation. In Nelson Textetbook of pediatrics. 7th Edetion. P:556, 2004.

Klekamp, J.G., Jarzecka, K. and Perkett, E.A. : Exposure to hyperoxia decreases the expression of vascular endothelial growth factor and its receptors in adult rat lungs. Am. J. Pathol. 154: 823 ,1999.

Koivisto, M., Marttila, R. and Saarela, T. : Wheezing illness and re-hospitalization in the first two years of life after neonatal respiratory distress syndrome. J. Pediatr. 147:486 ,2005.

Koivisto, M., Marttila , R. and Kurkinen-Raty , M. : Changing incidence and outcome of infants with respiratory distres syndrome ,a population-based survey. Acta. Paediatr. 93: 184, 2004.

Kotecha, S. : Lung growth: Implications for the newborn infant. Arch. Dis. Child. Fetal. Neonatal. Ed. 82:F69–74, 2000.

Koyama, S., Sato, E. and Tsukadaira, A. : Vascular endothelial growth factor mRNA and protein expression in airway epithelial cell lines in vitro. Eur. Respir. J .20:1449 ,2002.

Lantuejoul, S., Constantin, B. and Drabkin, H. : Expression of VEGF, semaphorin SEMA3F, and their common receptors neuropilins NP1 and NP2 in

pre-invasive bronchial lesions, lung tumours, and cell lines. J. Pathol . 200:336 ,2003.

Lassus, P., Ristimäki, A. and Ylikorkala, O. : Vascular endothelial growth factor in human preterm lung. Am. J. Respir. Crit. Care. Med. 159: 1429 , 1999.

Lassus, P., Turanlahti, M. and Heikkilä. : Pulmonary vascular endothelial growth factor and Flt-1 in fetuses, in acute and chronic lung disease, and in persistent pulmonary hypertension of the newborn. Am. J. Respir. Crit .Care. Med. 164, 2001.

Le Cras, T.D., Markham, N.E. and Tuder, R.M. : Treatment of newborn rats with a VEGF receptor inhibitor causes pulmonary hypertension and abnormal lung structure. Am. J. Physiol. Lung. Cell. Mol. Physiol . 283:L555 ,2002.

Le Cras, T.D., Spitzmuller, R.E. and Albertine, K.H. : VEGF causes pulmonary haemorrhage, hemosiderosis, and air space enlargement in neonatal mice. Am. J. Physiol. Lung. Cell. Mol. Physiol . 287:L134 ,2004.

Ley, D., Wide-Svensson, D. and Lindroth, M. : Respiratory distress syndrome in infants with impaired intrauterine growth. Acta. Paediatr. 86:1090 ,1997.

Lin, C., Wang, S. and Hsu, Y. : Risk for respiratory distress syndrome in preterm infants born to mothers complicated by placenta previa. Early. Hum. Dev;60(3):215-24,2001.

Locke, R.G. : Inadvertent administration of positive end-expiratory pressure during nasal cannula flow. Pediatrics. 91:135,1993.

Luttun, A. and Carmeliet, P. : Angiogenesis and lymphangiogenesis: highlights of the past year. Curr. Opin. Hematol. 11: 262 ,2004.

Maeda, S., Suzuki, S. and Suzuki, T. : Analysis of intrapulmonary vessels and epithelialendothelial interactions in the human developing lung. *Lab. Invest.* 82: 293 ,2002.

Malloy, M.H. and Freeman, D.H. : Respiratory distress syndrome mortality in the United States, 1987 to 1995. *J. Perinatol.* 20:414 ,2000.

Maniscalco, W.M., Watkins, R.H. and Pryhuber, G.S. : Angiogenic factors and alveolar vasculature: development and alterations by injury in very premature baboons. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 282: L811, 2002.

Martin, J. A., Kochanek, K. D. and Strobino, D. M . : Annual summary of vital statistics. *Pediatrics.* 115: 634, 2005.

Matsumoto, T. and Claesson-Welsh, L. : VEGF receptor signal transduction. *Sci.* 112:RE21,2001.

McIntire, D.D., Bloom, S.L. and Casey, B.M. : Birth weight in relation to morbidity and mortality among newborn infants. *N. Engl. J. Med.* 340:1234 , 1999.

Mendelson, C.R. and Boggaram, V. : Hormonal control of the surfactant system in fetal lung. *Annual Review of Physiology.* 53,415 ,1991.

Moglia, B.B and Phelps, D.S. : Changes in surfactant protein A mRNA levels in a rat model of insulin-treated diabetic pregnancy. *Pediatr. Res.* 39:241 ,1996.

Monacci, W. T., Merrill, M. J. and Oldfield, E. H.: Expression of vascular permeability factor/vascular endothelial growth factor in normal rat tissue. *Am. J. Physiol.* 264: 1002. 1993.

Moutquin, J.M. : Classification and heterogeneity of preterm birth. *BJOG.* Suppl. 20:30 ,2003.

Narayanan, M. : Prophylactic indomethacin: Factors determining permanent ductus arteriosus closure. *Pediatrics .* 136:330,2000.

Newnham, J.P., Moss, T.J. and Nitsos, I. : Antenatal corticosteroids: the good, the bad and the unknown. *Curr. Opin. Obstet. Gynecol.* 14:607 ,2002.

Norberto, A. Freddi, and Jose Oliva Proenca Filho. : Exogenous surfactant therapy in pediatrics . *J. Pediatr.* 2: S205 ,2003.

O’Rahilly, R. : The early prenatal development of the human respiratory system. In *Pulmonary Development from Intrauterine to Extrauterine Life. Lung Biology in Health and Disease*, vol. 27, pp. 3-18.1988.

Parker, T.A. and Abman, S.H. : The pulmonary circulation in bronchopulmonary dysplasia. *Semin. Neonatol.* 8: 51 ,2003.

Pavlova, M. : Factors affecting the prognosis of Ebstein’s anomaly during fetal life. *Am. Heart. J.* 135:1081,1998.

Piazzze, J.J., Anceschi, M.M. and Maranghi, L. : Fetal lung maturity in pregnancies complicated by insulin-dependent and gestational diabetes: a matched cohort study. *Eur. J. Gynekol. Reprod. Biol.* 83(2):145 ,1999.

Piper, J.M., Xenakis, E.M. and McFarland, M. : Do growthretarded premature infants have different rates of perinatal morbidity and mortality than appropriately grown premature infants? *Obstet. Gynecol.* 87:169 ,1996.

Pochedly, C. and Musiker, S. : Twin-to-twin transfusion syndrome. *Postgrad. Med.* 47:172,1979.

Poets, C.F. : Noninvasive monitoring of oxygenation in infants and children: Practical considerations and areas of concern. *Pediatrics.* 93:737,1994.

Possmayer, F. : Pulmonary surfactant. *Can. J. Biochem. Cell. Biol.* 62:1121, 1984.

Pramanik, A. : Respiratory distress syndrome. . *Postgrad. Med.* 3:7,2004.

Randell, S.H. and Young, S.L. : Structure of alveolar epithelial cells and the surface layer during development. Fetal and neonatal physiology. Philadelphia . 962,2001.

Raoul, W., Chailley-Heu, B. and Barlier-Mur, A.M. : Effects of vascular endothelial growth factor on isolated fetal alveolar type II cells. Am. J. Physiol. Lung. Cell. Mol. Physiol. 286: L1293 ,2004.

Rayani, H.H., Gewolb, I.H. and Floros, J. : Glucose decreases steady state mRNA content of hydrophobic surfactant proteins B and C in fetal rat lung explants. Exp. Lung. Res. 25, 69 ,1999 .

Ricardo, J. R. : Management of Respiratory Distress Syndrome: An Update. Respir. Care. 48:279, 2005.

Roberton, N.R.C. and Greenough, A. : Respiratory distress syndrome. In: Neonatal Respiratory Disorders. 4th edition. London. 238-80,1997.

Roberton, N.R.C. and Greenough, A. : Respiratory distress syndrome. In: Neonatal Respiratory Disorders. 4th edition. London. 238-80,1999.

Robinson, C.J., Stringer, S.E. : The splice variants of vascular endothelial growth factor (VEGF) and their receptors. J. Cell. Sci . 114:853 ,2001.

Rodriguez, R.J., Martin, R.J. and Fanaroff, A.A. : The respiratory distress syndrome and its management, 2002.

Rooney, S. : Regulation of surfactant associated phospholipid synthesis and secretion. Fetal and neonatal physiology. Philadelphia . 986,1992.

Rooney, S.A. : Regulation of surfactant secretion. Comp. Biochem. Physiol. A. Mol. Integr. Physiol. 129: 233 ,2001.

Shah, S.S., Ohlsson, A. and Halliday, H. : Inhaled versus systemic corticosteroids for the treatment of chronic lung disease in ventilated very low birth weight preterm infants. Cochrane. Database. Syst. Rev. 2003.

Shulenin, S., Noguee, L.M. Annilo, T. : The ABCA3 gene is frequently mutated in human newborns with fatal surfactant deficiency. N. Engl. J. Med. 350:1296, 2004.

Silverman, N.H. and Schmidt, K.G. : Ventricular volume overload in the human fetus: Observations from fetal echocardiography. J. Am. Soc. Echocardiogr. 3:20, 1990.

Sims, E.J., Vermillion, S.T. and Soper, D.E. : Preterm premature rupture of the membranes is associated with a reduction in neonatal respiratory distress syndrome. Am. J. Obstet. Gynecol. 187:268, 2002.

Soker, S., Takashima, S. and Miao, H.Q. : Neuropilin-1 is expressed by endothelial and tumour cells as an isoform-specific receptor for vascular endothelial growth factor. Cell. 92:735, 1998.

Soll, R.F. and Blanco, F. : Natural surfactant extract versus synthetic surfactant for neonatal respiratory distress syndrome. Cochrane. Database. Syst. Rev. 2, 2001.

Soll, R.F. and Morley, C.J. : Prophylactic versus selective use of surfactant in preventing morbidity and mortality in preterm infants. Cochrane. Database. Syst. Rev. CD000510, 2001.

Stoll, B.J. and Kliegman, R.M. : The fetus and the neonatal infant, respiratory tract disorder. Nelson Textbook of pediatrics. 16th edition. Philadelphia. P96-510, 2004.

Tan, K.L. : The twin transfusion syndrome. Clinical observations on 35 affected

pairs. Clin. Pediatr. (Phila). 18:111,1970.

Tang, K., Rossiter, H.B. and Wagner, P.D. : Lung-targeted VEGF inactivation leads to an emphysema phenotype in mice. J. Appl. Physiol. 97:1559,2004.

Thomson, M.A. : Early nasal continuous positive airways pressure (nCPAP) with prophylactic surfactant for neonates at risk of RDS. The IFDAS Multi-Centre Randomized Trial. Pediatr. Res. 379A:2204,2002.

Tischer, E., Mitchell, R. and Hartman, T. : The human gene for vascular endothelial growth factor. Multiple protein forms are encoded through alternative exon splicing. J. Biol. Chem. 266:11947– 54,1991.

Torday, J., Carson, L. and Lawson, E.E. : Saturated phosphatidylcholine in amniotic fluid and prediction of the respiratory-distress syndrome. N. Engl. J. Med. 301:1013,1979.

Tsao, P.N., Wei, S.C. and Chou, H.C. : Vascular endothelial growth factor in preterm infants with respiratory distress syndrome. Pediatr. Pulmonol. May;39(5):461, 2005.

Vain, N.E. : Regulation of oxygen concentration delivered to infants via nasal cannulas. Am. J. Dis. Child. 143:1458,1989.

Van Golde, L.M., Batenburg, J.J. and Robertson, B. : The pulmonary surfactant system: biochemical aspects and functional significance. Physiol. Rev. 68:374,1988.

Van Overmeire, B. : A comparison of ibuprofen and indomethacin for closure of patent ductus arteriosus. N. Engl. J. Med. 343:674,2000.

Verder, H. : Surfactant therapy and nasal continuous positive airway pressure

for newborns with respiratory distress syndrome. N. Engl. J. Med. 331:1051,2004.

Vitali, S.H. and Arnold, J.H. : Ventilator strategies to reduce lung injury - lessons from pediatric and neonatal intensive care. Crit. Care.;9(2):177-83,2005.

Voelkel, Norbert, F. and William Vandivier, R. : Vascular endothelial growth factor in the lung. Am. J. Physiol. Lung. Cell. Mol. Physiol. 290: L221, 2006.

Walsh, M.C. : Safety, reliability, and validity of a physiologic definition of bronchopulmonary dysplasia. J. Perinatol. 6:451,2003.

Wax, J.R., Herson, V. and Carignan, E. : Contribution of elective delivery to severe respiratory distress at term. Am. J. Perinatol. 19:81 ,2002.

Whitsett, JA., Rice, WR. and Warner, BB. : Avery's Neonatology Pathophysiology & Management of The Newborn. 6th edition. Philadelphia. 553–577, 2005.

Widness, J.A. : Clinical performance of an in-line point-of-care monitor in neonates. Pediatrics . 106:497,2002.

Wolf, E.J., Vintzileos, A.M. and Rosenkrantz, T.S. : Do survival and morbidity of very-low-birth-weight infants vary according to the primary pregnancy complication that results in preterm delivery? Am. J. Obstet. Gynecol. 169:1233 ,2002.

Yang, R., Thomas, G.R. and Bunting, S. : Effects of vascular endothelial growth factor on hemodynamics and cardiac performance. J. Cardiovasc. Pharmacol. 27:838 ,1996.

Ziadeh, S.M. and Badria, L.F. : Effect of mode of delivery on neonatal outcome of twins with birthweight under 1500 g. Arch. Gynecol. Obstet. 264:128 ,2000.