

References

References

A:

- **Abigail, A.S. and Dixie, D.W.(1994):** Antibiotics; mechanism of action and mechanisms of resistance in bacterial pathogenesis. A molecular approach: Abigail, A.S. and Dixie, D.W. (eds); ASM press, 8th Ed: 97-110.
- **Adcock, P.M., Pastor, P., Medley, F. and Patterson, J.E. (1998):**Methicillin resistant *Staphylococcus aureus* in two child care centers ,J .Infect. Dis., 178:577-580.
- http://www.aecom.yu.edu/id/micro/msagar_saureus-72.jpg
- **Alonso, A., Martin, P., Albarran, C., Garcia, P., Garcia, O., de Simon, L.F., Garcia-Hirschfeld, J., Sancho, M., de La Rua, C. and Fernandez-Piqueras, J. (2004):** Real-time PCR designs to estimate nuclear and mitochondrial DNA copy number in forensic and ancient DNA studies. Forensic. Sci. Int., 139: 141-149.
- **Ang, J.Y., Ezike, E. and Asmar, B.I ., (2004) :** Antibacterial resistance . The Indian Journal of Pediatrics , 71 (3) : 229-239
- **Apfalter, P., Assadian, O., Kalczyk, A., Lindenmann, V., Makristathis, A., Mustafa, S., Rotter, M., and Hirschl, A.M. (2002):** Performance of a new chromogenic oxacillin resistance screen medium (Oxoid) in the detection and presumptive identification of methicillin-resistant *Staphylococcus aureus*. Diagn. Microbiol. Infect. Dis. 44:209-211.
- **Arbeit, R.D. (1999):** Laboratory procedures for the epidemiologic analysis of microorganisms. In: Murray, P.M., Baron, E.J., Tenover, F.C., Tenover, F.C., Tenover, F.C., Tenover, F.C., eds; Manual of Clinical Microbiology. Washington: ASM Press.

- **Archer, G.L. and Bosilevac, J.M. (2001):** Signaling antibiotic resistance in staphylococci. *Microbiology. Science.* 291:1915–1916.
- **Archer, G.L. and Mayhall, C.G.(1983):**Comparison of epidemiological markers used in the investigation of an outbreak of methicillin-resistant *Staphylococcus aureus* infections.*J.Clin.Microbial*,18(2):395-399.
- **Archer, G. L. and Pennell, E. (1990):** Detection of methicillin resistance in staphylococci by using a DNA probe. *Antimicrob. Agents. Chemother.*, 34:1720-1724.
- **Ayliffe, G.A.J.(1994):**Prevention of spread of MRSA. In: *Postgraduate Doctor Middel East* .17(5):219-223.
- **Ayliffe, G.A.J. (1996):** Recommendation of the control of methicillin -resistant *Staphylococcus-aureus*.WHO/EMC/LTS/9

B:

- **Baggett, H.C., Hennessy, T., Bruden, D. and Hamlin, C.(2000):** Outbreak of community –acquired Methicillin Resistant *Staphylococcus aureus*(MRSA) skin infections among Alska natives .*International Conference on Emerging Infectious Diseases*, P72.
- **Baired, D.(1996):** *Staphylococci* :Cluster forming Gram – Positive cocci. In Collee, Fraser, Marmion and Simmons Cruicshank. *Practical Medical Microbiology*,14 th ed, Churchill-Livingstone 245-250.
- **Baker, C.N., Huang, M.B. and Tenover, F.C. (1994):** Optimizing testing of methicillin- resistant staphylococcus species. *Diag. Microbiol. Infect. Dis.*, 19:167-170.
- **Balows, A., Hausler, W.G., Herrmann ,M. L., Isenberg, H.D. and Shadomy, H.J.(1991):**Manual of clinical microbiology.D5 Washington DC ,American Society of Microbiology,(17):119-126.

- **Bannerman, T.L., Hancock, G.A., Tenover, F.C., Miller, J.M.(1995):** Pulsed-field gel electrophoresis as a replacement for bacteriophage typing of *Staphylococcus aureus*. *J. Clin. Microbiol.*, 33:551-5.
- **Barbarini, D., Fumagalli, P. and Marone, P.(2001):** Methicillin-resistant *Staphylococcus aureus* (MRSA) in an intensive care unit: a one year survey. *Infez. Med.* ,9:237-45.
- **Barletta, J.M., Edelman, D.C. and Constantine, N.T. (2004):** Lowering the detection limits of HIV-1 viral load using real-time immuno-PCR for HIV-1 p24 antigen. *Am. J. Clin .Pathol*, 122, 20-27.
- **Baron, E.J. and Chang, R.S (1994):** Antimicrobial agents in *Baron Medical Microbiology: A short course*, Wiley – Liss, N.Y. 73-131.
- **Barret, S (1998):** Trying to control MRSA causes more problem than it solves. *Journal of Hospital Infection*,39(2):85-93.
- **Barski, P., Piechowicz, L., Galinski, J. and Kur, J.(1996):** Rapid assay for detection of methicillin-resistant *Staphylococcus aureus* using multiplex PCR. *Mol .Cell Probes.*,10(6):471-5.
- **Bennett, C.D., Campbell, M.N., Cook, C.J., Eyre, D.J., Nay, L.M., Nielsen, D.R., Rasmussen, R.P. and Bernard, P.S. (2003):** The LightTyper: high-throughput genotyping using fluorescent melting curve analysis. *Biotechniques*, 34, 1288-1292, 1294-1285.
- **Bernard, P.S., Ajioka, R.S., Kushner, J.P. and Wittwer, C.T. (1998):** Homogeneous multiplex genotyping of hemochromatosis mutations with fluorescent hybridization probes. *Am. J. Pathol.*, 153, 1055-1061.
- **Bhadki, S., Murbly, M., Korom, S. and Hugo, F.(1989):** Release of interleukin -1b associated with potent cytotoxic action of staphylococcal alpha –toxin on human monocytes. *Infect. Immun*, 57:3512-3519.

- **Bischoff, F.Z., Marquez-Do, D.A., Martinez, D.I., Dang, D., Horne, C., Lewis, D. and Simpson, J.L. (2003):** Intact fetal cell isolation from maternal blood: improved isolation using a simple whole blood progenitor cell enrichment approach (Rosette Sep). *Clin. Genet*, 63, 483-489.
- **Blake, J.E. and Metcalfe, M.A.(2001):** A shared noncapsular antigen is responsible for false-positive reactions by *Staphylococcus epidermidis* in commercial Agglutination tests for *Staphylococcus aureus* .*J. Clin. Microbiol.*, 39(2):544-550.
- **Blanc, D.S., Lugeon, C., Wenger, A., Siegrist, H. and Francioli, P.(1994):** Quantitative antibiogram typing using inhibition zone diameters compared with ribotyping for epidemiological typing of methicillin –resistant *Staphylococcus aureus*.*J. Clin. Microbiol.*,32: 2505-2509.
- **Blanc, D.S., Struelens, M.J. and Deplano, A.(2001):** Epidemiological validation of pulsed-field gel electrophoresis patterns for methicillin-resistant *Staphylococcus aureus*. *J. Clin. Microbiol* , 39:3442-5.
- **Boyce, J.M., Landry, M., Deetz,T. R. and Dupont, L.H.(1980):**Nosocomial staphylococcal infections. *Ann. Intern. Med*, 95:141-142.
- **Brakstad, O.G. and Maeland, J.A.(1997):** Mechanisms of methicillin resistance in staphylococci., 105(4):264-76
- **Brakstad, O. G., Maeland, J. A. and Tveten, Y. (1993):** Multiplex polymerase chain reaction for detection of genes for *Staphylococcus aureus* thermonuclease and methicillin resistance and correlation with oxacillin resistance. *APMIS*, 101:681-688.
- **Bremer, C., Tung, C.H. and Weissleder, R. (2002):** Molecular imaging of MMP expression and therapeutic MMP inhibition. *Acad. Radiol*, 9 Suppl. 2, 314-315.
- **Brook, G.F., Butel, J.S. and Morse, S.A. (2001):** The *Staphylococci* in medical microbiology Appleton and Lange 23 Ed, 179-203.

- **Brown, D.F.J. (2001):** Detection of methicillin /oxacillin resistance in staphylococci. *J. Antimicrobial. Chemother.* , 48:65-70.
- **Brown, D.F.J and Walpole, E. (2001):** Evaluation of the Mastalex latex agglutination test for methicillin resistance in *Staphylococcus aureus* grown on different screening media. *J.Antimicrobiol.Chemother.*47:187-189.
- **Brown, D.F.J and Yates, V.S. (1986):** Methicillin susceptibility testing of *Staphylococcus aureus* on media containing five percent sodium chloride. *European J. Clin Microbiol and Infect. Dis*, 5:726-728.
- **Bryant, P.A., Li, H.Y., Zaia, A., Griffith, J., Hogg, G., Curtis, N. and Carapetis, J.R. (2004):** Prospective study of a real-time PCR that is highly sensitive, specific, and clinically useful for diagnosis of meningococcal disease in children. *J. Clin. Microbiol*, 42: 2919-2925.
- **Busch, U. and Nitschko, H. (1999):** Methods for the differentiation of microorganisms. *J. Chromatogr. B: Biomed. Sci. Appl.* ,722:263-78.
- **Bustin, S.A. (2000):** Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J. Mol. Endocrinol*, 25, 169-193.

C:

- **Caelli, M.(2000):**Tea tree oil as an alternative topical decolonization agent for methicillin-resistant *Staphylococcus aureus*. *Journal of Hospital. Infection*,46(3):236-237.
- **Cavassini, M., Wenger, A., Jatobn, K. and Blanc, D.S (1999):** Evaluation of MRSA –Screen, a simple Anti-PBP2a slide latex agglutination kit, for rapid detection of Methicillin resistance in *Staphylococcus aureus*. *J. Clin. Microbiol.*, 37(5):1591-1594.

- **Chambers, H.F. (1988):** Methicillin resistant staphylococci. Clin. Microbiol.rev.,1:173-186
- **Chambers, H.F (1997):** Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. Clin Microbiol. Rev., 10(4):781-91.
- **Chambers, H.F. (2001):** The changing epidemiology of Staphylococcus aureus. Emerging. Infect. Dis., 7(2):178-182.
- **Chambers, H.F., Hackbarth, C.J., Drake, T.A., Rusnack, M.G. and Sanda, M.A.(1984):**Endocarditis due to methicillin resistant Staphylococcus aureus in rabbits :expression of resistance to beta -lactam antibiotics in vitro and in vivo. J.Infect .Dis., 149:894-903.
- **Chang, S., Sievert, D.M., Hageman, J.C., Boulton, M.L., Tenover, F.C., et al. (2003):** Infection with vancomycin-resistant Staphylococcus aureus containing the vanA resistance gene. N. Engl. J. Med., 348:1342-1347.
- **Chapin, K.C. and Musgnug, M.C.(2004):**Evaluation of Penicillin Binding Protein 2a Latex Agglutination Assay for Identification of Methicillin-Resistant Staphylococcus aureus Directly from Blood Cultures. Journal of Clinical Microbiology, p. 1283-1284, Vol. 42, No. 3
- **Chen, X., Zehnbaauer, B., Gnirke, A. and Kwok, P.Y. (1997):** Fluorescence energy transfer detection as a homogeneous DNA diagnostic method. Proc .Natl. Acad. Sci. U S A, 94: 10756-10761.
- **Chomyarin,C., Siripornmongcolchi, T., Choicumpor, K., Limpai boon, T., Wongkhom, C. and Yutanaboonthai, W., (2004):**Evaluation of polymerase chain reaction ,conventional and MRSA sceen latex agglutination methods for detection of methicillin resistant ,borderline and susceptible Staphylococcus aureus .Southeast Asian ,J.Trop. Med. Public .Health, 35(4):879-885.
- <http://www.chromagar.com/products/o157.htm/>

- **Chung, M., de Lencastre, H., Matthews, P., Tomasz, A., et al. (2000):** Molecular typing of methicillin resistant *Staphylococcus aureus* by pulsed gel electrophoresis: Comparison of results obtained in a multi laboratory effort using identical protocols and MRSA strains .*Microb .Drug Resist.* ,6;189-198.
- **Collier, L. and Balows, A., (1998):** *Staphylococcus and Micrococcus* In: Collier, T., Balows, A. and Sussman, M., eds. *Topley and Wilson's Microbiology and Microbial Infections*.Vol.2, 9th Ed: 577-611.
- **Cormican, M.G, Wilke, W. and Barrett, M.(1996):** Phenotypic detection of *mec A* positive *Staphi* blood stream isolates .*Diagn. Microbiol. Infect. Dis* ,25:107-112
- **Cosgrove, S. E., Sakoulas, G., Perencevich, E. N., Schwaber, M. J. , Karchmer, A. W. and Carmeli, Y. (2003):** Comparison of mortality associated with methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* bacteremia: a meta-analysis. *Clin. Infect. Dis.* 36:53-59.
- **Coupry, I., Monnet, L., Attia, A.A., Taine, L., Lacombe, D. and Arveiler, B. (2004):** Analysis of CBP (CREBBP) gene deletions in Rubinstein-Taybi syndrome patients using real-time quantitative PCR. *Hum Mutat*, 23: 278-284.
- **Couto, I.(1995):** Unusually large number of methicillin-resistant *Staphylococcus aureus* clones in a Portuguese hospital. *J. Clin. Microbiol*, 33:2032–2035.
- **Couto, I., Wu, S.W., Tomasz, A. and de Lencastre, H. (2003):** Development of methicillin resistance in clinical isolates of *Staphylococcus sciuri* by transcriptional activation of the *mecA* homologue native to the species. *J. Bacteriol.*, 185:645–653
- **Croize, J., Gialanella, P., Monnet, D. and Okaada, J. (1993):** Improved identification of *Staphylococcus aureus* using a new agglutination test, Results of an international study APMIS 101:487-491.

D:

- **Deplano, A., Schuermans, A. and van Eldere, J.(2000):** Multicenter evaluation of epidemiological typing of methicillin-resistant *Staphylococcus aureus* strains by repetitive-element PCR analysis. *J. Clin. Microbiol.* , 38:3527-33.
- **Deplano, A., Vaneechoutte, M., Verschraegen, G. and Struelens, M.J. (1997):** Typing of *Staphylococcus aureus* and *Staphylococcus epidermidis* strains by PCR analysis of inter-IS256 spacer length polymorphisms. *J. Clin. Microbiol.*, 35:2580-7
- **De Silva, N. and Mendis, K.(1998):** One-day general practice morbidity survey in Sri Lanka. *Fam Pract.* , 15(4):323-31.
- **Dietmaier, W. and Hofstadter, F. (2001):** Detection of microsatellite instability by real time PCR and hybridization probe melting point analysis. *Lab. Invest.*, 81: 1453-1456.
- **Domanski, T.L. and Bayles K. W (1995):** Analysis of *Staphylococcus aureus* genes encoding penicillin binding protein 4 and an ABC –type transporter gene.167:111-113.
- **Domann, E., Hossain, H., Fussle, R. and Chakraborty, T.(2000):** Rapid and reliable detection of multiresistent *Staphylococcus aureus* (MRSA) by multiplex PCR. *Dtsch. Med. Wochenschr.*, 19; 125(20):613-8.
- **Duckworth, G. and Healthcock, R. (1995):** Working Party Report. Guidelines on the control of methicillin –resistant *Staphylococcus aureus* in the community. *Journal of Hospital Infection*, 31(1):1-12.
- **Duckworth,G., Lothian, J.L. and Williams,J.D.(1998):** Working Party Report .Revised guidelines for the control of methicillin –resistant *Staphylococcus aureus* infection in hospitals. *Journal of Hospital Infection*, 39(4):253-290.

- **Druham, D .R. and Kloos, W. E.(1978):** A comparative study of the total cellular fatty acids of staphylococcal species of human origin .Int .J. Syst. Bacteriol, 28:233-8

E:

- **Esmon, C.S. and Goodfellow, M.G.(1990):**Staphylococcus and Micrococcus. Cited from Topley and Wilson's .Principals of Bacteriology, Virology and Immunotype.8th ed ,2:161-168.
- **El-Adhami,W. and Stewart, P.R.(1997):**Genome organization of Staphylococcus aureus isolates from different populations. J.Med.Microbiol.46:297-306.
- **Enright, M.C., Day, N.P. and Davies, C.E. (2000):** Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of Staphylococcus aureus. J. Clin. Microbiol., 38:1008-15.
- **Eveillard, M., Quenon, J.L., Rufat, P., Mangeol, A., Fauvelle, F.(2001):**Association between hospital-acquired infections and patients' transfers. Infect Control Hosp Epidemiol. 22(11):693-6

F:

- **Fang, H. and Hedin, G., (2003):** Rapid screening and identification of methicillin –resistant Staphylococcus aureus from clinical samples by Selective –Broth and Real –Time PCR Assay. J. Clin. Microbiol. , P: 2894-1899.
- **Farber, J.M.(1996):** An introduction to the hows and whys of molecular typing. J. Food. Prot .59:1091-101.
- **Felten, A., Grandy, B., Lagrange, P.H .and Casin, I. (2002):** Evaluation of three techniques for detection of low level methicillin resistance Staphylococcus aureus (MRSA): a disk diffusion method with cefoxitin and moxalactam, the vitek 2 system and the MRSA Screen latex agglutination test. J. Clin. Microbiol., 40:2766-2771.

- Fey, P.D., Salim, B. S., Rupp, M.E., Hinrichs, S.H., Boxrud, D.J., Davis, C.C., Kreiswirth, B.N. and Schlievert, P.M.(2003): Comparative molecular analysis of community or hospital acquired methicillin resistant *Staphylococcus aureus*. *Antimicrobial agents and Chemotherapy*, P.196-203.
- Fierobe, L., Decre, D., Muller, C. and Lucet, J.C . (1999): Methicillin resistant *Staphylococcus aureus* as a causative agent of postoperative intra-abdominal infection: Relation to nasal colonization. *Clin. Infect. Dis.*, 29:1231-1238.
- Fluit, A.C., Visser, M.R. and Schmitz, F.J.(2001):Molecular detection of antimicrobial resistance .*Clinical Microbiology Reviews*, 14(4):836-871.
- Frebource, N.B., Nouet, D., Lemee, L., Martin, E. and Lemeland, J.(1998):Comparison of ATB Staph ,Rapid ATB Staph,Vitek ,and E-test methods for detection of oxacillin heteroresistance in staphylococci possessing *mecA*. *J.Clin.Microbiol*, 36:52-57.
- Frenay, H.M., Theelen, J.P., Schouls, L.M. and Vandenbroucke- Grauls, C.M.(1994):Discrimination of epidemic and nonepidemic methicillin-resistant *Staphylococcus aureus* strains on the basis of protein A gene polymorphism. *J Clin Microbiol.*, 32(3):846-7.
- Fridkin, S.K., Hegeman, J. and McDougal, L.K.(2003): Epidemiological and microbiological characterization of infections causes by *Staphylococcus aureus* with reduced susceptibility to vancomycin, United States,. *Clin. Infect. Dis.* 36:429-439

G:

- Gial, L.W. and Washington, J.A. (1995): Antibacterial susceptible tests: Dilution and disk diffusion method .In: *Manual of clinical microbiology* .Patrik, R., Ellen, B., Michael, A., and Robert, H.,(eds);ASM press 6th Ed:1327-1341.

- **Gaston, P.S., Duff, J., Naidoo, K., Ellis, J.I., Roberts, S., Richardson, J.F., Marples, R.R. and Cooka, E.M. (1988):** Evaluation of electrophoretic methods for typing methicillin – resistant *S.aureus* .J. Med. Microbiol., 26:189-197.
- **Ge, M., Chen, Z., Onish, H.R., Kohler, J .and Sliver, L.L.(1999):**Vancomycin derivative that inhibit peptidoglycan biosynthesis without binding D-Ala.NZ Med. J. ,112:105-108.
- **Geary, G., Jordens, J., Richardson. J. and Hawcroft, D.(1997):**J. Med. Microbiol –.46:195-203.
- **Ghuysen, J.M. (1994):** Molecular structures of penicillin-binding proteins and beta-lactamases. Trends Microbiol; 2:372–380.
- **Gibbs, P.J., Tan, L.C., Sadek, S.A. and Howell, W.M. (2003)** Comparative evaluation of 'TaqMan' RT-PCR and RT-PCR ELISA for immunological monitoring of renal transplant recipients. Transpl. Immunol. , 11: 65-72.
- **Gill, V.J., Selepak, A.T., Williams E.D. (1984):** Species identification and antibiotic susceptibilities of coagulase negative staphylococci isolated from clinical specimens .J. Clin. Microiol 18:1314.
- **Ginzinger, D.G., Godfrey, T.E., Nigro, J., Moore, D.H., Suzuki, S., Pallavicini, M.G., Gray, J.W. and Jensen, R.H. (2000):** Measurement of DNA copy number at microsatellite loci using quantitative PCR analysis. Cancer Res., 60: 5405-5409.
- **Giulietti, A., Overbergh, L., Valckx, D., Decallonne, B., Bouillon, R. and Mathieu, C. (2001):** An overview of real-time quantitative PCR: applications to quantify cytokine gene expression. Methods, 25: 386-401.
- **Goering, R.V. (1995):** The application of pulsed electrophoresis to analysis of the global dissemination of methicillin –resistant *Staphylococcus aureus*, methicillin resistant staphylococci, eds Brun –Buisson, C., Casewell, M.W. et al., Medicine-Science Flammarion ,Paris ,75-81.

- **Gonzalez-Zorn, B., Courvalin, P. (2003):** vanA-mediated high level glycopeptide resistance in MRSA. *Lancet. Infect. Dis.*, 3:67-68.
- **Gould, G.C. (1975):** Test for sensitivity to antibiotics for the laboratory .In: Mackie and Mc-Cartney Practical Medical Microbiology, Dujuid, J.P., Marimon, B.P. and Swan I.H.A.(Eds)12th ed,volum(2),Churchill Livingstone.P:199.
- **Grace, M.B., McLeland, C.B., Gagliardi, S.J., Smith, J.M., Jackson, W.E., 3rd and Blakely, W.F. (2003):** Development and assessment of a quantitative reverse transcription-PCR assay for simultaneous measurement of four amplicons. *Clin Chem*, 49, 1467-1475.
- **Greenwood, D.(2000):**,Antimicrobial Chemotherapy. Oxford Press.
- **Gregory, P.D., Lewis, R.A., Curnock, S.P. and Dyke, K.G.(1997):** Studies of the repressor (Blal) of beta-lactamase synthesis in *Staphylococcus aureus*. *Mol. Microbiol.*;24:1025-1037.
- **Griethuysen, V.A., Bes, M., Elienne, J., Zbinden, R. and Kluytmans, J.(2001):** International multicenter evaluation of latex agglutination tests for identification of *Staphylococcus aureus* . *J.Clin.Microbiol.*, 39(1):86-89.
- **Grundmann, H., Hori, S., Winter, B., Tami, A. and Austin, D.J. (2002):** Risk factors for transmission of methicillin – resistant *Staphylococcus aureus* in an adult intensive care unit. *J.Infect. Dis.*, 185:481-488.
- **Gueudet, T. and Lemble, C.(2004):** Comparison of five usual techniques for detection of methicillin resistant *Staphylococcus aureus* .*Path.Biol.*,52(10):617-621.
- **Guy, R.A., Payment, P., Krull, U.J. and Horgen, P.A. (2003):** Real-time PCR for quantification of *Giardia* and *Cryptosporidium* in environmental water samples and sewage. *Appl. Environ. Microbiol.* , 69: 5178-5185.

H:

- **Hackbarth, C.J., Micck, C., and Chambers, H.F.(1994):** Altered production of penicillin binding protein 2a can affect phenotypic expression of methicillin resistance in *Staphylococcus aureus*. *Antimicrobiol Agents Chemother.* 38:2578-2571.
- **Haddadin, A.S., Fappiano, S.A. and Lipsett, P.A.(2002):** Methicillin resistant *Staphylococcus aureus* (MRSA) in the intensive care unit . *Postgraduate Medical Journal*, 78:385-392.
- **Hanaki, H. (1998):** Increase in glutamine-non-amidated muropeptides in the peptidoglycan of vancomycin-resistant *Staphylococcus aureus* strain Mu50. *J. Antimicrob. Chemother.* 42:315-320.
- **Harries, L.W., Wickham, C.L., Evans, J.C., Rule, A., Joyner, M.V. and Ellard, S. (2004):** Analysis of haematopoietic chimaerism by quantitative real-time polymerase chain reaction. *Bone Marrow Transplant*, (in press).
- **Hartshorn, C., Rice, J.E. and Wangh, L.J. (2002):** Developmentally-regulated changes of Xist RNA levels in single preimplantation mouse embryos, as revealed by quantitative real-time PCR. *Mol. Reprod. Dev.*, 61: 425-436.
- **Hazbon, M.H. and Alland, D. (2004):** Hairpin primers for simplified single-nucleotide polymorphism analysis of *Mycobacterium tuberculosis* and other organisms. *J. Clin. Microbiol*, 42: 1236-1242.
- **Heid, C.A., Stevens, J., Livak, K.J. and Williams, P.M. (1996):** Real time quantitative PCR. *Genome Res*, 6: 986-994.
- **Heldman, A.W., Hartert, T. and Ray, S.(1996):** Oral antibiotic treatment of right-sided staphylococcal endocarditis injection drug users .*Am.J.Med* ,101:68-76.

- **Herold, B., Immergluck, L.C. and Maranan, M.C. (1998):** Community acquired methicillin resistant *Staphylococcus aureus* in children with no identified predisposing risk. *JAMA*, 279:593-598.
- **Higashi, Y., wakabayashi, A., Matsumoto, Y., Watanabe, Y. and Ohno, A.(1999):** Role of inhibition of penicillin binding proteins and cell wall cross-linking by Beta-Lactam antibiotics in low-and high-level methicillin resistance of *Staphylococcus aureus*. *J.Infect.Chemotherapy* 45:37-47.
- **Higuchi, R., Fockler, C., Dollinger, G. and Watson, R. (1993):** Kinetic PCR analysis: real-time monitoring of DNA amplification reactions. *Biotechnology (N Y)*, 11: 1026-1030.
- **Hiramatsu, K. (1995):** Molecular evaluation of MRSA. *Microbiol. Immunol.* 39:531-543.
- **Hiramatsu, K.,(1997):** Dissemination in Japanese hospitals of strains of *Staphylococcus aureus* heterogeneously resistant to vancomycin. *Lancet.* , 350:1670-1673.
- **Hiramatsu, K., Cul, L., Kuroda, M., and Ito, T(2001):**The emergence and evolution of methicillin-resistant *Staphylococcus aureus*. *Trends Microbiol.* P:486-493.
- **Hiramatsu, K., Konodo, N., and Ito, T. (1996):** Genetic basis for molecular epidemiology of MRSA. *J. Infect. Chemother.* 2:119-129.
- **Hiramatsu, K., Okuma, K., Ma, X.X., Yamamoto, M., Hori Sand Kapi, M.(2002):**New trends in *Staphylococcus aureus* infections: glycopeptide resistance in hospital and methicillin resistance in the community. *Curr. Opin. Infect. Dis.*
- **Holland, D., Ikram, R., Morris, A. and Sally, R. (2002):** Draft guidelines for the control of Methicillin Resistant *Staphylococcus aureus* in New Zealand by the Ministry of Health PO Box 5013 ,Wellington , New Zealand ISBN 0-478-27066-6 URL:<http://www.moh.govt.nz>.

References

- **Holland, P.M., Abramson, R.D., Watson, R. and Gelfand, D.H. (1991):** Detection of specific polymerase chain reaction product by utilizing the 5'-----3' exonuclease activity of *Thermus aquaticus* DNA polymerase. *Proc. Natl. Acad. Sci. U S A*, 88: 7276-7280.
- **Holt, J.G.(1994):** Gram –positive cocci .In Bergey,s manual of Determinative Bacteriology,9th ed Baltimore Williams and Wilkins .
- **Hooper, D.C.(2002):** Fluoroquinolone resistance among Gram-positive cocci. *Lancet Infect. Dis.*, 2:530–538.
- **Hung, M., Baker, P.N. and Tenver, F.C.(1992):**Accuracy of the E-test for determining antimicrobial susceptibility of *Staphylococcus aureus* . *J. Clin. Microbiol.*, 30:3243-3248.
- **Huletsky, A., Giroux, R., Rossbach, V. and Ganon, M.(2004):** New real time PCR assay for rapid detection of methicillin resistant *Staphylococcus aureus* directly from specimens containing a mixture of staphylococci. *J.Clin. Microbiol. P*: 1875-1884.
- **Huang, M.B., Gay, T.E., Baker, C.N., Banerjee, S.N and Tenover, F.C.(1993):**Two percent sodium chloride is required for susceptibility testing of staphylococci with oxacillin when using agar based dilution methods , *J. Clin. Microbiol.* , 31:2683-2688.
- **Hunt, C., Dionne, M., Derlorme, M. and Murdock, D. (1999):**Four pediatric deaths from community acquired methicillin resistant *Staphylococcus auerus* –Minnesota and North Dakota ,*Arch Dermartol.* 135:1566-8.
- **Hwa, H.L., Ko, T.M., Yen, M.L. and Chiang, Y.L. (2004):** Fetal gender determination using real-time quantitative polymerase chain reaction analysis of maternal plasma. *J. Formos. Med. Assoc.*, 103: 364-368.

I:

- **Ibelings, M.M. and Bruining, H.A.(1998):**Methicillin-resistant *Staphylococcus aureus*: acquisition and risk of death in patients in the intensive care unit. ,164(6):411-8.
- **Idzik, D., Wojtyczka, R., Kepa, M., Wydmuch, Z., Pacha, J., Glab, S., Wroblewska, I. and Skowronek-Ciolek, O.(2000):** Evaluation of methicillin -resistance in *Staphylococcus aureus* by the agar disk diffusion method and PCR. *Med Dosw Microbiol.* 52(4):327-32.
- **Iinuma, Y. (2001):** Identification and epidemiologic investigation of bacteria by the Riboprinter Microbial Characterization System, the automated ribotyping system. *Rinsho.Byori.*49:1129-32.
- http://www.inside-hospitals.co...ents/oxoid/ORSAB_RESAMPLED.jpg

J:

- **Jaffe, R., Lane, J., Albury S.V. and Niemeyer D.M.(2000):** Rapid extraction from and direct identification in clinical samples of methicillin resistant staphylococci using the PCR. *J. Clin. Microbiol.*, 38(9):3407-3412.
- **Jana, M.S., Janet, A.H. and Lance, R.P.(1995):** Special test for detecting antimicrobial resistance in manual of clinical microbiology Patrick, R., et al ., (eds)ASM press 6th Ed 1359-1360.
- **Jana, M., Portia,S., Williams, P., Killgore, G. and Tenover F.C.(2001):** Performance of eight methods , including two new rapid methods, for detection of oxacillin resistance in a challenge set of *Staphylococcus aureus* organism. *J. Clin. Microbiol.*P.3785-3788.

- **Jayaratne, P, and Rutherford, C.(1999):**Detection of methicillin-resistant *Staphylococcus aureus* (MRSA) from growth on mannitol salt oxacillin agar using PCR for nosocomial surveillance. *Diagn. Microbiol Infect Dis.* ,35(1):13-8.
- **Jeršek, B., Tcherneva, E., Rijpens, N. and Herman, L.(1996):** Repetitive element sequence-based PCR for species and strain discrimination in the genus *Listeria*. *Lett .Appl. Microbiol.* 23:55-60.
- **Johonson, A. P. and Speller, D.E.(1997):**Epidemiology of antibiotic resistance :blood and cerebrospinal fluid .In *Antibiotic resistance* Edited by SGB Amyes and CD Gemmell.J. *Med.Microbiol*,46:445-447.
- **Joklik, W.H., Willet, H.P., Aos, D.B. and Wilfert, C.M.(1988):** Zinsser Microbiology Nineteenth Edition, Appleton and Lange. Norwalk, Connecticut/Scan Mateo, California,(9) :343-354.
- **Jonas, D., Speck, M., Daschner, F.D. and Grundmann, H.(2002):** Rapid PCR-Based Identification of Methicillin-Resistant *Staphylococcus aureus* from Screening Swabs. *J. Clin. Microbiol.*, 40(5): 1821–1823.
- **Jong, W.U., Hung, H., Hwa,J. and Tsung, C. (1997):** Rapid detection of oxacillin resistant *Staphylococcus aureus* in blood culture by an impedance method. *J .Clin .Microbiol.*, 35(6):1460-1464.

K:

- **Kampf, G., Lecke, C., Cimbali, A.C. and Weist, K. (1998):** Evaluation of mannitol salt agar for detection of oxacillin resistance in *Staphylococcus aureus* by disk diffusion and agar screening. *J.Clin. Microbiol.*, 36(8):2254-2257.
- **Kampf, G., Weist, K., Kegel, M., Swidsinski, S. and Ruden, H. (1997):** Comparison of screening methods to identify

methicillin-resistant *Staphylococcus aureus*. *Eur. J. Clin. Microbiol. Infect. Dis.* 16:301-307.

- **Katayama, Y., Ito, T. and Hiramatsu, K. (2000):** A new class of genetic element, staphylococcus cassette chromosome mec, encodes methicillin resistance in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.*, 44:1549–1555.
- **J Kearns, A.M., Graham, C., Burdess, D., Heatherington, J. and Freeman, R. (2002):** Rapid real-time PCR for determination of penicillin susceptibility in pneumococcal meningitis, including culture-negative cases. *J Clin Microbiol*, 40, 682-684.
- **Kearns, A. M., Seiders, P. R., Wheeler, J., Freeman, R. and Steward, M. (1999):** Rapid detection of methicillin-resistant staphylococci by multiplex PCR. *J. Hosp. Infect.*, 43:33-37.
- **Kernodle, D.S. (2000):** Mechanisms of resistance to β -lactam antibiotics. In *Gram-positive pathogens*. Fischetti, V.A., Novick, R.P., Ferretti, J.J., Portnoy, D.A. and Rood, J.I. Editors. American Society for Microbiology. Washington, DC, USA. 609–620.
- **Kipp, F., Becker, K., Peters, G. and Von Eiff, C. (2004):** Evaluation of different methods to detect methicillin resistance in small colony variants of *Staphylococcus aureus*. *J. Clin. Microbiol.* P:1277-1279.
- **Kirst, H.A, Thompson, D.G. and Nicas, T.I.(1998):** Historical yearly usage of vancomycin. *Antimicrob. Agents Chemother.*, 42:1303–1304.
- **Kloos, W.E. and Jorgensen, J.H. (1985):** *Staphylococci*. *Manual of Clinical Microbiology*, 4th ed(EH), Lennete, A., Balows, E.J., Hausler, H.J., Shadomy (eds), Washington DC, American Society for Microbiology, P:143-153, 1985.
- **Kluytmans, J., Griethuysen, V.A., Willemse, P. and Van Keulen, P. (2002):** Performance of CHROMagar selective medium and oxacillin resistance screening agar base for

identifying *Staphylococcus aureus* and detecting methicillin resistance. *J.Clin.Microbiol.*, 40(7):2480-2482.

- **Kluytmans, J., Van Belkum, A., and Verbrugh, H. (1997):** Nasal carriage of *Staphylococcus aureus*: epidemiology underlying mechanisms, and associated risks. *Clin. Microbiol. Rev.*, 10(3):505-520.
- **Kolbert, C. P., Connolly, J. E., Lee, M. J. and Persing, D. H. (1995):** Detection of the staphylococcal *mecA* gene by chemiluminescent DNA hybridization. *J. Clin. Microbiol.* 33:2179-2182.
- **Koneman, E.W., Allen, S.D., Janda, W.M., Schreckenberger, P. C. and Jr, W.C.(1997):**The gram positive cocci: staphylococci and related organisms .In color atlas and textbook of diagnostic microbiology.5th edition,JB Lippincott company,Philadelphia:539-557.
- **Kosarsky, P.E.,Rimland,D., Terry, P.and Wachsmuth, K.(1986):**Plasmid analysis of simultaneous nosocomial outbreaks of methicillin resistant *Staphylococcus aureus*. *Infect.Control*,7:577-581.
- **Kühn, I., Burman, L. G., Hæggman, S., Tullus, K. and Murray, B. E.(1995):** Biochemical fingerprinting compared with ribotyping and pulsed-field gel electrophoresis of DNA for epidemiological typing of enterococci. *J. Clin. Microbiol.*, 33:2812-2817.
- **Kumari, D.N.P., Keer, V. and Hawkey, P.M.(1997):** Comparison and application of ribosome spacer DNA amplicon polymorphisms and pulsed-field gel electrophoresis for differentiation of methicillin-resistant *Staphylococcus aureus* strains. *J. Clin. Microbiol.* , 35:881-5.

L:

- **Lacey, R.W., Barr, V.E. and Inglis, T. J. (1986):** Properties of methicillin resistant *Staphylococcus aureus* colonizing patients in a burn unit. *J.Hosp .Infect.* , 7:137-148.
- **Lally, R.T., Ederer, M.N. and Woolfrey, B.F.(1985):** Evaluation of mannitol salt agar with oxacillin as a screening medium for methicillin resistant *Staphylococcus aureus*. *J.Clin.Microbiol.*22:501-504.
- **Lancoster, H. and Tomasz, A. (1994):** Number of auxiliary genes essential for expression of high level resistance in *S.aureus*. *Antimicrobial .Agents .Chemother.*, 38:5290-5298.
- **Lareu, M.V. and Ruiz-Ponte, C. (2004):** Genotyping SNPs with the LightCycler. *Methods Mol. Biol.*, 297: 127-140.
- **Le Dell, K., Naimi, T., Borchardt, S. and Boxrud, D. (2002):** Community-Onset and Healthcare-Associated Methicillin Resistant *Staphylococcus aureus* in Minnesota. *International conference on emerging infectious disease*, p 74.
- **Lee, L.G., Connell, C.R. and Bloch, W. (1993):** Allelic discrimination by nick-translation PCR with fluorogenic probes. *Nucleic Acids Res*, 21: 3761-3766.
- **Leruez-Ville, M., Minard, V., Lacaille, F., Buzyn, A., Abachin, E., Blanche, S., Freymuth, F. and Rouzioux, C. (2004):** Real-time blood plasma polymerase chain reaction for management of disseminated adenovirus infection. *Clin Infect. Dis*, 38, 45-52.
- **Leslie, C., Albert, B. and Max, S.(1998):** *Staphylococcus* and *Micrococcus* .Cited from Topley and Wilson's principles of Bacteriology ,Virology and Immunology .9th ed . Microbial infection chapter 27(2):577-611.
- **Leung, W., Iyengar, R., Triplett, B., Turner, V., Behm, V.G., Holladay, M.S., Houston, J. and Handgretinger, R. (2005):** Comparison of Killer Ig-Like Receptor Genotyping and

Phenotyping for Selection of Allogeneic Blood Stem Cell Donors. *The Journal of Immunology*, 174: 6540-6545.

- **Ligozzi, M., Rossolini, G. M., Tonin, E. A. and Fontana, R.(1991):** Nonradioactive DNA probe for detection of gene for methicillin resistance in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 35:575-578.
- **Lim, D. and Strynadka, N.C. (2002):** Structural basis for the beta lactam resistance of PBP2a from methicillin-resistant *Staphylococcus aureus*. *Nat. Struct. Biol.*, 9:870-876.
- **Livak, K.J., Flood, S.J., Marmaro, J., Giusti, W. and Deetz, K. (1995):** Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl*, 4: 357-362.
- **Livak, K.J. and Schmittgen, T.D. (2001):** Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, 25: 402-408.
- **Louie, L., Matsumura, S.O., Choi, E., Louie, M. and Simor, A.E.(2000):**Evaluation of three rapid methods for detection of methicillin resistance in *Staphylococcus aureus*. *J.Clin.Microbiol.*, 38:2170-2174.
- **Louie, L., Goodfellow, J., Mathieu, P., Glatt, A., Louie, M., and Simor, A.E.(2002):** Rapid Detection of Methicillin-Resistant *Staphylococci* from Blood Culture Bottles by Using a Multiplex PCR Assay. *Clin Microbiol.* ,40(8): 2786-2790.
- **Lovatt, A. (2002):** Applications of quantitative PCR in the biosafety and genetic stability assessment of biotechnology products. *J. Biotechnol.* , 82, 279-300.
- **Lowy F.D.(2000):**Is *Staphylococcus aureus* an intracellular pathogen?.*Trends Microbial.*, 8:341-343.
- **Lowy F.D (2003):** Antimicrobial resistance: the example of *Staphylococcus aureus*. *J. Clin. Invest.* 1; 111(9): 1265-1273.

M:

- **Mackenzie, A.M., Richardson, H., Langigan, R. and Wood, D.(1995):** Evaluation for the disk diffusion test for detection of low expression class MRSA .J. Clin. Microbiol. ,33(7):1909-1911.
- **Maiden, M.C., Bygraves , J.A. and Feil , E.(1998):** Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms.Proc.Natl.Acad.Sci.,95:3140-5.
- **Maki, H., Yamaguchi, T. and Murakami, K.(1994):**Cloning and characterization of a gene affecting methicillin resistance level and the autolysis rate in Staph.aureus. J. Bacteriol., 176:4993-5000.
- **Malouin, F., Blais, J., Chamberland, S. and Hoang, M. (2003):**RWJ-54428(Mc-02,479), a New Cephalosporin with high affinity for Pencillin-Binding Proteins ,including PBP2a. Antimicrob. Agents Chemother.,47(2):658-664.
- **Marples, R. and Rosdhi, V., (1997):** International quality control of phage typing of Staphylococcus aureus .Med. Microbiol. 46:511-516.
- **Maslow, J.N., Mulligan, M.E. and Arbeit R.D.(1993):** Molecular epidemiology: application of contemporary techniques to the typing of microorganisms. Clin Infect Dis., 17:153-64.
- **Massanari, R.M., Pfaller, M.A., Wakefield, D.S., Hammons, G.T., McNutt,L.A., Woolson,R.F. and Helms,C.M.(1988):** Implications of acquired oxacillin resistance in the management and control of Staphylococcus aureus infections. J Infect Dis.,158(4):702-9
- **Mc Niel, S.A., Mody, L. and Bradley, S.F.(2002):**Methicillin-resistant Staphylococcus aureus. Management of asymptomatic

colonization and out breaks of of infection in long term care.
Geriatrics .57(6):16-27.

- **Mengelle, C., Pasquier, C., Rostaing, L., Sandres-Saune, K., Puel, J., Berges, L., Righi, L., Bouquies, C. and Izopet, J. (2003) :** Quantitation of human cytomegalovirus in recipients of solid organ transplants by real-time quantitative PCR and pp65 antigenemia. *J Med Virol*, 69: 225-231.
- **Merlino, J., Ieroi, M., Bradbury, R., Veal, D., and Harbour, C.(2000):**New chromogenic identification and detection of *Staphylococcus aureus* and methicillin resistant *S.aureus*. *J.Clin.Microbiol.*, 38:2378-2380.
- **Merlino, T., Gill, R. and Robertson, G.J.(1996):**Application of lipovitellin mannitol salt agar for screening ,isolation ,and presumptive identification of *staphylococcus aureus* in a teaching hospital. *J.Clin.Microbiol.*34:3012-3015.
- **Merrer, J., Santoli, F. and Appere de Vecchi, C. (2000):** Colonization pressure and risk of acquisition of methicillin-resistant *Staphylococcus aureus* in a medical intensive care unit. *Infect.Control Hosp.Epidemiol* . 21:718-723.
- **Michael, O. D. and Bean, P. (1999):** Principles and Applications of Methods for DNA-Based Typing of Microbial Organisms *Journal of Clinical Microbiology*, 37(6):1661-1669.
- **Milne, L.M., Crow, M., Emptage, A.G. and Selkon, J.B.(1993):**Effects of culture media on detection of methicillin resistance in *Staphylococcus aureus* and coagulase negative staphylococci by disc diffusion methods .*J.Clin. Pathol* .46:394-397.
- **Mir, N., Sanchez, M. and Morgan, T.(1998):** Evaluation of ORSAB as a selective medium for detection of MRSA. *J.Clin.Microbiol.*36:986-989.

- Mizobuchi, S., Minami, J., Jin, F., Matsushita and Okabe A(1994) :Comparison of the virulence of methicillin –resistant and methicillin sensitive Staphylococcus aureus. Microbiol. Immunol, 38(8): P: 599-605.
- Moreno, F., Crisp, C., Jorgensen, J.H. and Patterson, J.E.(1995): Methicillin-resistant Staphylococcus aureus as a community organism. Clin Infect Dis.,21(5):1308-12.
- Moriki, S., Iwata, Y., Ishikura, H. and Endo, J.(1995): Detection of heterogeneous MRSA by using the PCR method and population analysis. Rinsho Byori.;43(5):487-92.
- Morrison, T.B., Weis, J.J. and Wittwer, C.T. (1998): Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. Biotechniques, 24: 954-958, 960,-962.
- Mulder, J.G.(1996):Comparison of disk diffusion ,the E-test,and detection of mecA for determination of MRSA. European J of Clinical Microbiology and Infect Dis.,15(7):567-573.
- Mulvey, M.R., Chui, L., Ismail, J. and Louie, L.(2001):Development of a Canadian Standardized Protocol for subtyping methicillin –resistant Staphylococcus aureus using Pulsed –Field Gel electrophoresis. J. Clin. Micobiol., 39(10):3481-3485.
- Murakami, K., Minamide, W., Wada, K., Nakamura, E., Teraoka, H. and Watanabe, S. (1991): Identification of methicillin-resistant strains of staphylococci by polymerase chain reaction. J. Clin. Microbiol. 29:2240-2244.
- Mylotte, J.M.(1994): Control of MRSA: The ambivalence persists. Infect.Cont.Hospi.Epidemiol.15:73-77.

N:

- NCCLS (National Committee for Clinical Laboratory Standards) (1993): Performance standards for antimicrobial

susceptibility testing. 5th Ed Approved Standard. National Committee for Clinical Laboratory Standards, Villanova, Pa.

- **NCCLS (National Committee for Clinical Laboratory Standards) (1999):** Performance standards for antimicrobial susceptibility testing. 9th informational supplement. National Committee for Clinical Laboratory Standards, Villanova, Pa.
- **NCCLS (National Committee for Clinical Laboratory Standards) (2000):** Performance standards for antimicrobial susceptibility testing. Approved Standard M100-S10, NCCLS, Wayne, Pa.
- **Ng, E.Y., Trucksis, M. and Hooper, D.C. (1996):** Quinolone resistance mutations in topoisomerase IV: relationship to the flqA locus and genetic evidence that topoisomerase IV is the primary target and DNA gyrase is the secondary target of fluoroquinolones in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.*, 40:1881–1888.
- **Nicola, F., Bantar, C. and Canigia, L.F. (2000):** Comparison of several methods to determine MRSA. *Diag. Microbiol. Infect. Dis.* 36(2):91-93.
- **Nicolle, L.E. (2001):** Infection control programs to contain antimicrobial resistance WHO/CDS/CSR/DRS/2001.7.
- **Niemeyer, D.M., Pucci, M.J., Thanassi, J.A., Sharma, V.K. and Archer, G.L. (1996):** Role of *mecA* transcriptional regulation in the phenotypic expression of methicillin resistance in *Staphylococcus aureus*. *J. Bacteriol.*; 178:5464–5471.
- **Novak, S.M., Hindler, J. and Bruckner, D.A. (1993):** Reliability of two novel methods Almer and E test for detection of MRSA. *J. Clin. Microbiol.*, 31:305-307.

O:

- **Olive, D.M., Bean, P. (1999):** Principles and applications of methods for DNA-based typing of microbial organisms. *J. Clin.*

Microbiol.37:1661-9.

- **O'Sullivan, N. and Kean, C.(2000):**The prevalence of methicillin-resistant *Staphylococcus aureus* among the resident of six nursing homes for the elderly . *Journal of Hospital Infection*.45 (4):322-329.

P:

- **Patrick, R.M., George, S.K., Michael, A.p. and Ken, S.R. (1994):** Medical microbiology, second Ed .Mosby-Year Book Inc., LE International Edition, (17)167-179.
- **Peacock, J.E., Marsik F.J. and Wenzel R.P.(1980):** *Staphylococcus aureus* :introduction and spread within a hospital .*Ann.Intern.Med*,93:526-523.
- **Perry, J.D., Rennison, C., Butterworth, L.A., Hopley, A.L. and Gould, F.K.(2003):**Evaluation of *S.aureus* ID, a New Chromogenic Agar medium for detection of *Staphylococcus aureus*. *Journal of Clinical Microbiology*, p. 5695-5698, Vol. 41, No. 12.
- **Personne, P., Bes, M., Lina, G. and Vandenesch, F. (1997):** Comparative performance of six agglutination kits assessed by using typical and atypical strains of *Staphylococcus aureus*. *J.Clin. Microbiol.*, 35:1138-1140.
- **Persson, K., Hamby, K. and Ugozzoli, L.A.(2005):** Four-color multiplex reverse transcription polymerase chain reaction-overcoming its limitations ,*Analytical Biochemistry* , 344: 33-42.
- **Pfaffl, M.W. (2001)** A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res*, 29, e45.
- **Pinho, M.G., de Lancoster, H. and Tomasz, A.(2000):** Cloning, Characterization ,and Inactivation of the Gene *pbpC*, Encoding penicillin –binding protein 3of *Staphylococcus aureus*. *Journal of Bacteriology*, 182(4):1074-1079.

- **Rajeevan, M.S., Vernon, S.D., Taysavang, N. and Unger, E.R. (2001):** Validation of array-based gene expression profiles by real-time (kinetic) RT-PCR. *J Mol Diagn*, 3: 26-31.
- **Rampling, A., Wiseman, S. and Davis, L. (2001):** Evidence that hospital hygiene is important in the control of Methicillin –resistant *Staphylococcus aureus* . *J.Hosp.Infect.*49:109-116
- **Rayan, K.J.(1994):**Staphylococci in Rayan K.J (edit) Shetties Medical Microbiology 3rd Ed ,Prentice-Hall Inc. London.253-263.
- **Rayner, D. (2003):** MRSA: an infection control overview. *Nursing Standard*.17 (45):47-53.
- **Rego V.H., Hageman J.C., Sitz J. and Baughman W. (2002):** The Epidemiology of Community –Onset Methicillin –Resistant *Staphylococcus aureus*(MRSA)in the Atlanta Metropolitan Area. International Conference on emerging infectious disease P.76.
- **Reischl , U., Linde , H. J. , Metz, M., Leppmeier , B. and Lehn , N.(2000):**Rapid identification of methicillin-resistant *Staphylococcus aureus* and simultaneous species confirmation using real-time fluorescence PCR . *J. Clin. Microbiol.* 38:2429-2433.
- **Resende, C.A. and Figueiredo, A.M.(1997):**Discrimination of methicillin resistant *Staphylococcus aureus* from borderline resistant and susceptible isolates by different methods. *J.Med.Microbiol.*46:145-149.
- **Ririe, K.M., Rasmussen, R.P. and Wittwer, C.T. (1997):** Product differentiation by analysis of DNA melting curves during the polymerase chain reaction. *Anal Biochem*, 245: 154-160.
- **Roberts, J. I. and Gaston , M .A.(1987):**Protein A and coagulase expression in epidemic and non-epidemic *Staphylococcus aureus* J Clin Pathol.; 40(8): 837–840.
- **Rosato, A.E. (2003):** mecA-blaZ co-repressors in clinical *Staphylococcus aureus* isolates. *Antimicrob. Agents Chemother.*; 47:1460–1463.

S:

- **Safdar, N., Narans, L., Gordon, B. and Maki,D.G.(2003):** Comparison of culture screening methods for detection of nasal carriage of methicillin resistant *Staphylococcus aureus* :a prospective study comparing 32 methods . J. Clin. Microbiol. 41(7):3163-3166.
- **Safah, A.S., Wassef,M. and Sayed, A.(2001):**Analysis of risk factors causing resistance to oxacillin and ofloxacin among clinical isolates of MRSA. Faculty of Medicine Cairo University.
- **Safdur,N. and Maki D.G.(2002):** The commonality of risk factors For nosocomial colonization and infection with antimicrobial – resistant *Staphylococcus aureus*, enterococcus, gram-negative bacilli, *Clostridium difficile* and candida .Ann. Intern. Med.136:834-844.
- **Salisbury, S. M., Sabatini, L. M. and Spiegel, C. A.(1997):** Identification of methicillin-resistant staphylococci by multiplex polymerase chain reaction assay. Am. J. Clin. Pathol. 107:368-373.
- **Salmenlinna, S., Lyytikainen, Q. and Varkila, J.V.(2002):** Community Acquired Methicillin Resistant *Staphylococcus aureus*. Finland Emerging Infectious Disease, 8(6):602-607.
- **Saunders, N.J. (1995):**Vancomycin administration and monitoring reappraisal. J.Antmicrob.Chemother.36:279-282.
- **Schuermans, H., Peetermans, W.E., Struelens, M.J. and Van – Lierde, S. (1997):** Contribution of a typing methods based on IS256 probing of *Sma*I –digested cellular DNA to discrimination of European type 77 methicillin –resistant *Staphylococcus aureus* strains J. Clin. Microbiol., 2514-2520.
- **Schwartz, D.C., Cantor, C.R. (1984):** Separation of yeast chromosome–sized DNAs by pulsed field gradient gel electrophoresis cell .37:67-75.

- **Sekiguchi, k., Saito, M. and Yahjima, R.(1995):** Detection of MRSA with antibodies against synthetic peptides derived from PBP2 ,Microbiology and Immunology ,39(8):545-550.
- **Shiff B. (1999):** VRE and MRSA, putting Bad Bugs out of Business. Nursing Mangement, 30(6).
- **Shopsin, B.O. and Kreiswirth, B.N.(2001):**Molecular epidemiology of methicillin –resistant Staphylococcus aureus. Emerging Infection Diseases. 7(2):223-226.
- **Showsh, S.A., De Boever, E.H., Clewell, D.B.(2001):** Vancomycin resistance plasmid in Enterococcus faecalis that encodes sensitivity to a sex pheromone also produced by Staphylococcus aureus. Antimicrob. Agents. Chemother., 45:2177-2178.
- **Sieradzki, K., Roberts, R.B., Haber, S.W., Tomasz, A.(1999):** The development of vancomycin resistance in a patient with methicillin- resistant Staphylococcus aureus infection. N. Engl. J. Med., 340:517–523.
- **Simor, A.E., Goodfellow, J., and Marie, L.L.(2001):**Evaluation of a new medium ,Oxacillin Resistance Screening Agar Base ,for the detection of methicillin –resistant Staphylococcus aureus from clinical specimens. J.Clin. Microbiol. 39(9):3422-3424.
- **Siraj, A.K., Ozbek, U., Sazawal, S., Sirma, S., Timson, G., Al-Nasser, A., Bhargava, M., El Solh, H., Bhatia, K. and Gutierrez, M.I. (2002):** Preclinical validation of a monochrome real-time multiplex assay for translocations in childhood acute lymphoblastic leukemia. Clin Cancer Res, 8: 3832-3840.
- **Skinner, S., Inglis, B., Matthews, P.R. and Stewart, P.R.(1988):**Mercury and tetracycline resistance genes and repeats associated with methicillin resistance on the chromosome of Staphylococcus aureus .Mol. Microbiol., 2:289-297.
- **Smith, T. L., Pearson, M. L., Wilcox, K. R., Cruz, C., Lancaster, M. V., Robinson-Dunn, B., Tenover, F. C., Zervos, M. J. , Band, J. D., White, E. and Jarvis, W. R.(1999):**

Emergence of vancomycin resistance in *Staphylococcus aureus*. N. Engl. J. Med. 340:493-501.

- **Smyth, R.W., Kahlmeter, G., Liljequist, B.O. and Hoffman, B.(2001):**Methods for identifying methicillin resistance in *Staphylococcus aureus* .J.Hosp.Infect.48(2);103-107.
- **Soares, M.J., Tokumarw-Miyazaki, N.H., Noieto, A.L. and Figueriedo, A.M. (1997):**Enterotoxin production by *S.aureus* clones and detection of Brazilian epidemic MRSA clone among isolates from food handlers .J.Med .Microbiol ,46:214-221.
- **Soares, M.J., Soares, C., Mendes, C.A., Guimar, M.L ., Cabeda, J.M. and Amarim, J.M.(2004):**Assessment of three rapid methods for the detection of methicillin –resistant *staphylococcus aureus* .Enferm. Infecc .Micrbiol. Clin 22:390-391.
- **Solinas, A., Brown, L.J., McKeen, C., Mellor, J.M., Nicol, J., Thelwell, N. and Brown, T. (2001):** Duplex Scorpion primers in SNP analysis and FRET applications. Nucleic Acids Res, 29, E96.
- **Sparham S., P.D., Lobban, D.I. and Speller, .C.E. (1978):** Isolation of *Staphylococcus aureus* from sputum in cystic fibrosis. J.Clin. Pathol. , 31:913-918.
- **Stevens, D.L.(2002):**Linezolid versus Vancomycin for the treatment of methicillin-resistant *Staphylococcus aureus* infections. Clin. Infect .Dis.34:1481-1490.
- **Stones, S., Beric, V., Quick, A., Balestrini, A.A. and Kibbler, C.C.(2001):**Controlling infection in British nursing homes :it is time for a national strategy .British Medical Journal. 314(7258)505.
- **Struelens, M.J.(1998):** Molecular epidemiologic typing systems of bacterial pathogens: current issues and perspectives. Mem. Inst. Oswaldo.Cruz.93:581-5.
- **Struelens, M.J., Nonhoff, C., Auwera P.V., Mertens, R., et al., (1995):** Evaluation of Rapid ATB Staph for 5-hour antimicrobial susceptibility testing of *Staphylococcus aureus*. J. Clin. Microbiol. 33:2395-2399

T:

- **Tanguchi, H., Kaneko, T., Onozaki, M., Kubo, R. and Kamiya, S.(2004):**CHROM agar ^{IM} MRSA :Evaluation of new chromogenic medium for isolation of MRSA .Journal of the Japanese Association for Infective Diseases ,P:24-58.
- **Tan, T .Y., Corden, S., Barnes, R., and Cookson, B.(2001):** Rapid Identification of Methicillin-Resistant Staphylococcus aureus from Positive Blood Cultures by Real-Time Fluorescence PCR J .Clin. Microbiol., 39(12): 4529–4531.
- **Tang, Yi-Wie ., Waddington, M.G., Smith, D.H. and Manahan, J.M. (2000):**Comparison of protein A gene sequencing with pulsed –field gel electrophoresis and epidemiologic data for molecular typing of methicillin –resistant Staphylococcus aureus J. Clin. Microbiol., 32:407-415.
- **Tenover, F.C., Arbeit, R.D. and Goering, R.V. (1997):** How to select and interpret molecular strain typing methods for epidemiological studies of bacterial infections: a review for healthcare epidemiologists. Infect.Control.Hosp.Epidemiol.18:426-38.
- **Tenover, F.C. and Gaynes, R.P.(2000):**The epidemiology of staphylococcus infection P:414-421.In Fischetti V A., Novick R.P., Ferrlti J.J ., Portnoy P.A. and Rood J I(ed),Gram –positive pathogen .American Society for Microbiology, Washington D.C.
- **Terry, C.F., Shanahan, D.J., Ballam, L.D., Harris, N., McDowell, D.G. and Parkes, H.C. (2002):** Real-time detection of genetically modified Soya using Lightcycler and ABI 7700 platforms with TaqMan, Scorpion, and SYBR Green I chemistries. J AOAC Int, 85, 938-944.
- **Thomassin, H., Kress, C. and Grange, T. (2004)** MethylQuant: a sensitive method for quantifying methylation of specific cytosines within the genome. Nucleic Acids Res, 32: e168.
- **Toku, Y., Shoji, S., Satoh, K., Watanabe, A. and Motomiye, M,(1991):** Detection of methicillin resistance staphylococcus

aureus using polymerase chain reaction amplification . Tohokue,J.
Exp.Med.P:31-37.

- **Tomasz, A., Drugeon, H. B. , de Lencastre, H. M. , Jabes, D., McDougall, L. and Bille, J.(1989):** New mechanism for methicillin resistance in *Staphylococcus aureus*: clinical isolates that lack the PBP2a gene and contain normal penicillin-binding protein with modified penicillin-binding capacity. *Antimicrob. Agents Chemother.* ,33: 1869-1874.
- **Towner, K. J., Talbot, D. C. S., Curran, R., Webster, C. A. and Humphreys, H.(1998):** Development and evaluation of a PCR-based immunoassay for the rapid detection of methicillin-resistant *Staphylococcus aureus*. *J. Med. Microbiol.* 47:607-613.
- **Trindade, P.A., McCulloch, J.A., Oliveira, G.A. and Mamizuka, E.M.(2003):**Molecular techniques for MRSA typing :current issues and perspectives .*Braz J Infect Dis* vol .7 no .1 Salvador.

U:

- **Ubukata, K., Nakagami, S., Nitta, A. , Yamane, A., Kawakami, S. , Sugiura, M. and Konno, M.(1992):** Rapid detection of the *mecA* gene in methicillin-resistant staphylococci by enzymatic detection of polymerase chain reaction products. *J. Clin. Microbiol.* 30:1728-1733.
- **Ünal, S., Hoskins, J., Flokowitsch, J. E., Wu, C. Y., Preston, D. and Skatrud, P. L. (1992):** Detection of methicillin-resistant staphylococci by using the polymerase chain reaction. *J. Clin. Microbiol.* , 30:1685-1691.
- **Ünal, S., Werner, K., DeGirolami, P., Barsanti, E., and Eliopoulos, G.(1994):** Comparison of tests for detection of methicillin-resistant *Staphylococcus aureus* in a clinical microbiology laboratory. *Antimicrob. Agents Chemother.* 38:345-347.

V:

- **Van belkum, A., Kluitmans, J. and Van leeuwen, W. (1995):** Multicenter evaluation of arbitrarily primed PCR for typing of *Staphylococcus aureus* strains. *J .Clin. Microbiol.*, 33:1537-47.
- **Van der Velden, V.H., Hochhaus, A., Cazzaniga, G., Szczepanski, T., Gabert, J. and Van Dongen, J.J. (2003):** Detection of minimal residual disease in hematologic malignancies by real-time quantitative PCR: principles, approaches, and laboratory aspects. *Leukemia*, 17: 1013-1034.
- **Van leeuwen, W., Verbrugh, H. and Van der Velden, J. (1999):** Validation of binary typing for *Staphylococcus aureus* strains. *J.Clin.Microbiol.*, 37:664-74.
- **Van Wannel, W.J., Fliut, A.D., Wardstom, T. and Dijk, V.H. (1995):** Phenotypic characterization of epidemic versus sporadic strains of methicillin –resistant *Staphylococcus aureus*. *J.Clin.Microbiol.*, 33:1769-1774.
- **Van Enk, R.A. and Thompson, K.D.(1992):** Use of a primary isolation medium for recovery of methicillin –resistant *Staphylococcus aureus* . *J.Clin.Microbiol.* 30:504-505.
- **Velasco, D., Tomas, M.M., Cartelle, M. and Beceiro, A. (2005):** Evaluation of different methods for detecting methicillin (oxacillin) resistance in *staphylococcus aureus* *J.Clin .Microbiol.*, 55:379-382.
- **Versalovic, J., Kapur, V. and Lupski, J.R.(1991):** Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Res.* 19:6823-31.
- **Vet, J.A. and Marras, S.A.(2005):** Oligonucleotide Synthesis: Design and Optimization of Molecular Beacon Real-Time Polymerase Chain Reaction Assays, ISBN *Methods in Molecular Biology.*, (288):273-290.

- **Von Ahsen, N., Armstrong, V.W. and Oellerich, M. (2004):** Rapid, Long-Range Molecular Haplotyping of Thiopurine S-methyltransferase (TPMT) *3A, *3B, and *3C. Clin Chem.
- **Vouillamoz, J., Entenza, M.J., Hohl, P. and Moreillon P. (2004):** LB11058, a New Cephalosporin with High Penicillin-Binding Protein 2a Affinity and Activity in Experimental Endocarditis Due to Homogeneously Methicillin-Resistant *Staphylococcus aureus*. Antimicrob. Agents Chemother. November; 48(11): 4322–4327.
- **Vrettou, C., Traeger-Synodinos, J., Tzetis, M., Palmer, G., Sofocleous, C. and Kanavakis, E. (2004):** Real-time PCR for single-cell genotyping in sickle cell and thalassemia syndromes as a rapid, accurate, reliable, and widely applicable protocol for preimplantation genetic diagnosis. Hum Mutat, 23: 513-521.

W:

- **Wadlvoegel, F.A. (2000) :** *Staphylococcus aureus*. In :Mandell GL, Bannett JE, Dolin R, eds. Principles and practice of infectious disease .5th ed .Philadelphia: Churchill Livingstone, P:2072-3.
- **Walsh, C.T. (1993):** Vancomycin resistance :decoding the molecular logic Science 261:308-309.
- **Walsh, T.R., Howe, R.A. (2002):** The prevalence and mechanisms of vancomycin resistance in *Staphylococcus aureus*. Annu. Rev. Microbiol., 56:657–675.
- **Wang, X. and Seed, B. (2003):** A PCR primer bank for quantitative gene expression analysis. Nucleic Acids Res, 31: e154.
- **Watson, A. and Brown, D. F. J. (1998):** methods of determining susceptibility of *Staphylococcus aureus* to methicillin and oxacillin. In Abstracts of the thirty eighth interscience conference on antimicrobial agents and chemotherapy ,San Diego, CA, Abstract D 53 ,P.144 ,Washington DC ,American Society for Microbiology.

- **Watzinger, F., Suda, M., Preuner, S., Baumgartinger, R., Ebner, K., Baskova, L., Niesters, H.G., Lawitschka, A. and Lion, T. (2004):** Real-time quantitative PCR assays for detection and monitoring of pathogenic human viruses in immunosuppressed pediatric patients. *J Clin Microbiol*, 42, 5189-5198.
- **Webster, J.A. and Bannerman, T.L. (1994):** Identification of the *Staphylococcus sciuri* species group with *EcoR*I fragments containing rRNA sequences and description and description of *Staphylococcus vitulus* sp.nov., *Int .Syst Bacteriol*,44:454-60.
- **Weller, T.M.A(1999):**The distribution of *mecA*, *mecR*I and *mecI* and sequence analysis of *mecI* and the *mec* promoter region in staphylococci expressing resistance to methicillin. *Journal of Antimicrobial Chemotherapy*. 43: 15-22.
- **Weller, T.M.A.(2000):**Methicillin-resistant *Staphylococcus aureus* typing methods: which should be the international standard? *J. Hosp. Infect.*, 44:160-72.
- **Welsh, J. and McClelland, M. (1990):** Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res*; 18:7213-8.
- **Wenzel, R.P., and Per, T. (1995):** The significance of nasal carriage of *Staphylococcus aureus* and the incidence of postoperative wound infection. *J. Hosp. Infect.*,31:13-24.
- **Wertheim, H., Verbrugh, H.A, van Pelt,C., de Man, P., van Belkum, A. and Margreet C. (2001):**Improved detection of methicillin resistant *Staphylococcus aureus* using phenyl mannitol broth containing aztreonam and ceftizoxime *.J.Clin.Microbiol.*, P:2660-2662.
- **Whitener, C., Caputo, G., Weirekamp, M. and Karchmer, A.(1993):**Endocarditis due to coagulase negative staph: microbiology ,epidemiology and clinical consideration. *Infect .Dis. Clin. North Am.*7:81-96.
- <http://www.whosea.org/microbio/secbf.htm>

References

- **Wichelhaus, T.A., Kern, S., Schafer, V. and Brade, V. (1999):** Evaluation of modern agglutination tests for identification of methicillin –susceptible and methicillin –resistant *Staphylococcus aureus* .*Eur.J. Clin. Microbial, Infect .Dis* 18:756-758.
 - **Wielders, C. L. C., Fluit, A. C., Brisse, S., Verhoef, J. and Schmitz, F. J.(2002):** *mecA* Gene Is Widely Disseminated in *Staphylococcus aureus* Population. *Journal of Clinical Microbiology*, 40(11): 3970-3975.
 - **Williams J.G., Kubelik A.R. and Livak K.J.,(1990):** DNA polymorphisms amplified by arbitrary primers are useful genetic markers. *Nucleic Acids Res*; 18:6531-5.
 - **Wittwer, C.T., Reed, G.H., Gundry, C.N., Vandersteen, J.G. and Pryor, R.J. (2003):** High-resolution genotyping by amplicon melting analysis using LC Green. *Clin Chem*, 49: 853-860.
 - **Woods, G.L. and Yam, P.(1988):** Evaluation of MicroScan MIC panels for detection of oxacillin-resistant staphylococci. *J Clin Microbiol.*, 26(5): 816–820.
- Y:
- **Yamazumi , S.A.M., Wilke, W.W., Diekema, D.J. and Pfaller, M.A.(2001):** Comparison of the Vitek gram positive susceptibility 106card and the MRSA –Screen Latex agglutination test for determining oxacillin resistance in clinical bloodstream isolates of *Staphylococcus aureus* .*J.Clin.Micrbiol.*34:249-253.
 - **York, M.K., Gibbs, L., Chehab, F. and Brooks, G.F.(1996):** Comparison of PCR detection of *mec A* with standard susceptibility testing methods to determine methicillin resistance in coagulase –negative staphylococci. *J.Clin.Microbiol.*34:249-253.
 - **Yoshida, T., Kondo, N., Hanifah, Y.A. and Hiramatsu, K.(1997):** Combined use of ribotyping, PFGE and IS431 typing in

the discrimination of nosocomial strains of methicillin-resistant *Staphylococcus aureus*. *Microbiol. Immunol.* 41:687-95.

Z:

- **Zambardi, G., Reverdy, M. E. , Bland, S. , Bes, M. , Feney, J. and Fleurette, J.(1994):** Laboratory diagnosis of oxacillin resistance in *Staphylococcus aureus* by a multiplex-polymerase chain reaction assay. *Diagn. Microbiol. Infect. Dis.* 19:25-31.
- **Zhang, H.Z., Hackbarth, C.J., Chansky, K.M. and Chambers, H.F.(2001):** A proteolytic transmembrane signaling pathway and resistance to beta-lactams in staphylococci. *Science.* 291:1962–1965.
- **Zhou, L., Vandersteen, J., Wang, L., Fuller, T., Taylor, M., Palais, B. and Wittwer, C.T. (2004):** High-resolution DNA melting curve analysis to establish HLA genotypic identity. *Tissue Antigens*, 64: 156-164.
- **Zimmermann, B., Holzgreve, W., Wenzel, F. and Hahn, S. (2002):** Novel real-time quantitative PCR test for trisomy 21. *Clin Chem*, 48: 362-363.