

References

References

- ❖ *Adam, I.J; Mohamdee, M.O. and Martin, I.G.(1994) :*
Role of circumferential margin involvement in the local recurrence of rectal cancer.
Lancet 1994; 344: 707-11.
- ❖ *AJCC (1997):*
Colon and Rectum.
In : American Joint Committee on Cancer : AJCC Cancer Staging Manual.
Philadelphia,Pa: Lippincott-Raven publishers.
5th ed., 1997, pp: 83-99.
- ❖ *Anderberg, B.; Enblad, P. and Sjodahl, R. (1984):*
Recurrent rectal carcinoma after anterior resection and stapling.
Br J Surg 1984; 71: 98-100.
- ❖ *Astler, V. B. and Collier, F. A. (1954):*
The prognostic significance of direct extension of carcinoma of the Colon and Rectum. Ann. Surg. 1954; 119:846
- ❖ *Ault, G. and Castro, O. (1952):*
Clinical study of ligation of the inferior mesenteric artery in left colon resection.
Surg Gynecol Obstet 1952; 94: 223-8.
- ❖ *Bacon, H.; Dirbas, F. And Poncedeleon, F. (1958):*
Extensive lymphadenectomy and high ligation of the inferior mesenteric artery for carcinoma of the left colon and rectum.
Dis Colon Rectum 1958;1:457-65.

❖ ***Bacon, H. and Jackson, C. (1953):***

Visceral metastases from carcinoma of the distal colon and rectum.

Surgery 1953; 33: 495.

❖ ***Balslev, I. And Harling, H. (1983):***

Sexual dysfunction following operation for carcinoma of the rectum.

Dis Colon Rectum 1983; 26: 785-8.

❖ ***Baron, P.; Enker, W.E. and Zakowski, M. (1995):***

Immediate vs. salvage resection after local treatment for early rectal cancer.

Dis Colon Rectum. 1995; 38:177-181.

❖ ***Bentzen, SM.; Baslev, I. And Pedersen, M. (1988):***

A regression analysis of prognostic factors after resection of Duke's B and C carcinoma of the rectum and rectosigmoid. Does postoperative Radiotherapy change the prognosis.

Br J Cancer 1988; 58: 195-201.

❖ ***Beynon, J.; Foy, D.M. and Temple, L.N. (1986):***

The endosonic appearance of normal colon and rectum.

Dis Colon Rectum 1986; 29: 810-13.

❖ ***Beynon, J.; Mortenson, N.J. and Foy, D.M. (1989):***

Preoperative assessment of mesorectal lymph node involvement in rectal cancer.

Br J Surg 1989; 76: 276-285.

❖ ***Bishoff, J. (2002):-***

Laparoscopic Pelvic Lymph Node Dissection.

e-medicine, February 22-2002.

❖ ***Bokey, E.; Chapuis, P. and Hughes, W. (1984):***

Local recurrence following anterior resection for cancer rectum
with a stapled anastomosis.

Acta Chirurgica Scandinavica 1984; 150: 683-686.

❖ ***Bokey, E.; Chapuis, P. and Ojerkog, B. (1999):***

Local recurrence after curative excision of the rectum for cancer
Without adjuvant therapy: role of total anatomical dissection.

Br J Surg 86; 1164-1170.

❖ ***Bolkainy, N. (1991):***

Epidemiology of rectal cancer at NCI.

In: General Pathology of Cancer.

1st ed., 1991; Al Asedekaa graphic center.

❖ ***Brown, C.E. and Warren, S. (1938):***

Visceral metastases from rectal carcinoma.

Surg Gynecol Obstet 66, 611.

❖ ***Burgos, F.J.; Romero, J. and Fernandez, E. (1988):***

Risk factors for developing voiding dysfunction after
abdominoperineal resection for adenocarcinoma of the
rectum.

Dis Colon Rectum 1988; 31: 682 – 85.

❖ ***Carlsen, E.; schlichting, E. and heald, R. J. (1998):***

Effect of introduction of total mesorectal excision for the surgical treatment of cancer rectum.

Br J Surg 1998; 85: 526-529.

❖ ***Cawthorn, S.J.; Parums, D.V. and Gibbs, N.M. (1990):***

Extent of mesorectal spread and involvement of lateral resection margin as prognostic factors after surgery for rectal cancer.

Lancet 1990; 335: 1055-1059.

❖ ***Cedermark, B.J.; Shultz, S.S. and Bakshi, S. (1977):***

The value of liver scan in the follow-up study of patients with adenocarcinoma of the colon and rectum.

Surg Gynecol Obstet 1977; 144: 745.

❖ ***Choe, J.M.; Talavera, F. and Dan, T. (2002):-***

Lymph Node Dissection, pelvic.

e-medicine, January 25-2002.

❖ ***Conte, M.; Panici, B. and Guariglia, L. (1990):-***

Pelvic lymphocele following radical para-aortic and pelvic lymphadenectomy for cancer cervix.

Obestet Gynecol. 76: 268.

❖ ***Coller, F.; Kay, E. and MacIntyre, R. (1940):***

Regional lymphatic metastases of carcinoma of the rectum.
Surgery 1940; 8: 294-311. (Quoted from Goligher, 1984).

❖ ***Cosin, J.A.; Fowler, J.M. And Chen, M. (1998):-***

Pretreatment staging of patients with cancer cervix.
Cancer 83:2241.

❖ ***Creasman, W.; Morrow, C. And Bundy, B. (1987):-***

Surgical pathological spread pattern of endometrial carcinoma.
Cancer 60: 2035.

❖ ***Crespi, M. and caperle, M. (1991) :***

Trends in epidemiology of colorectal cancer.
J. Surg. Oncology Suppl., 2: 1-3.

❖ ***Curely, S.; Skibber, J and Key, A. (1996):-***

Survival factors in 186 patients younger than 40 years old
With colorectal adenocarcinoma.
J Am Coll Surg 183: August 1996.

❖ ***Danzi, M.; Ferulano, G. P. And Abate, S. (1983) :***

Male sexual function after abdominoperineal resection for
rectal cancer.
Dis Colon Rectum 1983; 26: 665-8.

❖ ***Dawson, P.M.; Habib, N.A. and Rees, H.C. (1987):***

Influence of sialomucin at the resection margin on local tumour
recurrence and survival of patients with colorectal cancer: a
multivariate analysis.
Br J Surg 1987; 74: 366- 369.

- ❖ *Decosse, J.; Wong, R. and Quan, S. (1989) :*
 Conservative treatment of distal rectal cancer by local excision.
 Cancer 1989; 63: 219-23.

- ❖ *Deddish, M. (1951):*
 Abdominopelvic lymph node dissection in cancer of the rectum and distal colon.
 Cancer 1951; 4: 1364-6.

- ❖ *Di-matteo, G.; Mascagni, D. and Tarroni, D. (1991):*
 Radical surgery for cancer rectum.
 J surg oncol supp. 2:32-35.

- ❖ *Douglas, E. and Arnold, M. (1984):*
 Cancer of the Colon and Rectum.
 In: Surgical Oncology.
 Edited by Yosef, H. P. Hamburg. Tokyo. PP: 596- 607.

- ❖ *Dukes, C. E. (1930):*
 The spread of cancer rectum.
 Br J Surg 1930; 17: 643. (Quoted from Goligher, 1984).

- ❖ *Dukes, C. E. (1940):*
 Carcinoma of the rectum: an analysis of 1000 cases.
 J. Path. Bact. 1940; 50: 527. (Quoted from Goligher, 1984).

❖ ***Duncan, J.L. and Kyle, J. (1982):***

Family incidence of carcinoma of the colon and rectum.
Gut 23,169.

❖ ***Dunstone, G.H. and Knaggs, T.(1972):***

Familial cancer of the colon and rectum.
J. Med. Genet.,9,451.

❖ ***Dwight, R.; Higgin, G. And Keehn, R. (1969):***

Factors influencing survival after resection in cancer of the
Colon and rectum.
Am. J. Surg. 1969; 117: 512-22.

❖ ***Dworak, O. (1991):-***

Morphology of lymph nodes in the resected rectum of patients
with rectal carcinoma.
Pathol Res Pract 1991; 187:1020- 1024.

❖ ***Enker, W.E(1995):-***

Total Mesorectal excision in the operative treatment of rectal
Cancer.
J Am Coll Surg 1995, 181: 335-346.

❖ ***Enker, W.; Havenga, K. and Cohen, A. (1996):***

Male and Female sexual and urinary function after total
Mesorectal excision with autonomic nerve preservation for
carcinoma of the rectum.
J Am Coll Surg June 1996. 182: 495-502.

❖ **Enker, W.; Pilpshen, S. and Heilweil, M.(1986):**

En bloc pelvic lymphadenectomy and sphincter preservation in the surgical management of rectal cancer.

Ann Surg 1986; 203: 426-33.

❖ **Enker, W.; Urban, L. and Block, G. (1979):**

Enhanced survival of patients with colic and rectal cancer is Based upon wide anatomic resection.

Ann surg 1979; 190: 350-60.

❖ **Ferguson, L.; Boland, J. and Thomen, F. (1962):**

Anterior segmental resection for carcinoma of the upper rectum, rectosigmoid and sigmoid.

Surgery 1962; 52: 741-6.

❖ **Gabriel, W. B. (1948):**

The Principles and Practice of Rectal Surgery.

4th ed, P. 309. London: H. K. Lewis. (Quoted from Goligher, 1984).

❖ **Gabriel, W. B.; Dukes, C. and Bussey, H.(1935):**

Lymphatic spread in cancer of the rectum.

Br J Surg 1935; 23: 395-413. (Quoted from Goligher,1984).

❖ **Ganong, W.F. (1993):**

Autonomic nervous system.

In: Review of medical physiology.

Edited by Appleton Lange, 16th ed. pp: 201-208.

- ❖ *Gilbertsen, A. (1962):-*
The result of the surgical treatment of rectal cancer.
Surg Gynecol. Obstet. 1962; 114: 313- 319.
- ❖ *Gilchrist, R. (1940):*
Fundamental factors governing lymphatic spread of carcinoma.
Ann Surg 1940; 111: 630-9.
- ❖ *Gilchrist, R. and David, V. (1938):*
Lymphatic spread of carcinoma of the rectum.
Ann Surg 1938; 108: 621-37. (Quoted from Goligher,1984).
- ❖ *Gilchrist, R. and David, V. (1947):*
A consideration of pathological factors influencing five- year
survival in radical resection of the large bowel and rectal
cancers.
Ann Surg 1947; 126: 421- 38.
- ❖ *Gilchrist, R. and David, V. (1948):*
Prognosis in carcinoma of the bowel.
Surg Gynecol Obstet 1948; 86: 359-71.
- ❖ *Glaser, F.; Kunt, Z. and Herfarth, C. (1993):*
Endorectal ultrasound for control of preoperative radiotherapy
of rectal cancer.
Ann Surg 1993; 86: 359-71.

❖ **Glaser, F.; Layer, G. and Zuna, I. (1990):**

Preoperative beurteilung pararectaler lymphknoten durch
Ultraschall.

Chirurg 1990; 61: 587-91.

❖ **Glass, R.; Ritchie, J. and Thompson, H.(1985):**

The results of surgical treatment of cancer of the rectum by
radical resection and extended abdomino-iliac lymph-
adenectomy.

Br J Surg 1985; 72: 599-601.

❖ **Glover, R. and Waugh, J. (1945):**

Retrograde lymphatic spread of carcinoma of the rectosigmoid
Region; its influence on surgical procedures.

Surg Gynecol Obstet 1945; 80: 434.

❖ **Goligher, J.C. (1984):**

Surgery of the anus, rectum and colon.

5th ed london bailliere tindall, 1984.

❖ **Goligher, J.C.; Dukes, C.E. and Bussey, H. (1951):**

Local recurrence after sphincter-saving excision for carcinoma
of the rectum and rectosigmoid.

Br J Surg 1951; 39: 199.

❖ **Greaney, M. and Irvin, T. (1977):**

Criteria for the selection of rectal cancers for local treatment :

A clinicopathological study of low rectal cancers.

Dis Colon Rectum 1977; 20: 463-6.

- ❖ *Gregory, P.M. and Barry, W.F (1999):*
Cancer of the colon rectum and anus.
In: The M.D. Anderson Surgical Oncology. Hand book.
Edited by lippincott, Philadelphia. London. 2nd ed. Pp: 185.

- ❖ *Grinnell, R. (1939):*
The grading and prognosis of carcinoma of the colon and rectum.
Br J Surg 1939; 109: 500. (Quoted from Goligher, 1984).

- ❖ *Grinnell, R. (1942):*
The lymphatic and venous spread of carcinoma of the rectum.
Ann Surg 1942; 116: 200-16.

- ❖ *Grinnell, R. (1950):*
Lymphatic metastases of carcinoma of the colon and rectum.
Ann Surg 1950; 131: 494-506.

- ❖ *Grinnell, R. (1965):*
Results of ligation of the IMA at the aorta in resections for carcinoma of the descending and sigmoid colon and rectum.
Surg Gynecol Obstet 1965; 120: 1031-6.

- ❖ *Grinnell, R. (1966):*
Lymphatic block with atypical and retrograde lymphatic metastases and spread in carcinoma of the colon and rectum.
Ann Surg 1966; 163: 272-80.

❖ *Grinnell, R. and Hiatt, R. (1952):*

Ligation of the IMA at the aorta in resections for carcinoma of the sigmoid and rectum.

Surg Gynecol Obstet 1952; 94: 526-34.

❖ *Havenga, K.; Enker, W.E. and Weivart, K. (1996):*

Anatomic bases of autonomic nerve preserving total mesorectal Excision for rectal cancer.

Br J Surg 1996; 83: 384-388.

❖ *Harnesberger, J. R.; Vernara, A.M. and Long, W.E. (1994):*

Radical Abdominopelvic Lymphadectomy: Historic Perspective and Current Role in the Surgical Management of rectal cancer.

Dis Colon Rectum 1994; 37: 73-87.

❖ *Harvey, H. and Auchincloss, H. (1968):*

Metastases to lymph nodes from carcinoma that were arrested.

Cancer 1968; 21:684-91.

❖ *Heald, R.J. (1986):*

Recurrence and survival after total mesorectal excision for rectal cancer.

Br J Surg 1986; i: 1479-1482.

❖ *Heald, R.J. (1995):*

Rectal cancer: The surgical options.

Eur. J. Cancer 1995; 31A: 1189-1192.

- ❖ *Heriot, A.G.; Grundy, A. and Kumar, D. (1999):*
Preoperative staging of rectal carcinoma.
Br J Surg 1999; 86: 17-28.

- ❖ *Herzog, U.; Von flue, M. and Tondelli, P.(1993):*
How accurate is Endorectal Ultrasound in the preoperative
Staging of Rectal Cancer.
Dis Colon Rectum 1993; 36:127-134.

- ❖ *Hildebrandt, U. and Fiefel, G. (1985):*
Preoperative staging of rectal cancer by intrarectal ultrasound.
Dis Colon Rectum.1985; 28: 42-48.

- ❖ *Hida, J.; Yastomi, M. and Tokoro, T. (1999):-*
Examination of nodal metastasis by clearing method supports
pelvic plexus preservation in rectal cancer surgery.
Dis Colon Rectum 42; 510- 514.

- ❖ *Hojo, K. (1981):*
Preservation of micturition and sexual functions in the surgery
of rectal cancer.
Igaku no ayumi 1981; 119: 716-23.

- ❖ *Hojo, K.; Anthony, M. and Katumata, K. (1991):*
 Preservation of urine voiding and sexual function after rectal cancer surgery.
 Dis Colon Rectum 1991; 34: 532-539.

- ❖ *Hojo, K. and Koyama, Y. (1982) :*
 The effectiveness of wide anatomical resection and radical lymphadenectomy for patients with rectal cancer.
 Jpn J Surg 1982; 12: 111-17.

- ❖ *Hojo, K.; Koyama, Y. and Moriya, Y. (1982):*
 Lymphatic spread and its prognostic value in patients with rectal cancer.
 Am J Surg 1982; 144: 350- 4.

- ❖ *Hojo, K.; Sawada, T. and Moriya, Y. (1989):*
 An analysis of survival and voiding , sexual functions after wide iliopelvic lymphadenectomy in patients with carcinoma of the rectum ,compared with conventional lymphadenectomy.
 Dis Colon Rectum 1989; 32: 128-33.

- ❖ *Horn, A.; Dahl, O. and Morild, I. (1990):*
 The role of venous and neural invasion on survival in rectal adenocarcinoma.
 Dis Colon Rectum 1990; 33: 598-601.

- ❖ *Horn, A.; Dahl, O. and Morild, I. (1991):*
 Venous and Neural Invasion as predictors of Recurrence in Rectal cancer.
 Dis Colon Rectum 1991; 34: 798-804.

❖ ***Jamieson, J. K. and Dobson, J. F. (1909):***

Lectures on the lymphatic system of the caecum and appendix.

Lancet, 1, 1137. (Quoted from Goligher, 1984)

❖ ***Jinnai, D (1982):***

Personal communications; on work by him and his colleagues;
in preparation.

❖ ***Katsura, Y.; Yamada, K. and Ishizawa, T.(1992):***

Endorectal Ultrasonography for the assessment of wall invasion
and lymph node metastasis in Rectal Cancer.

Dis Colon Rectum 1992; 35: 362-368.

❖ ***Kettlewell, M.G.W. (1994):***

Colorectal cancer and benign tumours of the colon.
Oxford textbook of surgery.

Edited by Morris, P.J. and Malt, R.A. pp: 1060-1073.

❖ ***Kodner, I. J. (1997):***

Rectal cancer.

In: Maingot's Abdominal Operations.

Edited by Zinner, M.J. Boston, 10th ed. pp: 1455-78.

❖ ***Koyama, Y.; Moriya, Y. and Hojo, K. (1984):***

Effect of extended systemic lymphadenectomy for
adenocarcinoma of the rectum-significant improvement of
survival rate and decrease in local recurrence.

Jpn J. Clin. Oncol. 1984; 14: 623-32.

❖ ***Leggeri, A.; Roseano, M. and Balani, A. (1994):***

Lumboaortic and Iliac Lymphadenectomy: What is the Role Today.

Dis Colon Rectum 1994; 37(suppl): s54- s61.

❖ ***Lovett, E. (1976):***

Familial cancer of the gastrointestinal tract.

Br J Surg 1976; 63: 19.

❖ ***Lynch, H.; Harris, R. and Organ, C. (1978):***

Familial cancer: indication of surgical management in high risk patients.

Surgery, 83: 104.

❖ ***Makela, J.; Hankipuro, K and Laitinen, S. (1989):-***

Surgical treatment for recurrent cancer rectum.

Arch surg 124; 1029-1032.

❖ ***MacFarlane, J.K.; Ryall, R.D.H. and Heald, R.J. (1993):***

Mesorectal excision for rectal cancer.

Lancet 1993; 341:457-460.

❖ ***Marcio, J.; Wexer, J and Maga, N. (1997):-***

Anatomy and physiology of the rectum and anus.

Eur. J Surg 1997; 163:723-731.

- ❖ *Mascagin, D.; Hojo, K. and Moriya, Y. (1992):*
 Pattern of lymph node metastases from low rectal cancer
 (meeting abstr).
 Dis Colon Rectum 1992; 35: p6.

- ❖ *Mathis, M. (1962):*
 Familial carcinoma of the colon, a family tree from the canton
 of Argau.
 Schweiz. Med. Schr., 92, 1673.

- ❖ *Mc Call, J. (1995):*
 Analysis of local recurrence rates after surgery alone for cancer
 rectum.
 J colorectal Dis 1995; 126: 132.

- ❖ *McElwain, J.; Bacon, H. and Trimpi, H. (1954):*
 Lymph node metastases: Experience with aortic ligation of the
 IMA in cancer of the rectum.
 Surgery 1954; 35: 513-31.

- ❖ *Michelassi, F. and Block, G. (1992):*
 Morbidity and mortality of wide pelvic lymphadenectomy for
 rectal carcinoma.
 Dis Colon Rectum 1992; 35: 1143-50.

- ❖ *Michelassi, F. and Block, G. and Vannucci, L. (1988):*
 A 5-to 21- year follow up and analysis of 250 patients with
 rectal adenocarcinoma.
 Ann Surg 1988; 208:379-89.

❖ *Michelassi, F.; Vannacci, L. and Goldberg, R. (1990):-*

Local recurrence after curative resection for colorectal adenocarcinoma.

Surgery 108; 787-793.

❖ *Miles, W.E. (1908):*

A method of performing Abdominopelvic excision for carcinoma of the rectum and of the terminal portion of the pelvic colon.

Lancet, 1908; 2: 1812-15. (Quoted from Goligher, 1984).

❖ *Miles, W.E. (1910):*

The radical abdomino-perineal operation for cancer of the rectum and the pelvic colon.

Br Med J 1910; 2: 941. (Quoted from Goligher, 1984).

❖ *Miles, W.E. (1926):*

Cancer of the rectum.

London: Harrison (Quoted from Goligher, 1984).

❖ *Moore, K. (1992):-*

The abdomen and pelvis.

In: Clinically Oriented Anatomy.

Edited by Timothy S. Satterfield. 3rd edition.

❖ *Moreira, Y. (1997):*

Importance of extended lymphadenectomy with lateral node dissection for advanced lower rectal cancer.

World J Surgery 1997, 21(7): 728-732.

❖ ***Moreira, L.; Hizuta, A. and Iwagaki, H. (1994):***

Lateral lymph node dissection for rectal carcinoma below the peritoneal reflection.

Br J Surg 1994; 81: 293-296.

❖ ***Morgan, N. and Griffiths, D. (1959):***

High ligation of the IMA during operations for carcinoma of the distal colon and rectum.

Surg Gynecol Obstet 1959; 108: 641-50.

❖ ***Morrow, P. and Curtin, J. (1996):-***

Surgery for cervical neoplasm.

Gynecological cancer surgery, New York Churchill Livingstone, 1996.P: 522.

❖ ***Morson, B.C. and Dawson, I.M. (1979):***

Prevention of colorectal cancer.

Gastrointestinal Pathology.

2nd edition. Oxford: Blackwell scientific.

❖ ***Morson, B.C.; Vaughan, E.G. and Bussey, H.J. (1963):***

Pelvic recurrence after excision of the rectum for carcinoma.

Br Med J 1963; 2: 13.

❖ ***Morikawa, E. Yasutomi, M. and Shindou, K. (1994):-***

Distribution of Metastatic lymph Nodes in Colorectal Cancer by The Modified Clearing Method.

Dis Colon Rectum 1994;37:219-223.

❖ ***Moriya,Y.; Hojo,K. and Koyama,Y. (1989):***

Significance of lateral node dissection for advanced rectal carcinoma at or below the peritoneal reflection.

Dis Colon Rectum 1989; 32: 307-15.

❖ ***Moynhan, B. (1908):***

The surgical treatment of cancer of the sigmoid flexure and rectum.

Surg Gynecol Obstet 1908; 6: 463-6. (Quoted from Harnsberger et al. 1994).

❖ ***Northover,J.M. and Scholefield,J.H.(1995):***

Surgical management of rectal cancer.

Br. J Surg. 1995;82:745-748.

❖ ***Orr,J.; Holimon, J. and Orr,P.(1997):-***

Stage I corpus cancer: is teletherapy necessary?

Am J Obstet Gynecol 176:777.

❖ ***Orr,J.; Holimon, J. and Orr,P.(1997):-***

Stage I corpus cancer: is teletherapy necessary?

Am J Obstet Gynecol 176:777.

❖ ***Paker, S.L.; Tong, T. and Bolder, S.(1996):***

Cancer statistics, 1996.

CA Cancer J Clin 1996; 46: 5-27.

❖ ***Panici,B.; Maneschi,F. and Baiocchi ,G. (1993):-***

Radical hysterectomy.

Gynecol Oncol 50: 226.

- ❖ ***Panici,B.; Maneschi,F. and Catello,G. (1997):-***
 Lymphadenectomy in early epithelial ovarian cancer.
 Int J Gynecol. Cancer 7 (suppl):17.

- ❖ ***Panici,B.; Maneschi,F. and Cutillo,G. (2001):-***
 Pelvic and Aortic Lymphadenectomy.
 S Clin. North America; Gynecological oncology.
 volume 81: 4; 841-858. Augst 2001.

- ❖ ***Panici,B.; Maneschi,F. and D'Andrea,G. (2000):-***
 Early cervical cancer, role of lymphadenectomy.
 Cancer 88; 2267.

- ❖ ***Panici,B.; Maneschi,F. and Scambia,G. (1996):-***
 Lymphatic spread of cervical cancer.
 Gynecol Oncol 62:19.

- ❖ ***Peyster, F.A. and Gilchrist, R.K. (1982):-***
 Pathology of cancer of the colon and rectum.
 S Clin. North America. 1295-305.

- ❖ ***Pezim, M. and Nicholls, R. (1984):-***
 Survival after high and low ligation of the IMA during curative
 surgery for rectal cancer.
 Ann Surg 1984; 200: 729-33.

- ❖ ***Pfeiffer, D. and Miller, D.(1950):-***
 Cancer of the rectum; 5 and 10 year follow study of cases of
 cancer below the peritoneal reflection.
 Surg Gynecol Obstet 1950; 91: 319-22.

- ❖ ***Pilipshen, S.; Heilweil, M. and Quan, S. (1984):***
 Pattern of pelvic recurrence following definitive resection of rectal cancer.
 Cancer 1984; 53: 1354- 62.

- ❖ ***Poirier, P.; Cuneo, B. and Delamere, G. (1903):***
 The lymphatics.
 London, p: 351, constable. (Quoted from Goligher, 1984).

- ❖ ***Quentmeier, A.; Schlag, P. and Smok, M. (1990):-***
 Re-operation for recurrent colorectal adenocarcinoma: the importance of early diagnosis for respectability and survival.
 European J Surg Oncol 16; 319-325.

- ❖ ***Quirke, P.; Durdey, P. and Dixon, M. (1986):***
 Local recurrence of rectal adenocarcinoma due to inadequate surgical resection: histopathological study of lateral tumour spread and surgical excision.
 Lancet, 1986; 2: 996-8.

- ❖ ***Raghaviah, N.V. and Jordan, W. P. Jr. (1979):***
 Prostatic lymphography.
 J. Urol 1979; 121: 178-183.

- ❖ ***Ratto, G.; Sofo, L. and Merico, M. (1999):***
 Accurate lymph node detection in colorectal specimens resected for cancer is of prognostic significance.
 Dis Colon Rectum 1999; 43: 143-58.

❖ ***Robert, W. and Beart, Jr. (1995);***

Colon and Rectum.

In: clinical oncology.

Edited by Martin, D. Churchill living stone, London.

PP: 1267-1272.

❖ ***Robin, P (1992):***

A multidisciplinary approach for physicians and student.

In: Clinical oncology.

7th edition, 1992 philadelphia, saunders, pp: 576-587.

❖ ***Romanes, G.J. (1987):-***

Thorax and Abdomen.

In: Cunningham's Manual of practical anatomy.

Edit. by Oxford university press., 5th edition. PP:179.

❖ ***Rosi, P.; Cahill, W. and Carey, J. (1962):***

A ten year study of hemicolectomy in treatment of carcinoma of the left half of the colon.

Surg Gynecol Obstet 1962; 114: 14-24.

❖ ***Sack, J and Rothman, J (2000):-***

Colorectal cancer. Natural history and Management.

Hospital physician (internet). October 2000; p: 64-73.

❖ ***Salvat, J. (2003):-***

Endoscopic lymphadenectomy.

Edited by Aldo Compana, March 13.2003.

❖ *Satoh, N.; Ihara, M. and Sarashins, H. (1989):*

Studies of diagnosis of rectal cancer using CT, MRI, and intrarectal ultrasonography.

Rinsbo hosbasen 1989; 34: 573-581.

❖ *Sauer, I and Bacon, H. (1951):*

Influence of lateral spread of cancer of the rectum on radicality of operation and prognosis.

Am J Surg 1951; 81: 111- 20.

❖ *Sauer, I and Bacon, H. (1952):*

A new approach for excision of carcinoma of the lower portion of the rectum and anal canal.

Surg Gynecol Obstet 1952; 95: 229-42.

❖ *Sauer, I and Bacon, H. (1958):*

Surgical treatment of cancer of the bowel.

Cancer 1950; 3 : 773-8.

❖ *Schulten, M.; Heiskeli, C. and Shields, T. (1976):*

The incidence of solitary pulmonary metastases from carcinoma of the large intestine.

Surg Gynecol Obstet 1979; 143: 727.

❖ *Schardey, H.M. (1991):*

Intraluminal vs. extraluminal local recurrence of colorectal carcinoma. Absence of predisposing histopathological factors after RO resection. ACA(suppl.) 93: 21-22.

❖ *Scott, N.; Jackson, P. and Al-Jaberi, T. (1995):*

Total mesorectal excision and local recurrence: A study of tumour spread in the mesorectum distal to rectal cancer.

Br J surg 1995; 82:1031-1033.

❖ *Scott, N.; Rainwater LM. and wieand, HS.(1987):*

The relative prognostic value of flow cytometric DNA analysis and conventional clinicopathological criteria in patients with operable rectal cancer.

Dis Colon Rectum 1987; 30: 513-320.

❖ *Seow- Choen,F.; EU.K-W,and Ho,J.M.S. (1998):*

Local recurrence following rectal resection for cancer.

J.R. Coll. Surg. Edinb. 43. December, 1998, 393-396.

❖ *Shand, w. s. (1981):*

Surgical anatomy and physiology of the rectum and anal canal.

In: operative surgery and manegement.

Edit. by G.Keen Bristol-London, Boston.1981: 171-173.

❖ *Sholefield, J.H. and Northover, J.M.(1995):-*

Surgical management of rectal cancer.

Br J Surg 1995; 82: 745-748.

❖ *Sitzler,P.; Choen, F. and Ho,Y. (1997):-*

Lymph node involvement and tumour depth in rectal cancers.

An analysis of 805 patients.

Dis Colon Rectum 1997; 40: 1472- 1476.

❖ ***Soliman, A.; Bondy, M. and Hamza, R. (1997):***

Colorectal cancer in Egyptian patients under 40 years of age.
Int. J. Cancer 1997; 71(1): 26-30.

❖ ***Soloman, MJ. and Mcleod, R.S. (1993):***

Endoluminal transrectal ultrasonography: accuracy, reliability,
and validity.
Dis Colon Rectum 1993;36: 200-205.

❖ ***State, D. (1951):***

Combined abdominoperineal excision of the rectum-a plan for
standarization of the proximal extent of dissection.
Surgery 1951; 30: 349-54.

❖ ***Stelle, G.; Ravikuma, S. and Benotti, P. (1990):-***

New surgical treatment for recurrent colorectal cancer.
Cancer 65; 723-730.

❖ ***Stipa, S.; Nicolani, V. and Botti, C. (1991):-***

Local recurrence after curative resection of rectal cancer:
frequency, risk factors and treatment.
J surg oncol. Supp. 2; 155-160.

❖ ***Streans, M. and Deddish, M. (1959):***

Five year results of abdominopelvic lymph node dissection
for carcinoma of the rectum.
Dis Colon Rectum 1959; 2: 169-72.

❖ ***Sugar, P.M. and Pemberton, J.H. (1996):-***

Surgical management of locally recurrent cancer rectum.
Br J Surg 1996, 83; 293-304.

❖ ***Takahashi, T. and Kajitani, T. (1983):-***

Some consideration on the lateral lymphatic metastases from
rectal cancer.
Personal communications.

❖ ***Talbot, I.C.; Ritchie, S. and Leighton, M. (1980):-***

The clinical significance of invasion of veins by rectal cancer.
Br J Surg 1980; 67: 439.

❖ ***Tang, R.; Wang, J. and Chen, J. (1995):-***

Survival impact of lymph node metastasis in TNM stage III
carcinoma of the colon and rectum.
J AM Coll Surg 1995; 180: 705- 712.

❖ ***Todd, O.R. (1996):-***

The Abdomen and Pelvis.
In: A.D.A.M. Student Atlas of Anatomy.
Edited by Williams and Wilkins, Philadelphia, London. 3rd ed.

❖ ***Tschmelitsch, J; Kronberger, P. and Glaser, K. (1994):-***

Survival after surgical treatment of recurrent carcinoma of the
rectum.
J Am Coll Surg 1994; 179: 54-58.

❖ ***Tsuchiya, S. (1982):***

Autonomic nerve preservation in cancer rectal surgery.

Surgery 1982; 44: 1367-73.

❖ ***Turnbell,R.; Kyle, K. and Watson, F. (1967):***

Cancer of the colon: the influence of non touch isolation technique on survival rates.

Ann Surg 1967; 166: 420-427.

❖ ***Villemin, F.; Huard, P. and Montagne, M. (1925):***

Recheres anatomiques sur les lymphatiques du rectum et de l'anus; leurs applications dans le traitement chirurgical du cancer.

Rev. Chir. Paris 1925; 63: 39. (Quoted from Goligher, 1984).

❖ ***Walsh, P.C. and Schlegel, P. (1988):***

Radical pelvic surgery with preservation of sexual function.

Ann Surg 1988; 208: 391-400.

❖ ***Walsh, P.C. and Schlegel, P. N. (1991):***

Radical cystectomy.

In: Operative urology.

Edited by Marshall. F. Saunder company. Toronto. Pp: 132-137.

❖ ***Waugh, J. and Kinklin, J. (1949):***

The importance of the level of the lesion in the prognosis and treatment of carcinoma of the rectum and low sigmoid colon.

Ann Surg 1949; 129: 22-3.

❖ ***Westhues, H.(1934):***

Die pathologisch-anatomischen Grundlagen des
Rectumkarzinoms.

Leipzig: Thieme. (Quoted from Goligher, 1984).

❖ ***Williams, N.S.; Dixon, M.F. and Johnston, D. (1983):-***

Reappraisal of the 5 cm rule of distal excision for carcinoma of
the rectum: a study of distal intramural spread and patients
survival.

Br J Surg 1983; 70: 150-154.

❖ ***Willis, R.A. (1948):***

The pathology of tumours, pp. 422-425. London : Butterworths.

❖ ***Wilmink, A.B. (1997):***

Overview of the epidemiology of colorectal cancer.

Dis Colon Rectum 1997; 40: 483-493.

❖ ***Wood, W.Q. and Wilkie, D.P. (1933):***

Carcinoma of the rectum. An anatomico-pathological study.

Edinb. Med. J., 1933; 40: 321. (Quoted from Goligher, 1984).

❖ ***Yasutomi, M.(1997):***

Advances in rectal cancer surgery in Japan.

Dis colon rectum. 1997, 40: No., 10 (suppl).