



REFERENCES



REFERENCES

Adam G. and Schwartz, B. (1980):

Increased fluorescein filling defects in the wall of the optic disc cup in glaucoma.

Arch. Ophthalmol 98 : 1590-1592.

Anderson, D. R. (1969):

Ultrastructure of human and monkey lamina cribrosa and optic nerve head.

Arch. Ophthalmol 82: 800-814.

Anderson, D. R. and Hoyt, W. F. (1969):

Ultrastructure of the intraorbital portion of human and monkey optic nerve.

Arch. Ophthalmol 82 : 506.

Anderson, D. R. (1970):

Vascular supply to the optic nerve of primates.

Am. J. Ophthalmol 70: 341-351.

Anderson, D. R. and Hendrickson, A. (1974):

Effect of intraocular pressure on rapid axoplasmic transport in monkey optic nerve.

Invest. Ophthalmol 13 : 771-783.

Anderson, D. R. and Braverman, S. (1976):

Re-evaluation of the optic disc vasculature.
Am. J. Ophthalmol. 82 : 165-174.

Becker-Shaffer (1989):

Diagnosis and therapy of glaucomas.
The C.V. Mosby Company.

Beauvieux, J. and Ristitch, K. (1924):

Les vaisseaux centraux du nerf optique.
Arch. Ophthalmol 41 : 352.
Quoted from : François J. and Neetens A. (1954):
Vascularization of the optic pathway. I-Lamina cribrosa
and optic nerve.
Br. J. Ophthalmol 38: 472-488.

Best, M.; Blumenthal, M.; Galin, M. A. and Toyofuku, H. (1972):

Fluorescein angiography during induced ocular
hypertension in glaucoma.
Br. J. Ophthalmol 56: 6-12.

Best, M. and Toyofuku, H. (1972):

Ocular hemodynamics during induced ocular

hypertension in man.

Am. J. Ophthalmol 74 : 932-939.

Blumenthal, M.; Gitter, K. A.; Best, M. and Galin, M. A. (1970):

Fluorescein angiography during induced ocular hypertension in man.

Am.J. Ophthalmol 69: 39-43.

Blumenthal, M.; Best, M. and Galin, M. A. (1971 a):

Ocular circulation analysis of the effect of induced ocular hypertension on retinal and choroidal blood flow in man.

Am. J. Ophthalmol 71 : 819-825.

Blumenthal, M.; Best, M.; Galin, M.A. and Toyofuku, H. (1971 b):

Peripapillary choroidal circulation in glaucoma.

Arch Ophthalmol 86: 31-38.

Cohen, A. I. (1973):

Is there a potential defect in the blood-retinal barrier at the choroidal level of the optic nerve canal.

Invest. Ophthalmol. 12 : 513.

Cristini, G. (1951):

Common pathological basis of the nervous ocular

symptoms in chronic glaucoma.

Br. J. Ophthalmol. 35 : 11.

Drance, S. M. (1972):

Some factors in the production of low tension glaucoma.

Br. J. Ophthalmol 56: 229-242.

Emery, J. M.; Landis, D.; Paton, D.; Boniuk, M. and Craig, J.M.

(1974):

The lamina cribrosa in normal and glaucomatous human eyes.

Trans. Am. Acad Ophthalmol Otolaryngol. 78: 290-296.

Ernest, J. T. and Potts, A.M. (1968):

Pathophysiology of the distal portion of the optic nerve. 2-

Vascular relationships.

Am. J. Ophthalmol 66: 380.

Fishbein, S. L. and Schwartz, B. (1977):

Optic disc in glaucoma. Topography and extent of

fluorescein filling defects.

Arch Ophthalmol. 95: 1975-1979.

François, J. and Neetens, A. (1954):

Vascularization of the optic pathway.

I-Lamina cribrosa and optic nerve.

Br. J. Ophthalmol. 38: 472-488.

François, J.; Neetens, A. and Colette, J. M. (1958):

Vascularization of the primary optic pathways.

Br. J. Ophthalmol 42: 65.

François J. and Neetens, A. (1963):

Central retinal artery and central optic nerve artery.

Br.J. Ophthalmol. 46: 21.

François J. and Neetens, A. (1964):

Vascularity of the eye and the optic nerve in glaucoma.

Arch. Ophthalmol 71: 219-225.

François, J. and Neetens, A. (1965):

Vascularization of the intraorbital part of the optic nerve.

Am.J. Ophthalmol 60: 62-67.

Gillian, L. A (1961):

The collateral circulation of the human orbit.
Arch. Ophthalmol 65: 684.

Harrington, D. O. (1959):

The pathogenesis of the glaucoma field.
Am. J. Ophthalmol. 47: 177-184.

Harrington, D. O. (1964):

The visual fields 2nd ed.
p.p. 109, 136, 137, 155 Mosby, St Louis.

Hayreh, S. S. (1962):

The ophthalmic artery. III. Branches.
Br. J. Ophthalmol 46: 212-247.

Hayreh, S. S. (1969):

The blood supply of the optic nerve head and its role in
optic atrophy, glaucoma and oedema of the optic disc.
Br. J. Ophthalmol, 53: 721-748.

Hayreh, S. S. (1970):

Pathogenesis of visual field defects.

Role of ciliary circulation.

Br. J. Ophthalmol 54: 289-311.

Hayreh, S. S. (1972):

Colour and fluorescence of the optic disc.

Ophthalmologica 165: 100-108.

Hayreh, S. S. and Dass, R. (1960):

The central artery of the retina.

2-study of its distribution and anastomoses.

Br. J. Ophthalmol. 44: 280.

Hayreh, S. S. and Walker, W. M. (1967):

Fluorescent fundus photography in glaucoma.

Am. J. Ophthalmol 63: 982-989.

Hayreh, S. A.; Revie, I.H.S. and Edwards, J. (1970):

Vasogenic origin of visual field defects.

Br. J. Ophthalmol 54: 461-472.

Hayreh, S.S. and Baines, J.A.B. (1972):

Occlusion of the posterior ciliary artery. III. Effect on the optic nerve head.

Br. J. Ophthalmol 56: 754-764.

Henkind, P. and Levitzky, M. (1969):

Angioarchitecture of the optic nerve. I-The papilla.
Am. J. Ophthalmol 68: 979-986.

Hitchings, R. A. and Spaeth, G. L. (1977):

Fluorescein angiography in chronic simple and low-tension glaucoma.
Br. J. Ophthalmol 61: 126-132.

Hoyt, W. F. and Tudor, R. C. (1963):

The course of peripapillary temporal retinal axons through the anterior optic nerve.
Arch Ophthalmol 69: 503-507.

Kalvin, N. H.; Hamasaki, D. I. and Gass, J.D.M. (1966):

Experimental glaucoma in monkeys.
II studies of intraocular vascularity during glaucoma.
Arch Ophthalmol 76: 94-103.

La Grange, H. and Beauvieux (1925):

Anatomie pathologique et pathogenie de l'excavation glaucomateuse.
Arch Ophthalmol 42: 129-145.
Quoted from: Spaeth GL. (1971): Pathogenesis of visual

loss in patients with glaucoma. *Trans Am Acad Ophthalmol and Otolaryngol* 75: 296-316.

Lieberman, M. F.; Maumenee, A. E. and Green, W. R. (1976):

Histologic studies of the vasculature of the anterior optic nerve.

Am. J. Ophthalmol 82: 405-423.

Levitzky, M. and Henkind, P. (1969):

Angioarchitecture of the optic nerve.

2- Lamina cribrosa.

Am.J. Ophthalmol 68: 986-995.

Levy, N. S. (1974 a):

The effect of elevated intraocular pressure on slow axonal protein flow.

Invest. Ophthalmol. 12: 771.

Levy, N. S. (1974 b):

Functional implications of axoplasmic transport.

Invest. Ophthalmol. 13: 639.

Loebl, M.; Schwartz, B. (1977):

Fluorescein angiographic defects of the optic disc in

ocular hypertension.

Arch Ophthalmol 95 : 1980-1984.

McLean, J. (1957):

Management of the primary glaucomas.

Am. J. Ophthalmol 44 : 323-334.

Minckler, D. S. (1980):

The organization of nerve fiber bundles in the primate optic nerve head.

Arch Ophth. 98: 1630-1636.

Minckler, D. S. and Tso, M.O.M. (1976):

A light microscopic, autoradiographic study of axoplasmic transport in the normal rhesus optic nerve head.

Am.J. Ophthalmol 82: 1.

Minckler, D.S.; Mclean, I. and Tso, M.O.M.(1976):

Distribution of axonal and glial elements in the rhesus optic nerve head studied by electron microscopy.

Am.J. Ophthalmol. 82: 179.

Nanba, K. and Schwartz, B. (1988):

Nerve fiber layer and optic disc fluorescein defects in glaucoma and ocular hypertension.

Ophthalmology 95: 1227-1233.

Obstbaum, S. A. and Podos, S. M. (1974):

Axoplasmic transport.

Invest. Ophthalmol 12: 81.

Quigley, H. A.; Guy, J. and Anderson, D. R. (1979):

Blockade of rapid axoplasmic transport. Effect of intraocular pressure elevation in primate optic nerve.

Arch Ophthalmol 97: 525-531.

Quigley, H. A. and Addicks, E. M. (1981):

Regional differences in the structure of the lamina cribrosa and their relation to glaucomatous optic nerve damage.

Arch Ophthalmol 99: 137-143.

Radius, R. L. and Anderson, D. R. (1979):

The course of axons through the retina and optic nerve head.

Arch. Ophthalmol 97: 1154-1158.

Raitta, C. and Sarmela, T. (1970):

Fluorescein angiography of the optic disc and the peripapillary area in chronic glaucoma.

Acta Ophthalmol. 48: 303-308.

Reese, A. B. and McGavie, J. S. (1942):

Relation of field correlation to blood pressure in chronic primary glaucoma.

Arch Ophthalmol 27 : 845-850.

Rosen, E. S. and Boyd, T.A.S. (1970):

New methods of assessing choroidal ischaemia in open angle glaucoma and ocular hypertension.

Am. J. Ophthalmol 70: 912-921.

Schwartz B. (1976):

New techniques for the examination of the optic disc and their clinical application.

Trans Am Acad Ophthalmol and Otolaryngol. 81: 227-235.

Schwartz, B.; Riesser, J. C. and Fishbein, S. L. (1977):

Fluorescein angiographic defects of the optic disc in glaucoma.

Arch. Ophthalmol. 95: 1961-1974.

Sebag, J.; Thomas, J. V.; Ebstein, D. L. and Grant, W. M. (1986):

Optic disc cupping in arteritic anterior ischaemic optic neuropathy resembles glaucomatous cupping.

Ophthalmology 93: 357-361.

Sonty, S. and Schwartz, B. (1980):

Two-point fluorophotometry in the evaluation of glaucomatous optic disc.

Arch. Ophthalmol 98: 1422-1426.

Spaeth, G. L. (1971):

Pathogenesis of visual loss in patients with glaucoma.

Trans Am. Acad Ophthalmol and Otolaryngol 75: 296-316.

Swietliezko, I. and David, N.J.(1970):

Fluorescein angiography in experimental ocular hypertension.

Am.J.Ophthalmol. 70: 351.

Talusan, E. d. and Schwartz, B. (1977):

Specificity of fluorescein angiographic defects of the optic disc in glaucoma.

Arch Ophthalmol 95: 2166-2175.

Talusan, E. D.; Schwartz, B. and Wilcox, L. M. (1980):

Fluorescein angiography of the optic disc.

Arch. Ophthalmol 98: 1579-1587.

Wolff, E. (1976):

The anatomy of the eye and orbit.

7th Ed., Lewis, London.