

References

REFERENCES

Abdel Fattah, N.S. and El-Desoki, M.M. (1990):

Eye infection and contact lens during lens wear.

Bull. Ophthalmol. Soc. Egypt., 83 : 437–440.

Abdel Fattah, N.S.; El-Refaei, F.M.; El-Matarawy, M.A. and El-Sergany, T.M. (1991):

Biophysical studies of serum and tear immunoglobulins in bacterial and fungal corneal ulcers.

Bull. Ophthalmol. Soc. Egypt., 84 : 483–488.

Abdel Rehim, M.H. (1979):

Continuous wear soft lenses for aphakia.

Bull. Ophthalmol. Soc. Egypt., 72 : 137–140.

Abdel Rehim, M.H. and El-Samma, H.E. (1985):

Management of giant papillary conjunctivitis associated with soft contact lens wear.

Bull. Ophthalmol. Soc. Egypt., 78 : 393–399.

Abdel Rehim, M.H. and Sami, M. (1989):

Effect of different types of contact lenses on fungal flora of healthy eyes.

Bull. Ophthalmol. Soc. Egypt., 81 : 591–595.

Abdel Rehim, M.H.; Sami, M. and Ismail Z. (1990):

Giant papillary conjunctivitis in contact lens wearers. New trends in pathogenesis and management.

Bull. Ophthalmol. Soc. Egypt., 83 : 441–445.

Abdel Wehab, H.; Galal, E. and Khayyal, E. (1989):

Study of incidence of giant papillary conjunctivitis with different contact lenses.

Bull. Ophthalmol. Soc. Egypt., 82 : 405–407.

Abelson, M.B.; Udell, I.J. and Weston, J.H. (1981):

Normal human tear pH by direct measurement.

Arch. Ophthalmol., 99 : 301.

Allansmith, M.R.; Korb, D.R.; Greiner, J.V. et al. (1977):

Giant papillary conjunctivitis in contact lens wearers.

Am. J. Ophthalmol., 83 : 697–708.

Allansmith, M.R. (1990):

Treatment of giant papillary conjunctivitis.

S. Afr. Optom., 49 (4) : 194–197.

Quoted from: Contact Lens Update, 10 (3) : 35.

Association of Contact Lens Manufacturers Ltd. (1989):

Framework for classifying contact lens materials.

Branksome Park Road, Camberley, Surrey GU 15 2AQ.

Avisar, R.; Savir, H.; Sidi, Y.; Pinkhas, J. et al. (1977):

Tear calcium and magnesium levels of normal subjects and patients with hypocalcemia or hypercalcemia.

Invest. Ophthalmol. Visual Sci., 16 : 1150–1151.

Azen, E.A. (1976):

Genetic polymorphism of human anodal tear protein.

Biochem. Genet., 225–235.

Quoted from van Haeringen (1981): Clinical Biochemistry of Tears, *Surv. Ophthalmol.*, 26 (2) : 84–96.

Binder, P.S. and Woodward, C. (1980):

Extended wear hydrocurve and sauflon contact lenses.

Am. J. Ophthalmol., 90 : 309–311.

Binder, P.S.; Rasmussen, D.M. and Gordon, M. (1981):

Keratoconjunctivitis and soft contact lens solutions.

Arch. Ophthalmol., 99 : 87.

Bonanno, J.A. and Polse, K.A. (1987):

Measurement of in-vivo human corneal stromal pH.

Invest. Ophthalmol. Visual Sci., 28 : 522–530.

Bonavida, B. and Sapse, A.T. (1968):

Human tear lysozyme. II: Quantitative determination with standard Schirmer strips.

Am. J. Ophthalmol., 66 : 70–76.

Bonavida, B.; Sapse, A.T. and Sercarz, E.E. (1969):

A unique lacrimal protein absent from serum and other secretions, specific tear prealbumin.

Nature, 221 : 375–377.

Boot, N.; Kok, J. and Kijlstra, A. (1989):

The role of tears in preventing protein deposition on contact lenses.

Current Eye Research, 8 (2) : 185–188.

Botelho, S.Y. and Martinez, E.V. (1973):

Electrolytes in lacrimal gland fluids and in tears.

Am. J. Physiol., 225 : 606–609.

Broekuyse, R.M. (1974):

Tear lactoferrin; a bacteriostatic and complexing protein.

Invest. Ophthalmol. Visual Sci., 13 : 550–554.

Brunish, R. (1956):

The protein components of human tears.

Arch. Ophthalmol., 9 : 554–556.

Butts, B.L. and Rengstorff, R.H. (1990):

Antioxidant and vitamin A eye drops for giant papillary conjunctivitis.

Contact Lens. J., 18 (2) : 40–44.

Caldeira, J.; Luthold, W. and Sabbaga, E. (1982):

Tear calcium levels in normal subjects.

Ophthalmologica (Basel), 184 : 114–119.

Campell, F. (1951):

The influence of low atmosphere pressure on the development of retinal vessels in the rat.

Trans. Ophthalmol Soc. U.K., 71 : 287.

Carney, L.G. and Hill, R.M. (1976):

Human tear pH diurnal variations.

Arch. Ophthalmol., 94 : 821-824.

Carney, L.G. and Hill, R.M. (1979):

Human tear buffering capacity.

Arch. Ophthalmol., 97 : 951-952.

Cedarstaff, T.H. and Tomlinson, A. (1983):

A comparative study of evaporation rates and water content of soft contact lenses.

Am. J. Optom. Physiol. Opt., 60 : 167-174.

Chandler, J.W.; Leder, R.; Kaufman, H.E. and Caldwell, T.R. (1974):

Quantitative determination of complement components and immunoglobulins in tears and aqueous humour.

Invest. Ophthalmol. Visual Sci., 13 : 151-153.

Chang, F.W. (1977):

The possible adverse effects of over-the-counter medications on the contact lens wearer.

J. Am. Optom. Assoc., 48 : 319-323.

Chun, M.W. and Weissman, B.A. (1987):

Compliance in contact lens care.

Am. J. Optom. Physiol. Opt., 64 : 274.

Ciolletti, K.R. (1980):

Determination of thiomersol content in contact lens polymer.

Int. Contact Lens. Clin., 13 : 3.

Cohen, E.J.; Gonzalez, C.; Leavitt, K.G., et al. (1991):

Corneal ulcers associated with contact lenses including experience with disposable lenses.

CLAO J., 17 (3) : 173–176.

Coyle, P.K.; Sibony, P.A. and Johnson, C. (1989):

Electrophoresis combined with immunologic identification of human tear protein.

Invest. Ophthalmol. Visual Sci., 30 (8) : 1872–1878.

Crandall, D.C. and Leopold, I.M. (1979):

The influence of systemic drugs on tear constituents.

Ophthalmology, 86 : 115–125.

Dado, V.K.; Acharyee, S.; Mohan, M. et al. (1982):

Corneal diseases in soft lens wearers.

Acta. XXIV Int. Congress Ophthalmol., 2 : 1165–1168.

DeHass, E.B.H. (1960):

Lacrimal gland response to parasympathomimetics after parasympathetic denervation.

Arch. Ophthalmol., 64 : 34–43.

Dohlman, C.H.; Friend, J.; Kalevar, V. et al. (1976):

The glycoprotein content of tears from normal and dry eye patients.

Exp. Eye Res., 22 : 359–365.

Donshik, P.C. and Ballow, M. (1983):

Tear immunoglobulins in giant papillary conjunctivitis induced by contact lenses.

Am. J. Ophthalmol., 96 : 460–466.

Donzis, P.B. (1987):

Microbial contamination of contact lens care system.

Am. J. Ophthalmol., 104 : 325.

Dreifus, M.; Wichtele, O. and Lim, D. (1960):

Intracameral lenses of hydrocolloid acrylate

Cesk Oftalmol., 16 : 154–159.

Quoted from: Contact Lenses Procedures and Techniques by Lowther, G.E. Chapter 1, p. 7. Published by Butterworths, 1982.

El-Matarawy, M.A. and El-Sergany, T.M. (1992):

Immunoglobulin concentrations in human tears in keratomalacia and corneal graft reaction.

Bull. Ophthalmol. Soc. Egypt., 85 : 465–467.

Epstein, A. and Donnenfield, E. (1989):

Epithelial microcysts and oedema associated with Acuvue disposable contact lens.

Contact Lens Update, 8 (3) : 37.

Eriksen, S. (1975):

Cleaning hydrophilic contact lenses. An overview.

Ann. Ophthalmol., 7 : 1223–1232.

Quoted from: Farris, R.L. (1985).

Farris, R.L. (1985):

Tear analysis in contact lens wearers.

Tr. Am. Ophth. Soc., 83 : 502–544.

Ford, L.C.; Delange, R.J. and Petty, R.W. (1976):

Identification of a non-lysozymal bactericidal factor in human tears and aqueous humor.

Am. J. Ophthalmol., 81 : 30.

Fowler, S.A. and Allansmith, M.R. (1980):

Evolution of soft contact lens coatings.

Arch. Ophthalmol., 98 : 95–98.

Franklin, R.M.; Kenyon, K.R. and Tomasi, T.B. (1973):

Immunohistologic studies of human lacrimal gland.

J. Immunol., 110 : 984–992.

Fraunfelder, R.T (1976):

Drug induced ocular side effects.

Philadelphia, Lea & Febiger, p. 132.

Quoted from: Farris, R.L. (1985): Tear analysis in contact lens wearers. *Tr. Am. Ophth. Soc.*, 83 : 502–544.

Frederichs, F.T. (1846):

Thranensection.

In: Wagner R: *Handwörterbuch der Physiologie*.

Braunschweig F. Vieweg and Son, 1846, p. 617–630.

Quoted from: Farris, R.L. (1985): Tear analysis in contact lens wearers. *Tr. Am. Ophth. Soc.*, 83 : 506.

Frey, W.H.; DeSota-Johnson, D.; Hoffman, C. et al. (1987):

Effect of stimulus on the chemical composition of human tears.

Am. J. Ophthalmol., 92 : 559–567.

Fucrow and van Quelan (1791):

Chemical analysis of tears.

In: Wagner R: Handwörterbuch der Physiologie.

Braunschweig F. Vieweg and Son, 1846, p. 617–630.

Quoted from: Farris, R.L. (1985): Tear analysis in contact lens wearers. *Tr. Am. Ophth. Soc.*, 83 : 506.

Fullard, J. (1987):

Identification of proteins in small tear volumes with and without size exclusion HPLC fractionation.

Current Eye Research, 7 (2) : 163–179.

Gellatly, K.W.; Brennan, N.A. and Efron, N. (1988):

Visual decrement with deposit accumulation on HEMA contact lenses.

Am. J. Optom. Physiol. Opt., 65 (12) : 937–941.

Gitelman, H.F. (1967):

Calcium assay by cresolphthalein complexone.

Anal. Biochem., 18 : 521.

Gudmundsson, O.G.; Woodward, D.F.; Fowler, S.A. and Allansmith, M.R. (1985):

Identification of protein in contact lens surface deposit by immunofluorescence microscopy.

Arch. Ophthalmol., 103 : 193–197.

Hart, D.E.; Lane, B.C.; Josephon, J.E. et al. (1987):

Spoilage of hydrogel contact lenses by lipid deposits.

Ophthalmology, 94 : 1315–1321.

Hartstein, J. (1979):

Basics of Contact Lenses. 3rd edition. American Academy of Ophthalmology.

Heremans, J.F. (1968):

Immunoglobulin formation and function in different tissues.

Curr. Top. Microbiol. Immunol., 45 : 131.

Hill, R.M. (1975):

Tear cholesterol and your contact lens patient.

Aust. J. Optom., 58 : 300–303.

Hill, R.M. and Carney, L.G. (1977):

Tear pH and the soft contact lens patient.

Int. Contact Lens. Clin., 617 : 71–76.

Holden, B.A. and Mertz, G.W. (1984):

Critical oxygen levels to avoid corneal oedema for daily and extended wear contact lenses.

Invest. Ophthalmol. Visual Sci., 25 : 1161.

Holden, B.A.; Sweeney, D.F. and Sanderson, G. (1985):

The minimum precorneal oxygen tension to avoid corneal oedema.

Invest. Ophthalmol. Visual Sci., 26 : 1489.

Holden, B.A.; Ross, R. and Jenkins, J. (1987):

Hydrogel contact lenses impede carbon dioxide efflux from the human cornea.

Curr. Eye Res., 6 : 1283–1290.

Holden, B.A. (1990):

Factors affecting the corneal response to contact lenses.

CIBA Vision Monograph Series, (1) : 17.

Holden, B.A. and Sweeney, D.F. (1991):

The significance of the microcyst response: a review.

Optom. Vis. Sci., 68 (9) : 703–707.

Quoted from: Contact Lens Update, 10 (2) :26.

Holly, F.J. and Lemp, M.A. (1977):

Tear physiology and dry eye.

Surv. Ophthalmol., 22 : 69.

Holly, F.J. (1981):

Tear film physiology and contact lens wear.

Am. J. Optom. Physiol. Opt., 59 : 331–341.

Horowitz, G.; Lammers, J.W.; Sussman, S. and Botelho, S.Y. (1976):

Calcium in tears, lacrimal gland fluid and fluid from the other orbital glands.

Invest. Ophthalmol. Visual Sci., 15 (12) : 994–995.

Huth, S.W.; Hirano, P. and Leopold, I.H. (1980):

Calcium in tears and contact lens wear.

Arch. Ophthalmol., 98 : 122-125.

Huth, S.W.; Miller, M.A.; Michael, J. et al. (1981):

Calcium and proteins in tears: Diurnal variation.

Arch. Ophthalmol., 99 : 1628-1633.

Janssen, P.T. and van Bijsterveld, O.P. (1983):

Origin and biosynthesis of human tear fluid proteins.

Invest. Ophthalmol. Visual Sci., 24 : 623-630.

Jones, B. and Sack, R. (1990):

Immunoglobulin deposition on soft contact lenses.

Contact Lens. Assoc. Ophthalmol. J., 16 (1) : 43-48.

Jordan, A. and Baum, J. (1980):

Basic tear flow, does it exist?

Ophthalmology, 87 (9) : 920-930.

Kastle, P.R.; Refojo, M.F. and Dabezies, O.H. (1984):

Review of polymerization for the contact lens fitter.

In: Dabezies, O.H. (Ed.). *Contact Lenses: The CLAO Guide to Basic Science and Clinical Practice*. Vol. 1, Chapter 6.

New York: Grune and Stratton.

Kersley, H.; Kerr, C. and Pierse, D. (1977):

Hydrophilic lenses for continuous wear in aphakia:

Definitive fitting and problems.

Br. J. Ophthalmol., 61 : 38.

Klintworth, G.K.; Reed, J.W.; Hawkins, H.K. and Ingram, P. (1977):

Calcification of soft contact lenses in patient with dry eye
and elevated calcium concentrations in tears.

Invest. Ophthalmol. Visual Sci., 16 (2) : 158–161.

Klyce, S.D. (1981):

Stromal lactate accumulation can account for corneal
oedema following epithelial hypoxia in the rabbit.

J. Physiol., 321 : 49.

Korb, D.R.; Greiner, J.V.; Finnemor, V.M. and Allansmith, M.R.

(1983):

Biomicroscopy of papillae associated with wearing of soft
contact lenses.

Br. J. Ophthalmol., 7 : 733–736.

Lemp, M.A.; Holly, F.J.; Iwata, S. et al. (1970):

Factors in spreading and maintaining a continuous tear film
over the corneal surface.

Arch. Ophthalmol., 33 : 89–94.

Liotet, S. and Rouchy, J.P. (1970):

Etude des immunoglobulines des larmes humaines.

Arch. Ophthalmol. (Paris), 30 : 799–808.

Quoted from: van Haeringen, N.J. (1981): Clinical
biochemistry of tears. *Survey of Ophthalmol.*, 26 (2) : 84–
96.

Liotet, S.; Warnet, V.N. and Arrata, M. (1982):

Functional exploration of the lacrimal gland by tear electrophoresis.

Ophthalmologica (Basel), 184 : 87-91.

Liotet, S.; Guillaumin, D.; Cochet, P. et al. (1983):

The genesis of organic deposits on soft lenses.

Contact Intraocu. Lens Med. J., 99 : 1382-1386.

Lowry, O.H.; Rosenbough, N.J.; Farr, L.A. and Randall, R.J. (1951):

Protein measurement with the folin phenol reagent.

J. Biol. Chem., 193 : 265-275.

Lowther, G.E.; Miller, R.B. and Hill, R.M. (1970):

Tear concentrations of sodium and potassium during adaptation to contact lenses.

Am. J. Optom. Physiol. Opt., 47 : 261-265.

Lowther, G.E. (1982):

Contact lens terminology and types.

In: Lowther, G.E. (Ed): Contact Lenses Procedures and Techniques. Chapter 1, p. 31. Butterworth.

Lundh, R.L.; Liotet, S. and Pouliquen, Y. (1984):

Study of the human blood-tear barrier and the biochemical changes in tears of 30 contact lens wearers.

Ophthalmologica (Basel), 188 : 100-105.

Mackie, I.A. and Seal, D.Y. (1976):

Quantitative tear lysozyme assay in units of activity per microlitre.

Br. J. Ophthalmol., 60 : 70–74.

Mackie, I.A.; Seal, D.V. and Prescod, H.M. (1977):

Beta adrenergic receptor blocking drugs: Tear lysosyme and immunological screening of adverse reaction.

Br. J. Ophthalmol., 61 : 354–359.

Mancini, G.; Carbonera, A. and Heremans, J. (1965):

Immunochemical quantitation of single radial immunodiffusion.

Immunochem., 2 : 235.

Mandell, R.B. and Fatt, I. (1965):

Thinning of the human cornea on awakening.

Nature, 208 : 292.

Quoted from: Weissman, B. (Ed.): Contact Lens Primer.

Chapter 1, p. 8. Lea and Febiger, 1984.

Mannucci, L.L.; Pozzan, M.; Fregona, I. and Secchi, A. (1984):

The effect of extended wear contact lenses on tear immunoglobulins.

Contact Lens. Assos. Ophthalmol. J., 10 : 164–165.

McClellan, B.H.; Whitney, C.R.; Newman, L.P. et al. (1973):

Immunoglobulins in tears.

Am. J. Ophthalmol., 76 : 89–101.

McCulley, J.P. and Sciallis, G.F. (1977):

Meibomian keratoconjunctivitis.

Am. J. Ophthalmol., 84 : 788-793.

McEwen, W.K.; Kimura, S.J. and Feeney, M.L. (1958):

Filter paper electrophoresis of tears.

Am. J. Ophthalmol., 45 : 67-70.

McGill, J.; Liakos, G.; Gouldin, N. and Seal, D. (1984):

Normal tear protein profiles and age related changes.

Br. J. Ophthalmol., 68 : 316-320.

McMonnies, C.W. (1987):

Loss of tolerance in successful contact lens wearer.

Clin. Exp. Optom., 70 (4) : 121-127.

McMonnies, C.W. (1991):

Aids to improved patient compliance.

In: Contact Lens Yearbook. Chapter 2, p. 12-14. Ed.:

Christopher Kerr. Medical & Scientific Publishing Ltd.,

Kent, England.

McMurray, D.N.; Rey, H.; Casazza, L.J. and Watson, R.R. (1977):

Effect of moderate malnutrition on concentration of immunoglobulins and enzymes in tears and saliva of Columbian children.

Am. J. Clin. Nutr., 1944-1948.

Miller, R.B. (1970):

Tear concentrations of sodium and potassium during adaptation to contact lenses.

Am. J. Optom. Physiol. Opt., 47 : 773–779.

Minarik, L. and Rapp, J. (1989):

Protein deposits on individual hydrophilic contact lenses.

Contact Lens. Assoc. Ophthalmol. J., 15 (3) : 185–188.

Mishima, S. and Maurice, D.M. (1961):

Oily layer of tear film and evaporation from corneal surface.

Exp. Eye Res., 1 : 39–45.

Mishima, S. (1965):

Some physiological aspects of the precorneal tear film.

Arch. Ophthalmol., 73 : 233–241.

Mishima, S.; Gasset, A.; Klyce, S.D. et al. (1966):

Determination of tear volume and flow.

Invest. Ophthalmol. Visual Sci., 5 : 264–276.

Mishima, S. (1972):

Soft Contact Lenses.

St. Louis, CV Mosby Co., p. 19–36.

Molinari, J. (1989):

Chemotherapeutic management of giant papillary conjunctivitis.

Contact Lens. J., 17 (8) : 264.

Molinari, J. (1991):

Giant papillary conjunctivitis management: a review of management and a retrospective study.

Clin. Eye Vis. Care, 3 (2) : 68–72.

Mostafa, A.M.; Hegazy, M.A.; Elyan, S.A. and El-Atrebi, T.M. (1990):

Eye infection in contact lens wearers in Egypt.

Bull. Ophthalmol. Soc. Egypt., 83 : 111–117.

Mullen, W.; Shephard, W. and Labovitz, J. (1973):

Ophthalmic preservations and vehicles.

Surv. Ophthalmol., 17 : 469–471.

Neilson, N.V. and Jens, S. (1979):

Timolol transitory manifestations of dry eyes and long term treatment.

Arch. Ophthalmol., 57 : 418–424.

Ohnishi, S.T. and Barr, J.K. (1978):

A simplified method of quantitating proteins using the biuret and phenol reagents.

Anal. Biochem., 86 : 193.

O'Neal, M.R.; Polse, K.A. and Sarver, M.D. (1984):

Corneal responses to rigid and hydrogel lenses during eye closure.

Invest. Ophthalmol. Visual Sci., 25 : 837.

Parker, J. (1990):

The classification of contact lens materials.

In: Contact Lens Yearbook. Ed.: Christopher Kerr. Medical
& Scientific Publishing Ltd., Kent, England. P. 52-54.

Pietsch, R.L.; Pearlman, M.E. and Durham, S. (1973):

Human tear lysozyme variables.

Arch. Ophthalmol., 90 : 94-96.

Pipkin, F.B. (1984):

Medical Statistics Made Easy.

Churchill Livingstone: London, New York.

Pitts, R.E. and Krachmer, J.M. (1979):

Evaluation of soft contact lens disinfection in the home
environment.

Arch. Ophthalmol., 97 : 470.

Polse, K.A. and Mandell, R.B. (1970):

Critical oxygen tension at the corneal surface.

Arch. Ophthalmol., 84 : 505-508.

Rao, G.N. (1984):

Complications due to daily wear of soft contact lenses.

In: Dabezies, O.H. (Ed.): Contact Lenses: The CLAO Guide
to Basic Science and Clinical Practice. Vol. 2, Chapter 12.
New York: Grune and Stratton.

Refojo, M.F. and Holly, F.J. (1977):

Tear protein absorption on hydrogels: A possible cause of contact lens allergy.

Contact Intraoc. Lens Med. J., 3 : 23.

Rolando, M. and Refojo, M.F. (1983):

Tear evaporimeter for measuring water evaporation rate from the tear film in humans.

Exp. Eye Res., 36 : 25–33.

Ruben, M.; Tripathi, R.C. and Winder, A.F. (1975):

Calcium deposition as a cause of spoilage of hydrophilic soft contact lenses.

Br. J. Ophthalmol., 59 : 141–148.

Sack, A.; Jones, B. and Antignani, A. (1987):

Specificity and biological activity of the protein deposited on the hydrogel surface.

Invest. Ophthalmol. Visual Sci., 28 : 842–849.

Sapse, A.T.; Bonavida, B.; Stone, W. and Sercarz, E. (1969):

Proteins in human tears.

Arch. Ophthalmol., 81 : 815–819.

Schirmer, O. (1903):

Studies zur psychologie und pathologie der tränenabsonderung und tranenabfuhr.

Albrecht von Graefes Arch. Ophthalmol., 56 : 197–291.

Quoted from: van Haeringen (1981): Clinical biochemistry of tears. *Surv. Ophthalmol.*, 26 (2) : 84–96.

Schmidt, P.O.; Schoessler, T.P. and Hill, R.M. (1974):

Effect of hard contact lenses on chloride ion of tears.

Am. J. Physiol. Opt., 51 : 84–87.

Seal, D.V.; Mackie, I.A.; Coakes, R.L. and Faroogi, B. (1980):

Quantitative tear lysozyme assay: a new technique for transporting specimens.

Br. J. Ophthalmol., 64 : 700–704.

Sen, D.K.; Sarin, G.S.; Mathur, G.P. and Kunal Saha (1978):

Biological variations of immunoglobulin concentrations in normal human tears related to age and sex.

Arch. Ophthalmol., 56 : 439–444.

Sen, D.K. and Sarin, G.S. (1979):

Immunoglobulin concentrations in human tears in ocular diseases.

Br. J. Ophthalmol., 63 : 297–300.

Shebl, S.S.; Bayoumi, M.Y.; El-Desouki, E.S. and Zaki, R. (1992):

Practical insight on contact lens intolerance.

Bull. Ophthalmol. Soc. Egypt., 85 : 481–485.

Shultz, J.E. (1990):

Treating giant papillary conjunctivitis while wearing contact lenses.

Int. Contact Lens. Clin., 17 : 139–143.

Smolens, J.; Leopold, I.H. and Parker, J. (1949):

Studies of human tears.

Am. J. Ophthalmol., 32 : 402–416.

Smolin, G.; Kumato, M. and Nozik, R.A. (1979):

The microbial flora in extended wear soft contact lens wearers.

Am. J. Ophthalmol., 88 : 543–546.

Sommer, A. (1982):

Conjunctival appearance in corneal xerophthalmia.

Arch. Ophthalmol., 100 :.

Sørensen, T. and Jaagehøj Jensen, F. (1979):

Tear flow in normal human eyes.

Arch. Ophthalmol., 57 : 564–580.

Stuchell, R.N.; Farris, R.L. and Mandel, I.D. (1981):

Basal and reflex human tear analysis: II. Chemical analysis: Locatoferrin and lysozyme.

Ophthalmology, 88 : 858–862.

Stuchell, R.N.; Feldman, J.J.; Farris, R.L. and Mandel, I.D. (1984):

The effect of collection technique on tear composition.

Invest. Ophthalmol. Visual Sci., 2 : (3) : 374–377.

Takano, Y. and Higashi, M. (1979):

Correction of corneal astigmatism with soft contact lenses.

J. Jap. Contact Lens. Soc., 21 (3) : 88–90.

Tapasztó, I. (1973):

Pathophysiology of human tears.

Int. Ophthalmol. Clinic, 13 : 119–145.

Tapasztó, I.; Tapasztó, Z.S.; Tapasztó, B. (1989):

Biochemical changes in the human tears of hard and soft contact lens wearers.

Contact Lens Journal, 17 (1) : 5–7.

Temel, A.; Kazokoglua, H.; Taga, Y. and Orkan, A. (1991):

The effect of contact lens wear on the tear immunoglobulins.

Contact Lens. Assos. Ophthalmol. J., 17 (1) : 69–71.

Thaysen, J.H. and Thorn, N.A. (1954):

Excretion of urea, sodium and chloride in human tears.

Am. J. Physiol., 178 : 160–164.

Thompson, R. and Gallardo, E. (1941):

The antibacterial action of tears on staphylococci.

Am. J. Ophthalmol., 24 : 635–639.

Tiffany, J.M. (1978):

Individual variations in human meibomian glands.

Exp. Eye Res., 27 : 289–300.

Tripathi, R.C. and Ruben, M. (1973):

Degenerative changes in a soft hydrophilic contact lens.

Ophthalmic Res., 4 : 185.

Tripathi, R.C.; Tripathi, B.J. and Rubein, M. (1980):

The pathology of soft contact lens spoilage.

Ophthalmology, 87 (5) : 365–380.

Tripathi, R.C. and Tripathi, B.J. (1984):

Soft lens spoilage: Nature, aetiology and pathogenesis.

Ophthalmic Forum, 2 (2).

Tripathi, R.C.; Tripathi, B.J.; Silverman, R.A. and Rao, G.N. (1991):

Contact lens deposits and spoilage: Identification and management.

Int. Ophthalmol. Clinic., 31 (2) : 92–120.

Uotila, M.H.; Soble, R.E. and Savory, J. (1972):

Measurements of tear calcium levels.

Invest. Ophthalmol. Visual Sci., 11 : 258–259.

Vale, J. (1990):

The effects of drugs on the tear film and contact lenses.

Trans. Br. Contact Lens. Assoc. Sci. Meetings, p. 21–23.

van Haeringen, N.J. and Glasius, E. (1975):

Cholesterol in human tear fluid.

Exp. Eye Res., 20 : 271–274.

van Haeringen, N.J. and Glasius, E. (1977):

Collection method dependent concentrations of some metabolites in human tear fluid, with special reference to glucose in hyperglycemic conditions.

Albrecht von Graefes Arch. Klin. Exp. Ophthalmol., 202 : 1–7.

Quoted from: van Haeringen, N.J. (1981).

van Haeringen, N.J. (1981):

Clinical biochemistry of tears.

Survey of Ophthalmology, 26 (2) : 84–96.

Versura, P.; Maltarello, M.; Caramazza, R. and Laschi, R. (1988):

Immunocytochemical analysis of contact lens surface deposits in transmission electron microscopy.

Curr. Eye Res., 7 : 277–286.

Vinding, T.; Eriksen, T. and Nielsen, N. (1987):

The concentrations of lysozyme and secretory IgA in tears from healthy persons with and without contact lens use.

Acta. Ophthalmol., 65 (1) : 23–26.

Weissman, B.A. (1988):

Open-eye corneal swelling secondary to hydrogel contact lens wear.

Am. J. Optom. Physiol. Opt., 65 : 272.

Weissman, B.A. (1989):

Corneal infection and contact lens wear.

Contact Lens Update, 8 (3) : 46–48.

Wichtele, O. and Lim, D. (1960):

Hydrophilic gel for biological uses.

Nature (Lond.), 183 : 117.

Quoted from Lowther, G.E. (Ed.): Contact Lenses

Procedures and Techniques. Butterworth, 1982, Chapter 1, p. 7.

Wilson, G.; O'Leary, D. and Holden, B.A. (1989):

Cell content of tears following over-night wear of a contact lens.

Current Eye Res., 8 : 329–335.

Wolff, E. (1954):

Anatomy of eye and orbit. ed. 4. New York: Blakiston Co.,
p. 207–209.

Yamane, S.J. (1979):

Studies with a unit dose saline solution.
Contact Lens Forum, 4 : 91–95.

Young, W.H. and Hill, R.M. (1973):

Tear cholesterol levels and contact lens adaptation.
Am. J. Optom. Physiol. Opt., 50 : 12–16.

Zubareva, T.V. and Kiseleva, Z.M. (1977):

Catecholamine content of lacrimal fluid in healthy people
and glaucoma patients.
Ophthalmologica (Basel), 75 : 339–344.