

References

- Abd El Rahman, N.S. (1993):** Ecological studies on the distribution of zooplankton communities in the northern part of the Suez Gulf (Suez Bay). M. Sc. Thesis, Fac., Sci., Suez Canal Univ., 220 pp.
- Abd El-Azim, H. (1996):** Lead dynamics in Suez Bay. M. Sc. Thesis, Fac., Sci., Helwan Univ., 120 pp.
- Abdallah, M.A. (1979):** study of the currents and hydrographic structure of the water masses in front of Alexandria coast. M Sc. Thesis, Fac., Sci., Alex. Univ., 169 pp.
- Abott, W.S. (1925):** A method of computing the effectiveness of an insecticide. J. Econ. Entomol., 18: 265-267.
- Alaboster, J.S. and Lloyed, R. (1980):** Water quality criteria for freshwater Fish. London. Butterworths for FAO. pp: 80-85.
- APHA (1981):** Standard method for the examination of water and wastewater. 17th (ed.): American public Health association, Washington. pp: 1193.
- Arin, M.L. (1974):** Monitoring with the carbon analyzers. J. Environ. Sci. Tech., 7: 898-902
- Armstrong, D.A.; Chippendale, D.; Knight, A.W and Colt, J.G. (1972):** Interaction of ionized ammonia in a short-term survival and growth of prawn larvae. N. Rosenber. Gii. Biol. Bull., 154 (1): : 15- 31.

- Ashley, L.M. (1970): Pathology of fish fed aflatoxin and other antimetabolites. In: "A symposium on diseases of fishes and shellfishes". Snieizko, S.P. Ed., Special publ. S. Washington: Am. Fish. Soc., pp: 366- 379.
- Baker, J.T.P. (1969): Histological and electron microscopical observations on copper poisoning in the winter flounder (*Pseudopleuronectes americanus*): J. Fish. Res. Bd. Can., 26 (11): 2785 –2793.
- Baker-Cohen, K.F. (1969): Renal and other heteropic thyroid tissue in fishes. In: "Comparative endocrinology". A Gorbman (Ed), J. Wiley, New York, 283-301.
- Ball, T.R. (1967): The relative susceptibility of some species of freshwater fish to poisons. J. Am. Wat. Res., 1: 11-12.
- Belal, A.A.M. (1995): Ecological studies on Macrobenthic invertebrates, in the intertidal zone of the Suez region. M. Sc Thesis, Fac. Sci. Suez Canal Univ., 154pp.
- Beltagy, A. and Moussa, k.H.A. (1984): The distribution of organic matter in the. Sediments of the Gulf of Suez. Inst. Oceanogr. Fish. ARE., 10: 99 –109.
- Bower, C.E. and Bidwell, J.P. (1978): Ionization of ammonia in seawater, Effect of temperature, pH, and salinity. J. Fish. Res. Bd. Can., 35: 1012 – 1016.
- Bruno, D.W. and Ellis, A.E. (1986): Multiple hepatic cysts in farmed Atlantic salmon, *Salmo Salar* L. J. Fish Dis., 9: 79-82.
- Bucher, E. and Hofer, R.H. (1993): The effects of treated domestic sewage on three organs (gills, Kidney, liver): of brown trout *Salmo trutta*. J. Wat.Res., 27: 255-261.

- Burrow, R.E. (1964): Effects of accumulated excretory products on hatchery reared Salmonids. J. U.S. Fish. Wild. Res. Rep., 66: 1-12.
- Carballo, M. and Munoz, M.J. (1991): Effect of sublethal concentrations of four chemicals on susceptibility of Juvenile rainbow trout (*Oncorhynchus mykiss*): to saprolegniosis. Appl. Envir. Microbiol., 57: 1813-1816.
- Carline, P.; Benson, A.P. and Rother Bacher, H. (1987): Long term effects of treated domestic wastewater on Brown trout. J. Wat. Res., 21: 1409-1415.
- Colt, J.E. and Tchofanaglou, G. (1973): Evaluation of the short-term toxicity of nitrogenous compounds to channel catfish *Ictalurus punctatus*. J. Aquaculture, 3: 209-224.
- Cooper, L.H.N. (1958): Consumption of nutrient salts in the English Channel. In: A symposium. Int. Counc. Explor. Sea Bergen, 99 pp.
- DiMichele, L. and Taylor, H. (1978): Histopathological and physiological responses of *Fundulus heteroclitus* to naphthalene exposure. J. Fish. Res Bd., Canada, 35: 1060-1066.
- Doaoust, P.Y. and Ferguson, H.W. (1985): Nodular gill disease A unique form of proliferative gill disease in rainbow trout *Salmo gairdneri* Richardson. J. Fish. Dis., 8: 511-522-
- EAA (2000): The wastewater discharged in the Suez Bay. Annual report, Suez Governorate: 10pp (in Arabic).

- Galno, E.; Arillo, A. and Mensi, P. (1984): Involvement of the gill chloride cells of trout under acute nitrite intoxication. J. Comp. Biochem. Physiol., 77(A): 611-617.
- Ghittino, P. (1961): Eziologia palogenesie tentstivi di trasmissione della Oegenerazione lipoidea epatica Nella trutta (*Salmo gairdneri*). Vet. Ital., 12 :3 -16.
- Ghobashy, A.F.; Badawi, H. K.; El-Sayed, E.A and Belal, A.A. (2000): Growth and rate of settlement of *Hydroides elegans* (Haswell) in Suez Bay. Bull. Fac. Sci. Zagazig Univ., 22 (1): 204-220.
- Goldes, S.A.; Ferguson, H. W.; Moccia, R.D. and Daoust, P.Y. (1988): Histological effects of the inert suspended clay Kaolin on the gills of Juvenile rainbow trout, *Salmo gairdneri* Richardson. J. Fish. Dis., 11: 232-34.
- Goodman, L.R.; Hasen, D.J; Couch, J.A .and Forester, J. (1976): Effects of Heptachlor and Toxaphene on laboratory reared embryos and fry of the sheep head minnow. South Eastern. Association of Game and Fish commissions 13th Annual reference: 88-105
- Gardner, G.R. and Ycovich, P.P. (1970): Histological and haematological responses of an estuarine teleost to Cadmium. J. Fish. Res. Bd. Canada, 27: 2185-2196.
- Grassof, K. (1976): Methods of seawater analysis. Verlage Chemic. Weinberm (LTD): New York. 126pp.
- Grundy, K.D. (1971): Strategies for control of man-made Eutrophication. In: "Environmental Science and Technology". New York, 1184 - 1190.
- Hall, T.J.; Borton, D.L.; Walsh, A.H. and Walke, R.E. (1992): Histopathology of Rainbow trout (*Oncorhynchus mykiss*): after long-term exposure to biological treated bleached

- Kraft mill effluent in experimental stream channels. *J. Can. Aquat. Sci.*, 5: 49:1- 5.
- Hamed, M.F. (1992):** Seawater quality at the northern part of the Gulf of Suez and the nearby area of the Suez Canal. M. Sc. Thesis, Fac. Sci. Mansoura Univ., 71pp.
- Hartwell, S.I; Jin, H.J.; Chorry, D.S. and Carins, J. Jr. (1986):** Evaluation of statistical method for avoidance data of shoaling fish. *J. Hydro.*, 131 : 63-76.
- Heath, G.A. (1989):** Water pollution and fish physiology. CRC. Press, Inc. Weinberm (LTD) New York. 31– 46.
- Hendrson, P. (1979):** Utilization of Silver and Bighead carp for water quality improvements. In: "Aquaculture systems for wastewater treatment". Cstion B.K. and Reed S.C. (Project officers): Seminar proceedings and Engineering Assessment, 159 pp.
- Herbert, D.W.; Dowing, K.M. and Merkens, T.C. (1965):** Studies on the survival of fish in poisonous solutions. *J. Verb. Int. Ver. Sci. Limnol.*, 12: 789-794
- Hermann, H. and Klinke, R. (1965):** Fish pathology. T.F.H Pub. Inc. Ltd. Hong Kong. pp. 39-405.
- Higashi, R.M.; Gary, N. C.; Jonathan, M.S.; Jeffery, M.M. and Donald, B.C. (1992):** A polar high molecular mass constituent of bleached Kraft mill effluent is toxic to marine organisms. *J. Environ. Sci. Tech.*, 25 2413-2420.
- Hinton, D.E.; Baumann, P.C. and Gardner, G.R. (1992):** Histopathological biomarkes. In: "Biomarkers, Biochemical, Physiological and histological markers of anthropogenic". Hugget, R.J.,Kimerls, R.A.and Bergman,

- H.L. (Ed.). Stress. Lewis pub. Chelsea, MI. :155-209
- Hoover, K.L. (1984): Hyperplastic thyroid lesions in fish. Nat. Cancer. Inst., Monogr., 56: 275-287.
- Hussein, A.A.; Hagra, A.E.; Bishai, R.M. and El-Dyasty, M.E. (1999): effects of industrial pollution on some ecophysiological parameters of freshwater fish. J. Aqua. Biol. Fish., 4(2): 63-76.
- Hutuhinso, G.E. (1957): A treatise on limnology, Geography – physics and chemistry. John Wiley and Sons. Ltd., New York. 1 (XIV): pp 1015.
- Jaber, A.H. (1995): Industrial wastewater problems in Suez City. Rep. 29th, January. Int., Suez. 30 pp.
- Jayanagoudar, T.S. and Canapali, S.V. (1965): Some observations on the use of sewage stabilization lagoons in India. Acta Hydro. Limin. Parasitol., 26 :331-348.
- Johnson, L.L; Steher, C.M.; Olson Paul, O. and Myers, M.S. (1992): Chemical contaminants and hepatic lesions in winter flounder (*Pleuronectes americanus*): from the northeast coast of the United States. Environ. Sci. Tech., 27 (13): 2759-2771.
- Jordan, D.H.M. and Lloyed, R. (1964): The resistance of rainbow trout *Salmo gairadneri* and roach *Rutilus rutilus* to alkaline solutions. Inter. J. Air Water poll., 8: 405-408.
- Kandil, M.H. (1982): Hydrographical and Chemical Studies on the Red sea waters in front of Hurgada. M. Sc, Thesis, Fac., Sci., Alex. Univ., 120 pp.

- Karlsson-Norrgren, I.; Bjorklund, I. and Runn, P. (1986):** Acid water and aluminum exposure, gill lesions and aluminium accumulation in farmed brown trout *Salmo trutta*. J. Fish. Dis., 9:11-26.
- Kaur, K.; Dhawan, A. and Toor, H. S. (1993):** Toxic effect of industrial effluents on hatchability and viability of *cyprinus carpio* eggs. J. Bull. Environ. Toxic., 50 (5): 640-645.
- Khadre, S.E.M. and Shabana, M.B. (1991):** Histopathological and physiological disorders of acute crude oil toxicity in *Mugil cephalus*. Inter. Con. Mar. Fish. Mang. Develop. Alexandria, Egypt.
- Klinke, R.H. (1965):** Der phenolgehalt des wassers in seiner auswirkung auf den fischorganismus (Abstract in English) Arch. Fish. Wiss., 16:1- 16.
- Klontz, G.W; Stewart, B.C. and Kib, D.W. (1985):** On The etiology and pathophysiology of environmental gill disease in Juvenile salmonids. In: "Fish and shell fish pathology". T.F.H. Publ. Inc. Ltd. London. 20: 183-196.
- Korringa, P. (1976):** Safeguards in the exploitation of domestic effluents for aquaculture. In: "Harvesting polluted water, waste head and nutrient-laded effluents in the Aquaculture", Plenus Ltd. New York and London. pp 220-250.
- Kramer, J.R.; Herbs, S.E. and Allen, H.R. (1972):** phosphorus: Analysis of water, Biomass and sediment. In: "Nutrients in natural waters". Allen, N.E. and Kramer, J. R. (Eds.). Willey. Inter. Sci., Pub. pp: 51-100.

- Larmoyeux, J.D. and Piper, R.G. (1973): Effects of water reuse on rainbow trout in hatcheries. *Prog. Fish. Cult.*, 35: 2-8.
- Leatherland, J.F. (1994): Reflection on the thyroidology of fishes; from molecules to humankind." *Guelph. Ecthyol. Rev.*, 2: 1-67.
- Leino, R.L. and McCormick, J.H. (1984): Morphological and morphometrical changes in chloride cells in the gills of *Pimephales promelas* after chronic exposure to acid water. *Cell tissue Res.*, 36:121-129.
- Lloyed, R. (1961): The toxicity of ammonia to rainbow trout *Salmo garidneri* Richardson. *J. Wat. Waste. Treat.*, 8 : 278-279.
- Lowe, J.I. (1964): Chronic exposure of spot *leiostomus xanthurus* to sublethal concentrations of Toxaphene in Seawater. *J. Trans. Am. Fish. Soc.*, 93 :396-399.
- Mahmoud, N.M. (1994): Effect of El-Kabonon wastewater drain on the fingerlings of *Mugil seheli* in Suez Bay. M.Sc. Thesis. Fac. Sci. Suez Canal Univ., 100 pp.
- Mallat, J. (1985): Fish gill structural changes induced by toxicants and other irritants; a statistical review. *J. Can. Fish. Aquat. Sci.*, 42: 30 pp.
- Mancy, K.H. (1983): Assessing potential marine pollution in the Red sea. *Bull. Inst., Oceanogr. Fish.* 12 (9): 89-95.
- Mazher, M.A.; Ahsry, M.A. and Fathalla, M.M. (1987): Effects of environmental pollution by Crude oil on the Nile catfish (*Clarias lazera*). C.V.I. Mortality rates and haematological studies. *Proc. Zool. Soc., ARE.*, 14: 363 -372.

- Mercer, J.R.G.; Barker, D.E. and Khan, R.A. (1997):** Stress – related changes in (*Tautogolabrus adspersus*) living near a paper mill. J. Bull. Environ. Cont. toxic., 58 (3): 442-447.
- Meshal, A.H. (1967):** A physical study of the water pollution in Suez Bay (Hydrography of Suez Bay. M. Sc Thesis. Fac. Sci. Cairo Univ., 156 pp.
- Meshal, A.H. (1970):** Water pollution in Suez Bay. J. Inst. Oceanogr. Fish., 8 : 461-473.
- Middaugh, D.P. and Couch, J.A. (1977):** Responses of early life history stages of striped bass *Morone saxatilis* to chlorination. J. Chesapeake Sci., 18 (1): 141 – 153.
- Middaugh, D.P.; Burnett, L.E. and Couch, J.A. (1980):** Toxicological and physiological responses of the fish *Leiostomus xanthurus* exposed to chlorine-produced oxidants. J. Estu. Res. Fed., 2 : 132-141.
- Mitchell, S.J. and Cech, J.J. (1983):** Ammonia caused gill damage in channel Catfish (*Ictalurus punctatus*): confounding effects of residual chlorine. J. Can. Fish Aquat., Sci., 40: 242 pp.
- Molnar, K. (1982):** Nodular coccidiosis in the gut of the tench *Tinca tinca* L. J. Fish Dis., 5: 461-470.
- Morcos, S.A. (1960):** The tidal currents in the southern part of Suez Canal. Gcn. Ass. Helsinki, 51: 307-316.
- Morcos, S.A. (1970):** Physical and chemical oceanography of the Red sea. Oceanogr. Mar. Biol. Ann. Rev., 8: 73-202.

- River estuary Seattle Washington. J. Nat. Cancer. Inst., 60: 1445-1453.
- Redner, B.D. and Stickney, R.R. (1979):** Acclimation to ammonia by *Tilapia aurea*. J. Trans., Am. Fish. Soc., 108: 383-388.
- Roberts, R.J. (1989):** Neoplasia of fishes: In: "Fish pathology". Roberts, R. J. (Ed), London, Bailliere, Tindall. 232 pp
- Rogers, W.A. (1962):** A study of two streams receiving domestic sewage. Southeastern Assoc. of Game and fish Comm. 16th, annual conference, South Carolina. 449-463.
- Ryther, J.H. (1970):** Recycling human waste to enhance food production from the sea. Environ. Let., 10: 79-83.
- Saad, M.A.H. (1973):** Distribution of Phosphate in lake Maruit, a heavy polluted lake in Egypt. J. Wat. Air Soil Pollut., 2: 515 – 522.
- Saad, S.G.; Zaki, N.I. and Hamza, A.A. (1984):** Acute toxicity and pathological changes of the fish (*Tilapia nilotica*) exposed to different wastes of dyes processing. Environ. Manage. Develop. Countries. J., 11: 1-5.
- Said, T.O. (1992):** Study on Oil pollution in the northern part of the Gulf of Suez. M. Sc. Thesis, Fac. Sci. Mansoura Univ. Egypt, 120 pp.
- Sedrak, G.M. (1992):** Effect of Abu-Qir industrial effluents on the physiological characteristics of different stages of *Sparus auratus*. M. Sc Thesis, Fac. Sci. Alex. Univ., 168 pp.

- Shaker, B.S. and Khan, M.Z.A. (1983): Mathematical modeling of water quality of the Red Sea near Jeddah Coast. Sci. Res. Administration College of Engineering, King Abdelaziz Univ. Jeddah SA., 1 :1-9.
- Shriadah, M.M.A. (1982): Studies on the chemical composition of the Eastern Harbor in Alexandria. M. Sc Thesis, Fac. Sci. Alex., Univ., Egypt. 271 pp.
- Sis, K.F.; Ives, P.J.; Jons, D.M. and Lewis, D.H. (1979): The microscopic anatomy of oesophagus, stomach and intestine of the channel catfish (*Ictalurus punctatus*). J. Fish. Biol., 14:179.
- Skidmore, J.F. and Tovell, P.W.A. (1972): Toxic effects of Zinc sulphate on the gills of rainbow trout. J. Wat. Res., 6: 217-230.
- Skirrow, G. (1965): Chemical Oceanography. Acad. press. London. 1227 pp.
- Smart, G.R (1975): The Acute toxic mechanisms of Ammonia to the rainbow trout (*Salmo gairdneri*). J. Fish Biol., 8: 471-475.
- Smart, G.R. (1976): The effect of ammonia exposure on gill structure of the rainbow trout *Salmo gairdneri*. J. Fish. Biol., 8: 471-485.
- Smart, G.R. (1978): Investigation of toxic mechanisms of ammonia to fish-gas exchange in rainbow trout *Salmo gairdneri* exposed to acutely lethal concentrations. J. Fish. Biol., 12: 93-104.
- Smith, C.E. and Piper, R.G. (1975): Lesions associated with chronic exposure to ammonia. In: "The Pathology of fishes". Ribelin, W.E. and Migaki, G. (Ed.). Univ.

- Wisconsin Press, Madison, Wisconsin, USA. pp: 497 – 514.
- Spargue, J.B. (1971):** Measurement of pollutant toxicity to fish and sublethal effects and safe concentrations. *J. Wat. Res.*, 5 :245-266.
- Spies, K.B.; Slegeman, J.J.; Rice, D.W.; Woodin, B. and Thomas, P. (1990):** Sublethal responses of *Plathichthys stellatus* to organic contamination in San Francisco Bay with emphasis on reproduction. Roca Ratan, Lewis publishers, CRC. Press. 96 pp.
- Steher, C.M.; Johnson, L.L. and Myers, M.S. (1999):** Hydropic vacuolation in three species of fish from the US west coast; lesion description and relationship between occurrence and contaminant exposure. *J. Aqu. Dis.* 32: 119-135.
- Strin, J. (1972):** Possibilities for constructive use of domestic sewage (with an example of the lake of Tunis). In: "Marine pollution and sea life". Ruivo, M. (Ed.), Fishing News (book) Ltd., West. Byfleet. pp: 517-519.
- Tabat, K. (1962):** Toxicity of ammonia to aquatic animals with reference to the effect of pH and carbon dioxide. *Bull. Takai Prog. Fish. Res. Lab.*, 34-67.
- Takashi, H. (1982):** Fish histology. Kodansha, (Ltd). Tokyo, Japan. 63-64.
- Tarzwell, C.M. (1957):** The use of bioassays in the safe disposal of electroplating wastes. *Amer. Electroplaters, Sec. 44th Ann. Tech. Proc.*, 60 pp.

- Teh, J.; Adams, S.S.M. and Hinton, D.E. (1997): Histopathological biomarkers in feral freshwater fish population exposed to different types of contaminant stress. *J. Aquat. Toxic.*, 37 (1): 87-121.
- Thurston, R.V.; Russo, R.L. and Smith, C.E. (1981): Acute Toxicity of ammonia and nitrite to rainbow trout fry. *J. Trans. Am. Fish. Soc.*, 107:361-368.
- Wales, J.H. (1970): Hepatoma in winter flounder. *Special. Publical. Ann. Fish. Soc.*, 5: 351-665.
- Yassin, M. H. (1998): Biological and ecological studies on the pearl oyster, *pinctada radiata* (Mollusca, Lamellibranchia): from the red Sea, with special reference to its tolerance to water pollution. Ph.D. Thesis, Fac. Sci. Ain-Shams Univ., 158 pp.