

REFERENCES

REFERENCES

Abeyounis, C.J. (1981):

Malignancies. In: principles of Immunological Diagnosis in Medicine. Milgrom, F.; Abeyounis, P. and Kano, K. (eds). library of congress cataloging in publishing data, philadelphia, P.450.

Adolf, L.P. (1987):

Survillance of cancer patients with tumor markers.
J. tumor marker oncol., 2:1.

Ameral, A.L.; Nascimento, A. and Goellner, J.R. (1984):

Proliferating pilar (Trichelemmal cyts). Report of two cases. One with carcinomatous transformation and one with distant metastases.
Arch. Pathol. Lab. Med., 18 : 808.

Amon, R.B. and Goodkin, P. (1976):

Topical 5-fluorouracil and the basal cell nevus syndrome.
N. Engl. J. Med., 295 : 677.

Arnon, R.; Bustin, M. and Calef, E. (1976):

Immunologic cross-reactivity of antibodies to a synthetic undecapeptide analogous to the aminoterminal segment of the CEA, with the intact protein and with human sera.
Proc. Nati. Acad. Sci. U.S.A., 73: 2123.

Arnoud, J.P.; Koehl, C. and Adloff, M. (1980):

Carcinoembryonic antigen in Diagnosis and prognosis of colorectal carcinoma.

Dis. Colon Rectum, 141.

Banjo, C.; Shuster, J. and Gold, P. (1974):

Intermolecular Heterogeneity of the carcinoembryonic antigen.

Cancer Res., 34:2114.

Barsky, S.H.; Grossman, D.A. and Bhuta, S. (1987):

Desmoplastic Basal cell carcinomas possess unique basement membrane-degrading properties.

J. Invest. Dermatol., 88: 324.

Beastall, G.H.; Cool, B.; Rustin, G.J. and Jennings, J. (1991):

A review of the role of established tumor markers.

Ann. Clin. Biochem., 28:5.

Berger, M.S.; Locher, G.W. and Saurer, S. (1988):

Correlation of c-erb B-2 gene amplification and protein expression in human breast carcinoma with nodal status and nuclear grading.

Cancer Res., 48: 1238.

Birkmayer, G.D. (1988):

Tumor markers in diagnosis and therapy of cancer. Recent developments. Proceeding the 1st International conference of the Egyptian society of tumor marker oncol., Cairo, Egypt, P.1. (Meeting abstract).

Birkmayer, G.D. and Klavins, J.V. (1990):

Classification of tumor markers. The 2nd International conference of the Egyptian society of tumor marker Oncol., Cairo, Egypt, P. 46 (Meeting Abstract).

Bleehen, S.S. (1975):

Pigmented basal cell epithelioma. Light and electron microscopic studies on tumors and cell cultures.

Br. J. Dermatol., 93: 361.

Blewitt, R.W. (1980):

Why does basal cell carcinoma metastasize so rarely?

Int. J. Dermatol., 19: 144.

Bohn, H. and Dati, F. (1983):

Placental-related proteins. In: Protein in body fluids, Amino acids and tumor markers Diagnostic and clinical aspects., Ritzman S.E. and Killings Worth L.M. (Eds).

Alan R. Liss, New York, P. 331.

Borel, D.M. (1973):

Cutaneous basal squamous carcinoma. Review of the literature and report of 35 cases.

Arch. Pathol., 95: 293.

Broders, A.C. (1921):

Squamous-cell epithelioma of the skin.

Ann. Surg., 73: 141.

Quoted from lever, W.F. and Lever, G.S. (1990).

Brownstein, M.H. and Shapiro, L. (1976):

Verrucous carcinoma of the skin: Epithelioma cuniculatum plantare,

Cancer., 38: 1710.

Burton, P.; Von Kleist, S. and sabine, M.C. (1973):

Immunological localization of CEA and non specific cross reacting antigen in gasterointestinal normal and tumor tissues.

Cancer Res., 33:3299.

Caro, W.A. and Bronstein, B.R. (1985):

Tumors of the skin. In: Dermatology. Moschella, S.L. and Hurley, H.J. (eds). Library of congress cataloging in publicating data, philadelphia, P. 1533.

Caro, M.R. and Howell, J.B. (1952):

Marphea-like epithelioma.

Arch. Dermatol. Syph., 63: 471.

Quoted from Miller, S.G. (1991).

Clarke, C.; Hine, K.R.; Dykes, P.W.; Whitehead, T.P. and White field, A.G.W. (1980):

Carcinoembryonic antigen and smoking.

J. Rey. Coll. Phys. Lond., 14:227.

Constanza, M.E.; Pinn, V.; Schwartz, R.S. and Nathanson, L. (1973):

Carcinoembryonic antigen-antibody complexes in a patient with colonic carcinoma and nephrotic syndrome.

N. Engl. J. Med., 289:520.

Curigis, H.A.; Lynch, H.T. and Harris, R.E. (1978):

Carcinoembryonic antigen in the cancer family syndrome.
Cancer, 42:1574.

Dabska, M. (1969):

Malignant endovascular papillary angioendothelioma in childhood. Clinicopathologic study of 6 cases.
Cancer, 24:503.

Denk, H.; Tappeiner, G. and Eckerstorfer, R. (1972):

Carcinoembryonic antigen in gastrointestinal and extragastrointestinal tumors and its relationship to tumor-cell differentiation.
Int. J. Cancer, 10:262.

Deplech, A.; Deplech, B. and Girard, N. (1982):

Hyaluronectin in normal human skin and in Basal cell carcinoma.
Br. J. Dermatol. 106 : 561.

Disaia, P.J.; Marrow, C.P.; Haver back, B.J. and Dyce, B.J. (1977):

Carcinoembryonic antigen in cancer of the female reproductive system.
Cancer, 39:2365.

Dougherty, T.J. (1981):

Photoradiation therapy for cutaneous and subcutaneous malignancies.
J. Invest. Dermatol., 177: 122.

Dupree, W.B.; Langloss, J.M. and Weiss, S.W. (1985):

Pigmented dermatofibrosarcoma pro-tuberans (Bednar tumor): a pathologic, Ultrastructural, and immuno histochemical study.

Am. J. Surg. Pathol., 9:630.

Dvoretzky, I.; Fisher, B. and Haker, O. (1978):

Mutilating basal cell epithelioma.

Arch. Dermatol., 114: 239.

El-Ahmady, O.; Mahmoud, M.E.; El-Maraghy, A.; El-Hakim, M.S. and Hassan, G.Y. (1988):

Serum effusion alpha-2-globulin ratio as reliable marker in malignancy.

J. Tumor Marker Oncol., 3:441.

Fanger, H. and Parker, B.E. (1957):

Histochemical studies of some keratotic and proliferating skin lesions.

Arch. Pathol., 64: 143.

Farmer, E. and Helwig, E. (1985):

Metastatic Basal cell Carcinoma : A clinicopathologic study of seventeen Cases

Cancer, 46 : 798.

Flotte, T.J.; Hatcher, V.A. and Friedman-Kien, A.E. (1984):

Factor VIII-related antigen in Kaposi's sarcoma in young homosexual men.

Arch. Dermatol., 120:180.

Foot, N.C. (1947):

Adnexal carcinoma of the skin.

Am. J. Pathol., 23: 1.

Freeman, R. (1974):

The pathogenesis of pseudoepitheliomatous hyperplasia.

J. Cutan. Pathol., 1; 231.

Fuck, A.; Banjo, C.; Shuster, J.; Freedman, S.O. and Gold, P.
(1975):

Carcinomebryonic antigen: Molecular biology and clinical significance.

Biochem. Biophys. Acta, 417:123.

Fulling, U.F.; Strayer, D. and Santa, G. (1981):

Adenexal metaplasia in carcinoma in situ of skin.

J. Cutan Pathol., 8:79.

Gellin, G.; Kopf, A. and Garfinkel, L. (1965):

Basal cell carcinoma.

Arch. dermatol., 91: 38.

Giles, G.; Marks, R. and Faley, P. (1988):

Incidence of nonmelanocytic skin cancer treated in Australia.

Br. Med. J. Clin. Res., 296:13.

Gold, P. (1967):

Circulating antibodies against CEA of the human digestive system.

Cancer, 20:1663.

Gold, P. and Freedman, S.O. (1965):

Demonstration of tumor-specific antigens in human colonic carcinomata by immunological tolerance and absorption techniques.

J. Exp. Med., 121:439.

Gold, P.; Krupey, J. and Ansari, H. (1970):

Position of the CEA of the human digestive system in ultrastructure of tumor cell surface.

J. Nat. Cancer. Inst., 45:219.

Gomaa, A.; Hussein, A.; Hosny, M.; El-Naggar, T.; Khater, A.; Khalifa, A. and Helal, T.H. (1990):

Detection of CEA in tissue of Intrathoracic Malignancies.

Egypt. J. tumor marker oncol., 1 :21.

Gown, A.M.; Vogel, M.A.; Hoad, D. and McNutt, M.A. (1986):

Monoclonal antibodies specific for melanocytic tumors distinguish subpopulations of Melanocytes.

Am. J. Pathol., 123:195.

Gray, M.H.; Smoller, B.R. and McNutt, N.S. (1990):

Neurofibromas and neurotized melanocytic Nevi are immunohistochemically distinct Neoplasms.

Am. J. Dermatopathol., 12:234.

Greenway, H.T.; Carnell, R.C. and Tanner, D.J. (1986):

Treatment of basal cell carcinoma with intralesional interferon.

J. Am. Acad. Dermatol., 15: 437.

Grier, W. (1976):

Squamous cell carcinoma of the body and extremities. In:
Cancer of the skin, biology-diagnosis-management.
Andrade et al. (eds). Saunders, Philadelphia, P. 918.
Quoted from Harly, A. and Haynes, B. (1984).

Grimwood, R.; Ferris, C. and Mercill, D. (1986):

Proliferating cells of human Basal cell carcinoma are
located on the periphery of tumor nodules.
J. Invest. Dermatol., 86 :191.

Grinspan, D. and Abulafia, J. (1979):

Oral florid papillomatosis (verrucous carcinoma)
Int. J. Dermatol., 18: 608.

Guitart, J.; Bergfeld, W.F. and Tuthill, R.J. (1990):

Squamous cell carcinoma of the nail Bed.
Br. J. Dermatol., 123:215.

Gupta, A.K.; Cardella, C.J. and Haberman, H.F. (1986) :

Cutaneous malignant neoplasms in patients with renal
transplants.
Arch. Dermatol., 122: 1288.

Halprin, K.M.; Ohkawara, A. and Fuckui, K. (1968):

Enzymatic characteristics of human basal cell carcinoma.
Arch. Dermatol., 96:80.

Hansen, H.J.; Snijders, J.J. and Miller, E. (1974):

Carcinoembryonic antigen assays: A laboratory adjunct in the diagnosis and management of cancer.

J. Hum. Pathol., 5: 139.

Harach, H.R., Skinner, M. and Gibs, A.R. (1983):

Biological markers in human lung carcinoma: an immunopathological study.

Thorax, 38: 937.

Harley, A. and Haynes, B. (1984):

Pathophysiology of epidermal Malignant disorders. In: pathophysiology of Dermatological Diseases. Soter, N.A. and Baden, H.P. (eds). Library of Congress Cataloging in Publishing data. Philadelphia, P. 405.

Hartree, E.F. (1972):

Determination of a modification of the Lowery method that gives a linear photometric response.

Analyt. Biochem., 48: 422.

Headington, J.T. (1978):

Verrucous carcinoma.

Cutis, 21: 207.

Hellstrom, I.; Hellstrom, K.E. and Shepard, T.H. (1970):

Cell-mediated immunity against antigens common to human colonic carcinoma and fetal gut epithelium.

Int. J. Cancer, 6:346.

Herberman, R.B. (1976):

Immunologic approaches to the diagnosis of cancer.
Cancer, 37: 579.

Hori, M.; EGAM, A. and Maejima, K. (1978):

Electron microscopic study of sebaceous epithelioma.
J. Dermatol., 5:139.

Hoxtell, E.; Mandel, J. and Murray, S. (1977):

Incidence of skin carcinoma after renal transplantation
Arch. Dermatol., 113: 436.

Hyder, D.M. and Schnitzer, B. (1986):

Utility of Leu-M1 monoclonal antibody in the differential diagnosis of Hodgkin's disease.
Arch. Pathol. Lab. Med., 110:416.

Heyderman, E.; Graham, R.M.; Chapman, D.V.; Richardson, T.C. and Mckee, P.H. (1984):

Epithelial markers in primary skin cancer : an immunoperoxidase study of the distribution of epithelial membrane antigen (EMA) and CEA in 65 primary skin Carcinomas.
Histopathol., 8:423.

Hyman, A.B. and Michaelides, P. (1963):

Basal cell epithelioma of the palm.
Arch. Dermatol., 92: 571.

Johnson, W. and Helwig, E. (1966):

Adenoid squamous cell carcinoma (adenoacanthoma).
Cancer, 19: 1639.

Kerr, J.F.R. Bishop, C.J. and Searle, J. (1984):

Tumors. In: recent advances in histopathology.
Anthony P. and Macsueen RNM. (eds). Churchill, New York,
P.1.

**Khalifa, A.; Halim, H.; El-Ahmady, O.; Gomma, A.; Hussein,
H.A.; Safwat, T. and El-Naggar, T. (1988):**

The value of tumor markers in bronchial carcinoma.
J. Tumor Marker Oncol., 3:417.

Khoo, S.K. and Mackay, E.V. (1973):

Carcinoembryonic antigen in cancer of the female
reproductive system. Sequential levels and effects of
treatment.
Aust. N.Z.J. Obstet. Gynecol., 13:1.

Klee, G.G. and Go, V.L.W. (1987):

Carinoembryonic antigen and its role in clinical
practice. In: Tumor markers and tumor-Associated
antigens. Ghosh B.C. and Ghosh L. (eds). MC Grow-Hillbook
Company, New York, London, P. 43.

Knowles, D.M.II. (1985):

Lymphoid cell markers :Their distribution and usefulness
in the immunophenotypic of lymphoid neoplasms
Am. J. Surg. Pathol., 9:85.

Kumakiri, M. and Hashimoto, K. (1978):

Ultrastructural resemblance of basal cell epithelioma to primary epithelial germ.

J. Cutan. Pathol., 5: 53.

Lane, A.T.; Goldsmith, L.A. and McCoon, P.E. (1985):

Decreased anchoring-fibril antigens (AF1, and AF2) in basal cell carcinoma.

Arch. Dermatol. Res., 277: 499.

Lang, P.G.; McKelvey, A.C. and Nicholson, J.H. (1987):

Three dimensional reconstruction of the superficial multicentric basal cell carcinoma using serial sections and a computer.

Am. J. Dermatopathol., 9: 198.

Lattes, R. and Kessler, R.W. (1951):

Metastasizing basal-cell epithelioma of the skin. Report of two cases.

Cancer, 4: 866.

Lever, W.F. and Lever, G.S. (1990a):

Tumors and cysts of epidermis. In: Histopathology of the skin, Lever, W.F. and Lever, G.S. (eds). Vol. II J.B. Lippincott Company, Philadelphia, P. 523.

Lever W.F. and Lever, G.S (1990b):

Tumors of the epidermal appendages. In: Histopathology of the skin. Lever, W.F. and Lever, G.S.(eds).Vol.II J. B. Lippincott Company, Philadelphia, P. 578.

Lowenstein, M.S. and Zamcheck, N. (1978):

Carcinoembryonic antigen (CEA) levels in benign gastrointestinal disease states.
Cancer. 42: 1412.

Lund, H.Z. (1965):

How often does squamous cell carcinoma of the skin metastasize?
Arch. Dermatol., 92: 635.

Luri, B.B.; Lowenstein, M.S. and Zamcheek, N. (1975):

Elevated circulating CEA levels in benign extrahepatic biliary tract obstruction and inflammation.
J. Pathol., 88: 140.

Luzi, P.; Miracco, C. and Del vecchio, M. (1987);

Stereological study of desmosomes in basal cell carcinoma and seborrheic keratosis.
J. Submicrosc. Cytol., 19: 337.

Macadam, R.F. (1978):

An electron microscopic study of basal cell carcinoma.
J. Pathol., 126: 149.

Mackie, R.M. (1986):

Tumors of the skin. In: textbook of dermatology. Rook, A.;Wilkinson, D.S.;Ebling, F.J.G.;champion, R.H. and Burton, J.L. (eds). Vol. III. Blackwell scientific publications oxford London Edinburgh, P. 2375.

Madsen, A. (1965):

Studies on basal-cell epithelioma of the skin.

Acta. Pathol. Microbiol. 177: 9.

Quoted by Lever W.F. and Lever, G.S. (1990).

**Manivel, J.C.; Wick, M.R.; Swanson, P.E.; Patterson, K. and
Dehner, L.P. (1986):**

Endovascular papillary angioendothelioma of childhood: a
vascular lesion possibly characterized by high
endothelial cell differentiation.

Hum. Pathol., 17:1240.

Mark, R.; Wick, R.E.; Swanson, and Carlos, J. (1988):

Immunohistochemical findings in tumors of the skin. In:
Advances in immunohistochemistry. Ronald, A. (ed).
Dellellis, Raven Press, New York, P.395.

Martin, K.W. and Halpern, S.E. (1984):

Carcinoembryonic antigen production, Secretion and
kinetics in BALB/C mice and a nude mouse-human tumor
model.

Cancer Res., 44:5475.

Mazoujian, G.; Pikus, G.S. and Haagensen, D.E. (1984):

Extramammary paget's disease evidence for an apocrine
origin. An immunoperoxidase study of Gross cystic
disease fluid protein-15, CEA and Keratin proteins.

Am. J. surg, Pathol., 8:43.

McArdle, J.P.; Roff, B.T. and Muller, H.K. (1984):

The basal lamina in basal cell carcinoma, bowen's disease, squamous cell carcinoma and keratoacanthoma an immunoperoxidase study using an antibody to type IV collagen.

Pathol., 16: 67.

Mckenize, D. and Henderson, A.R. (1983):

Electrophoresis of lactate dehydrogenase isoenzymes.

Clin. Chem., 36:3476.

Miettinen, M.; Lehto, V.P.; Badley, R.A. and Virtanen, I. (1982):

Expression of intermediate filaments in soft tissue sarcomas.

Int. J. Cancer, 30:541.

Miettinen, M.; Lehto, V.P. and Virtanen, I. (1983):

Presence of fibroblastic-type intermediate filaments (vimentin) and absence of neuro filaments in pigmented nevi and malignant Melanomas

J. Cutan. Pathol., 10:188.

Milgrom, F.; ABE Younis, C.J. and KANO, K. (1981):

Fetal antigens as tumor markers. In : Principles of immunological Diagnosis in Medicine. Milgrom, F.; ABE Younis, C.D. and Kano, K. (eds). Library of congress cataloging in publishing data, Philadelphia, P. 452.

Miller, S.J. (1991):

Biology of basal cell carcinoma.

J. Am. Acad. Dermatol., 24:1.

Mohs, F. and Sahl, W. (1979):

Chemosurgery for verrucous carcinoma.

J. Dermatol. Surg. Oncol., 5:302.

Moll, R.; Franke, W. and Schiller, D. (1982):

The catology of human cytokeratins : Patterns of expression in normal epithelial, tumors and cultured cells.
Cell, 31 : 11.

Moller, R.; Reymann, F. and How-Jensen, K. (1979):

Metastases in dermatological patients with squamous cell carcinoma.

Arch. Dermatol., 115: 73.

Molnar, I.G.; Vandevoorde, P. and Gitnick, G.L. (1976):

Carcinoembryonic antigen levels in fluids bathing gasterointestinal tumor.

Gastroenterol., 70:513.

Montgomery, H. (1928):

Basal squamous cell epithelioma.

Arch. Dermatol. Syph., 18: 50.

Quoted from Lever, W.F. and Lever, G.S.(1990).

Muller, S.; Wilhelmj, C.J. and Harrison, E.J. (1974):

Adenoid squamous cell carcinoma,
Arch. Dermatol., 115:703.

Nakai, M. and Mori, M. (1986):

Immunohistochemical distribution of Monoclonal
antibodies against keratin in papillomas and carcinomas
from oral and nasopharyngeal regions.
Oral Surg., 62:292.

Nappi, O. and Wick, M.R. (1986):

Disseminated lobular capillary hemangioma (pyogenic
granuloma): a clinicopathologic study of two cases.
Am. J. Dermatopathol., 8:379.

Nilsson, O. (1989):

Mucin antigens as markers for gastrointestinal cancer.
Tumor Update, 4:3.

Ordonez, N.G. and Batsakis, J.G. (1984):

Comparison of Ulex europaeus I lectin and factor VIII-
related antigen in vascular lesions.
Arch. Pathol. Lab. Med., 108:129.

Parker, J.W. (1981):

A new look at malignant lymphoma.
Diagn. Med., 4:77.

Paver, K.; Poyzer, K. and Burry, N. (1973):

The incidence of basal cell carcinoma and their
metastases in Australia and New Zealand.
Aust. J. Dermatol., 14:53.

Peltonen, J.; Jaakkola, S. and Lask, G. (1988):

Fibronectin gene expression by epithelial tumor cells in Basal carcinoma: An immunocytochemical and in situ hybridization study.

J. Invest. Dermatol., 91 : 289.

Penneys, N.S.; Nadji, M. and Morales, A.R. (1982):

Carcinoembryonic antigen in benign sweat gland tumors.

Arch. Dermatol., 118: 225.

Pinkus, H. (1965):

Epithelial and fibroepithelial tumors.

Arch. Dermatol., 91:24.

Pinkus, G.S. and Kurtin, P.J. (1985):

Epithelial membrane antigen-a diagnostic discriminant in surgical pathology: immunohistochemical profile in epithelial mesenchymal and hematopoietic neoplasms using paraffin sections and monoclonal antibody.

Hum. Pathol., 16:929.

Pohl, A.L. (1983):

Present value of tumor markers in the clinic cancer.

Detect. Prev., 6:7.

Pohl, A.L. (1987):

Surveillance of cancer patients with tumor markers.

J. Tumor marker. oncol., 42:1.

Popkin, G.L. and Defea, C.P. (1976):

Basal cell epithelioma. In: Cancer of the skin, biology-diagnosis-management. Andrade et al. (eds). Saunders, Philadelphia, P. 821.

Quoted from Harley, A. and Haynes, B. (1984).

Posalaky, Z.; McGinley, D. and Cutler, B. (1979):

Intercellular junctional specializations in human basal carcinoma. A freeze, fracture study.

Virch. Arch., 384:53.

Reidbord, H.; Wechsler, H. and Fisher, E. (1971):

Ultrastructural study of basal cell carcinoma and its variants with comments on histogenesis.

Arch. Dermatol., 104:132.

Reynoso, G. (1981):

Carcinoembryonic antigen basic concepts, clinical applications.

Diagn. Med., 4:41.

Robinson, J.K. (1987):

Use of a combination of chemotherapy and radiation therapy in the management of advanced basal cell carcinoma of the head and neck.

J. Am. Acad. Dermatol., 17: 770.

Ronnen, M.; Suster, S. and Orrnstein, A. (1988):

Immunofluorescent localization of cytokeratin intermediate filaments as a means of defining the presence of recurrent Basal cell carcinoma.

cutis, 42:359.

Rosenthal, K.L.; Palmer, J.L. and Harris, J.A. (1975):

Antibody-induced redistribution of CEA on the cell surface : utilization in separation of CEA and isoantigen.

Am. J. Immunol. 115: 1049.

Sankowski, A.; Janik, P. and Jeziorska, M. (1987):

The results of topical application of 13-cis-retinoic acid on basal cell carcinoma. A correlation of the clinical effect with histopathological examination and serum retinal level.

Neoplasia, 34: 485.

Schurch, W.; Seemayer, T.A.; Lagace, R. and Gabbiani, G. (1984):

The intermediate filament cytoskeleton of myofibroblasts an immunofluorescence and ultrastructural study.

Virch. Arch., 403:3233.

Sell, S. (1990):

Tumor markers in Diagnostic pathology, cancer markers of the 1990, Comparison of the new generation of markers Defined by monoclonal antibodies and oncogene probes to prototypic markers.

Clin. Lab. Med., 10:1.

Sheibani, K.; Battitora, M. and Winberg, C.D. (1986):

Further evident that malignant angioendotheliomatosis is an angiotropic large cell lymphoma.

N. Engl. J. Med., 314:943.

Sherbet, G.V. (1970):

Epigenetic processes and their relevance to the study of neoplasia.

Advances in Cancer Res., 97:167.

Sibley, R.K. and Dahl, D. (1985):

Neuroendocrine carcinoma of the skin. An immunocytochemical study of 21 cases.

Am. J. Surg. Pathol., 9: 109.

Sommerville, J. and Milne, J. (1950):

Familial primary self-healing squamous epithelioma of the skin.

Br. J. Dermatol., 62: 485.

Stefansson, K.; Wollman, R.L. and Jenkovic, M. (1982):

S 100 protein in soft tissue tumors derived from schwann cells and melanocytes.

Am. J. Pathol., 106:261.

Steven, D.P. and Mackay, I.R. (1973):

Increased carcinoembryonic antigen in heavy cigarette smokers.

Lancet, 2:1238.

Stoll, H.J. (1979):

Squamous cell carcinoma. In: Text book dermatology in general medicine. Fitzpatrick, T.P.; Eisen, A.Z.; Wolff, C.; Freedberg, I. and Ousten, K.F. (eds), McGraw-Hill, New York, P. 362.

Terry, W.D.; Henkart, P.A.; Coligan, J.E. and Todd, C.W.
(1972):

Structural studies of the Major glycoprotein in
preparations with carcinoembryonic antigen activity.
J. Ex. Med., 136:200.

Terry, W.D.; Henkart, P.A.; Caligan, J.E. and Todd, C.W.
(1979):

Carcinoembryonic antigen characterization and clinical
applications.
Trans. Rev., 20:100.

Thomas, P. and Zamcheck, N. (1983):

Role of the liver in clearance and excretion of
circulating CEA.
Dig. Dis. Sci., 28:216.

Thomson, D.M.P.; Krupey, J.; Freedman, S.O. and Gold, P.
(1969):

The radioimmunoassay of circulating CEA of the human
digestive system.
Proc. Soc. Nat. Acad. Sci., 64: 616.

Thorpe, S.M.; Lykkasfelt, A.E.; Vinterbyl A. and Lons Dorfer.
M. (1986):

Quantitative immunological detection of estrogen receptor
in nuclear pellet from human breast cancer biopsy.
Cancer Res., 46: 4251.

Toker, C. (1972):

Trabecular carcinoma of the skin.
Arch. Dermatol., 105:107.

Tormey, D.C.; Davis, T.E. and Woalkes, T.P. (1982):

Tumor markers. In: Principles of cancer treatment. Carter S.K.; Glaststein E. and Livingston R.B. (eds). MacGraw-Hill, New York, P.170.

Tosca, A.D.; Lehou, J. and Panagiotopoulos, A. (1986):

Antigenic differences and further perspectives in the differential diagnosis of skin tumors.
J. Dermatol., 13: 366.

Turner, J. and Callen, J. (1981):

Aggressive behavior of squamous cell carcinoma in a patient with preceding lymphocytic lymphoma.
J. Am. Acad. Dermatol., 4: 446.

Turner, R.R.; Wood, G.S.; Beckstead, J.H.; Colby, T.V.;

Aorning, S.J. and Warnke, R.A. (1984):
Enzymatic heterogeneity.
Am. J. Surg. Pathol., 8: 485.

Van Nagell, J.R.; Mecker, W.R.; Parker, J.C. and Harralson, J.D. (1975):

Carcinoembryonic antigen in Patients with gynaecologic malignancy.
Cancer, 35:1372.

Van Scott, F.J. (1979):

Basal cell carcinoma. In: Text book dermatology in General medicine. Fitzpatrick, T.P.; Eisen, A.Z.; Wolff, C.; Freedberg, I.M. and Ousten, K.F. (eds) McGraw-Hill, New York, P. 377.

Varnaar, S.O.; Van Muijen, G.N.P.; Ruiter, D.J.; Ponac, M. and Huiskens-VanderMey, C. (1984):

Monoclonal antibodies with different specificities against cytokeratins, an immunohistochemical study of normal tissues and tumors.

Am. J. Pathol., 114:9.

Vasarinsh, P. (1982):

Squamous cell carcinoma. In: Clinical dermaotology Vasarinsh, P. (ed). Butter warth publishers, Boston, London, P. 548.

Vibra, R.; Alpert, E. and Isselbacher, K.J. (1975):

Carcinoembryonic antigen: Evidence for multiple antigenic determinants and isoantigens.

Proc. Nati. Acad. Sci. U.S.A., 72: 4602.

Virji, M.A.; Mercer, D.W. and Herberman, R.B. (1988):

Tumor markers in cancer Diagnosis and prognosis.

Cancer J. Clin., 38 :104.

Wade, T.R. and Ackerman, A.B. (1978):

The many faces of basal cell carcinoma.
J. Dermatol. Surg. Oncol., 4: 23.

Wagener, C. and Breuer, H. (1982):

Biochemical and immunochemical aspects of CEA and CEA related antigen: Current status and future perspectives.
J. Clin. Chem. Clin. Biochem., 20:705.

Wagner, R.F. and Cottle, W.I. (1990):

Treatment of squamous and basal cell carcinomas of the hand with mohs micrographic surgery.
Dermatol. and Venereol., 47: 66.

Ward, D.E.; Von Kuster, L.C. and Liwnicz, B.H. (1985):

Concanavalin A and lens culinaris binding pattern in normal and Neoplastic sebaceous epithelium.
Lab. Invest., 52:73.

Warner, T.F.; Uno, H. and Hafez, R. (1983):

Merkel cells and merkel cell tumors. Ultrastructure immunocytochemistry and Review of the literature.
Cancer, 52:238.

Wick, M.R. and Coffin, C.M. (1985):

Sweat gland and pilar carcinoma. In: Pathology of unusual Malignant cutaneous tumors. Wick, M.R. (ed).
Marcel Dekker, New York., P. 1.

Wick, M.R.; Kaye, V.N.; Sibley, R.K.; Tyler, R. and Frizzera, G. (1986):

Primary neuroendocrine carcinoma and small-cell malignant lymphoma of the skin: a discriminant immunohistochemical comparison.

J. Cutan. Pathol., 13:347.

Wick, M.R. and Manivel, J.C. (1986):

Epithelioid sarcoma and isolated necrobiotic granuloma : a comparative immunohistochemical study.

J. Cutan. Pathol., 13:253.

Wick, M.R. and Scheithauer, B.W. (1985):

Primary neuroendocrine carcinoma of the skin. In: Pathology of Unusual Cutaneous Tumors. Wick, M.R. (ed). Marcel Dekkey, New York, P. 107.

Wood, G.S.; Beckstead, J.H.; Turner, R.R.; Hendrickson, M.R.;

Kempson, R.L. and Warnke, R.A. (1986):

Malignant fibrous histiocyte tumor cells resemble fibroblasts.

Am. J. Surg. Pathol., 10:323.

Wood, M.G.; Pranich, K. and Becrman, H. (1958):

Investigation of possible apocrine gland component in basal cell epithelioma.

J. Invest. Dermatol., 30: 273.

Quoted from Miller, S.J. (1991).

Wu, J.T.; Herbst, J.J. and Bray, P.F. (1976):

Elevated plasma levels of carcinoembryonic antigen in cystic fibrosis.

Pediatr. Res., 10:235.

Yedema, C.; Kenemans, P.; Wobbes, T.; Op'tRoot, E.; Werheijen, R.; Poels, L. and Thomas, S. (1987):

Pretreatment serum levels of tumor-associated trypsin in 180 gynecological and surgical oncology patients. A comparison with other modern and classical markers.

Int. J. Cancer, 32: 718.

Zackheim, H.S. (1963):

Origin of the human basal cell epithelioma.

J. Invest. Dermatol., 40: 283.

Quoted by Miller, S.J. (1991).

Zarbo, R.J.; Crissman, J.D.; Venkat, H. and Weiss, M.A. (1986):

Spindle-cell carcinoma of the upper aerodigestive tract mucosa.

Am. J. Surg. Pathol., 10:741.

Zelickson, A.S.; Goltz, R.W. and Hartmann, J.F. (1961):

A histologic and electron microscopic study of basal cell epithelioma.

J. Invest. Dermatol., 136: 299.