

REFERENCES

- Ackermann - Schopf, C.M.A. (1976): Antibodies to basal epidermal cells in patients with basal or squamous cell carcinoma. *Arch. Dermatol. Res.*, 257 :149.
- Aggarwal,B.B., Eessalu,T.E. and Hass,P.E.(1985): Characterization of receptors for human tumor necrosis factor and their regulation by γ -interferon. *Nature*, 318 : 665.
- Aguet, M. (1980): High affinity binding of 125 I-labelled mouse interferon to a specific cell surface receptor. *Nature*, 284 : 459.
- Aguet,M., Groebke,M. and Dreiding, Ph.(1983): Down- regulation of interferon receptors. In:*The Biology of the interferon system 1983*.De Maeyer, E and Schellekens, H.(eds).Elsevier Science Publishers, Amsterdam. NewYork. Oxford., p161.
- Aguet, M. and Merlin,G.(1987): Purification of human γ - interferon receptors by sequential affinity chromatography on immobilized monoclonal antireceptor antibody and humen γ - interferon. *J.Exp.Med.*, 165 : 988.
- Aiyer, R.A., Serrano,L.E. and Jones, P.P.(1986): Interferon- γ binds to high and low affinity receptor components on murine macrophages.*J. Immunol.*, 136 : 3329 .
- Alkan, S.S., Towbin, H. and Hochkeppel,H-K. (1988): Enhanced antiproliferative action of interferon targeted by bispecific monoclonal antibodies. *J.Interferon Res.*, 8 : 25 .
- Alton,K., Stabinsky,Y., Richards,R., Ferguson,B., Goldstein,L., Altrock,B., Miller,L. and Stebbing,N. (1983): Production, Characterzation and biological effect of recombinant DNA derived human IFN- α and IFN- γ . analogs. In: *The Biology of the interferon system 1983*. De Maeyer,E., and Schellekens,H.(eds).Elsevier Science Publishers, Amsterdam. NewYork. Oxford., p.119
- Amon , R.B. and Goodkin , P.E.(1976):Topical 5- fluorouracil and the basal cell nevus syndrome. *N.Engl.J.Med.*, 295 : 677.
- Anderson, P., Yip, Y. K. and Vilcek, J. (1982) : Specific binding of 125 I-human interferon - γ to high affinity receptors on human fibroblasts. *J.Biol.Chem.*, 257 (19) : 11301.
- Antonelli , G., Currenti , M., Turriziani, O. and Dianzani, F. (1991) : Neutralizing antibodies to interferon - α : Relative frequency in patients treated with different interferon preparations.. *J.Infect.Dis.*, 163 : 882.

- Arnheiter, H. and Meier, E. (1990) : Mx proteins: Antiviral proteins by chance or by necessity ? *New Biol.*, 2 : 851.
- Barker, J.N.W.N., Allen , M.H. and Mc Donald, D.M. (1989) : The effect of *in vivo* interferon-gamma on the distribution of LFA -1 and ICAM-1 in normal human skin. *J Invest.Dermatol.*, 93 : 439.
- Barker, J.N.W.N., Allen, M.H. and Mc Donald,D.M. (1990) : Alterations induced in normal human skin by *in vivo* interferon - γ . *Br. J. Dermatol.*, 122:451.
- Baron, S., Tying, S.K., Fleischmann, W.R., Coppenhaver. D. H ., Niesal, D.W., Kimpel, G.R.,Stanton,G.J. and Hughes, T.K. (1991) : The interferons : Mechanisms of action and clinical application. *JAMA.* , 266 (10) : 1375 .
- Basham, T.Y. and Merrigan, T.C. (1983) : Recombinant interferon - γ increase HLA-DR synthesis and expression . *J. Immunol.*, 130:1492.
- Basham, T.Y., Nickoloff, B. J. , Merigan, T.C.and Morhenn,V.B.(1984a) : Recombinant - γ interferon induces HLA - DR expression on cultured human keratinocytes. *J.Invest .Dermatol.*, 83 : 88.
- Basham, T.Y., Smith, W. K. and Merigan, T.C. (1984b) : Interferon enhances antibody - dependent cellular cytotoxicity when suboptimal concentrations of antibody are used. *Cell Immunol.*, 88 : 393.
- Baylic, H.I. and Cies, W.A. (1975) : Complications of Mohs'chemosurgical excision of eyelid and canthol tumors. *Am.J.Ophthalmol.*, 80 : 116.
- Belardelli, F., Vignaux, F., Proietti E. and Gresser, I. (1984) : Injection of mice with with antibody to interferon renders peritoneal macrophages permissive for vesicular stomatitis virus and encephalomyocarditis virus (EMV) *Proc. Natl.Acad.Sci.USA.*, 81:602. Quoted from : Gresser, I (1990).
- Berglund, O., Engman, K., Ehrnst, A., Andersson, J., Lidman, L., Åkerlund, B., Sönnernborg, A. and Stannegård, Ö. (1991) : Combined treatment of symptomatic human immunodeficiency virus type 1 infection with native interferon - α and zidovudine *J.Infec.Dis.* , 163 : 710.

- Berman , B. and Duncan, R. (1989) : Short-term keloid treatment in *vivo* with human interferon alfa-2b results in a selective and presistent normalization of keloidal fibroblast collagen , glycosaminoglycan and collagenase production in *vitro*. *J.Am.Acad.Dermatol.*, 21 : 694.
- Beutler, B. (1990) : TNF in pathophysiology : Biosynthetic regulation. *J.Invest.Dermatol.* , 95 : 81s.
- Beutler, B. and Cerami, A. (1986) : Cachectin and tumour necrosis factor as two sides of the same biological coin. *Nature.*, 320 : 584.
- Beutler, B., Tkacenko, V., Milsark, I., Krochin, N. and Cerami, A., (1986): Effect of γ – interferon on cachectin expression by mononuclear phagocytes. *J.Exp.Med.*, 164 : 1791.
- Blalock, J. E., Georgiades, J. A., Langford, M. P. and Johnson, H. M. (1980): Purified human immune interferon has more potent anticellular activity than fibroblast or leukocyte interferon. *Cell Immunol.*, 49 : 390.
- Blewitt, R.W. (1980) : Why does basal carcinoma metastasize so rarely? *Int.J.Dermatol.*, 19 : 144.
- Blewitt, R.W., Aparicio, S. R. and Burrow, H. M. (1978) : Failure to detect circulating IgG or IgM antibodies to basal carcinoma by immunofluorescence. *Acta. Dermatol.Venereal.* , 58 : 82.
- Bocci, V. (1985) : The physiological interferon response. *Immunol.Today*, 6 : 7.
- Boguniewicz, M., Jaffe, H.S., Sullivan, M.J., York,D., Geha, R.S. and Leung, D.Y. M. (1990) : Recombinant gamma interferon in treatment of patients with atopic dermatitis and elevated IgE levels. *Am.J.Med.*, 88 : 365.
- Bollag,W. and Ott, F. (1975) : Vitamin A acid in benign and malignant epithelial tumors of the skin . *Acta Dermatol.Venereol.*, 74 : 163.
- Boneschi, V., Brambilla, L., Mozzanica, N., Cattaneo, A. and Finzi, A.F. (1989) : Treatment of basal cell carcinoma with intralesional alpha 2b recombinant interferon (Abstr.). *J.Invest.Dermatol.*, 93 : 542.

- Boutin, C., Viallat,J.R. and Van Zandwijk, N.,(1991) : Activity of intrapleural recombinant gamma- interferon in malignant mesothelioma. *Cancer*, 67 : 2033.
- Branca, A. and Baglioni, C. (1981) : Evidence that types I and II interferons have different receptors. *Nature*, 294 : 768.
- Bridgman, R., Rossi,C.R. and Campos, M. (1988) : The sensitivity of domestic animal cell lines to eight recombinant human interferons. *J Interferon Res.*, 8 : 1.
- Briggaman , R.A. (1982) : Biochemical composition of the epidermo-dermal junction and other basement membrane. *J.Invest Dermatol .*, 78 : 1.
- Brown, S. L., Miller, R. A., Horning,S. J., Czerwinski,D., Hart,S.M., Mc Elderry, R., Basham,T., Warnke, R. A ., Merigan, T.C. and Levy ,R . (1989) : Treatment of B-cell lymphomas with anti-idiotypic antibodies alone and in combination with alpha IFN. *Blood* , 73 : 651.
- Buechner, S.A. (1991) : Intralesional interferon alfa-2b in the treatment of basal cell carcinoma: Immunohistochemical study on cellular immune reaction leading to tumor regression . *J.Am.Acad.Dermatol.*, 24 (5) : 731.
- Bunn, P.A., Foon, K. A., Ihde, D.C., Longo, D. L., Eddy, J., Winkler, C. F., Veach, S.R., Zeffren, J., Sherwin, S. and Oldham,R. (1984) : Recombinant leukocyte A interferon : An active agent in advanced cutaneous T-cell lymphomas.*Ann.Int.Med.*, 101 : 484.
- Bunn, P.A., Ihde, D.C. and Foon , K.A. (1986) : The role of recombinant interferon alpha - 2a in the therapy of cutaneous T-cell lymphomas. *Cancer.*, 57 : 1689.
- Bunn, P.A. and Norris, D.A. (1990) : The therapeutic role of interferons and monoclonal antibodies in cutaneous T-cell lymphomas. *J.Invest.Dermatol.*, 95 : 209s.
- Camp, R.D.R., Fincham, N.J., Rosst, J.S., Bacon, K.B. and Gearing, A.J.H (1990): Leukocyte chemoattractant cytokines of the epidermis. *J.Invest.Dermatol.*, 95 : 108s.
- Carrel, S., Schmidt-Kessen, A. and Giuffre', L.(1985) : Recombinant interferon-gamma

- can induce the expression of HLA-DR and -DC on DR-negative melanoma cells and enhance the expression of HLA-ABC and tumor-associated antigens. *Eur.J.Immunol.*, 15:118. Quoted from: Mahrle, G., and Schuler, H. (1990).
- Carton, C.A. and Kilbourne, E.D (1952) : Activation of latent herpes simplex by trigeminal sensory - root section *N. Engl. J .Med.*, 246 : 172. Quoted from: HO, M. (1990).
 - Chebath, J., Benech, P., Hovanessian, A., Galabru, J. and Revel, M. (1987) : Four different forms of interferon-induced (2', 5') oligo(A) synthetase identified by immunoblotting in human cells. *J.Biol.Chem.*, 262 : 3852.
 - Chomyn, A., Cleeter, M.W. J., Ragan, C. I., Riley, M., Doolittle, R. F. and Attardi, G. (1986) : URF6 last unidentified reading frame of human mt DNA, codes for an NADH dehydrogenase subunit. *Science*, 234 : 614.
 - Clemens, M. J. and Mc Nurlan, M. A. (1985) : Regulation of cell proliferation and differentiation by interferons. *Biochem.J.*, 226 : 345.
 - Contente, S., Kenyon, K., Rimoldi, D. and Friedman, R.M. (1990) : Expression of gene *rrg* is associated with reversion of NIH 3T3 transformed by LTR-c-H-ras. *Science*, 249 : 796.
 - Cornell, R.C. (1992) : Reply : Intralesional injection therapy for basal cell carcinoma. *J.Am.Acad. Dermatol.*, 26(1) : 143.
 - Cornell, R.C., Greenway, H.T., Tucker, S. B., Edwards, L., Ashworth, S., Vance, C., Tanner, D.J., Taylor, E.L., Smiles, K. A. and Peets, E. A. (1990) : Intralesional interferon therapy for basal cell carcinoma. *J.Am.Acad.Dermatol.*, 23(4) : 694.
 - Coskey, R.J. and Mehregan, A.H., (1987) : The association of basal cell carcinoma with other tumors. *J.Dermatol. Surg.Oncol.*, 13 : 553.
 - Covelli, A., Cavalieri, R. and Coppola, G. (1987) : Recombinant leukocyte. interferon as initial therapy in mycosis fungoides and sezary syndrome (Abstr.). *Proc. Am. So. Clin. Oncol.*, 6 : 187. Quoted from: Bunn, P.A. and Norris, D.A. (1990).

- Creagan, E.T., Sohaid, D.J. , Ahmann, D.L. and Frytak, S. (1990) : Disseminated malignant melanoma and recombinant interferon: Analysis of seven consecutive phase II investigations. *J.Invest.Dermatol.*, 95 : 188s.
- Creasey,A A. and Merigan ,T.C. (1982): Advances in interferon research.*West. J.Med.*, 136 : 227.
- Cristofolini,M., Zumiani,G. and Scappini,P. (1984) : Aromatic retinoid in the chemoprevention of the progression of nevoid basal-cell carcinoma syndrome. *J.Dermatol. Surg.Oncol.*, 10 : 778.
- Czaja,M.J., Weiner,F.P. ,Eghbali, M., Giambrone,M.A., Eghbli,M.and Zern,M.A. (1987) : Differential effects of gamma-interferon on collagen and fibronectin gene expression. *J.Biol Chem.*, 262:13348.
- Czarniecki, C.W., Fennie, C.W., Powers, D.B. and Estell, D.A. (1984) : Synergistic antiviral and antiproliferative activities of *Escherichia coli* -derived human alpha,beta,and gamma interferon.*J.Virol.*, 49 : 490.
- Damme, J.V. and Opdenakker, G. (1990) : Interaction of interferons with skin reactive cytokines : From interleukin-1 to interleukin-8. *J.Invest.Dermatol.* , 95:90s.
- Davis, L.T.C.C. (1991) : HIV-specific treatment. *Dermatologic Clinics.*, 9 : 585.
- DeGrado, W.F., Wasserman, R. and Chowdhry, V. (1982) : Sequence and structural homologies among type I and type II interferons. *Nature.*, 300 : 379.
- Dellon, A.L. (1978) : Host tumor relationships in basal cell and squamous cell cancer of the skin. *Plast Réconstr. Surg.*, 62 : 37 Quoted from : Miller, S.J. (1991 II) .
- Detmar, M., Imcke, E., Ruszczak, Z. and Orfanos, C.E. (1990) : Effect of recombinant tumor necrosis factor - alpha on cultured microvascular endothelial cells derived from human dermis. *J. Invest.Dermatol.*, 95 : 219s.
- Detmar, M. and Orfanos, C.E. (1990) : Tumor necrosis factor-alpha inhibits cell proliferation and induces class II antigens and cell adhesion molecules in cultured normal human keratinocytes in vitro. *Arch.Dermatol.Res.*, 282 : 238.

- Dianzani, F., Antonelli, G., Amicucci, P., Cefaro, A. and Pintus, C. (1989) : Low incidence of neutralizing antibody formation to interferon α 2b in human recipients. *J. Interferon Res.*, 9 : 33s.
- Dianzani, F., Buckler, C.E. and Baron, S. (1969) : Effect of Cycloheximide on the antiviral action of interferon. *Proc.Soc.Exp.Biol.Med.*, 130 : 519 Quoted from: Richtsmeier, W.J. and Grossberg, S.E. (1989).
- Domzing, W., Stadler, B. M. and Herberman, R. B. (1983) : Interleukin-2 dependence of human natural killer (NK) cell activity. *J.Immunol.*, 130 : 1970.
- Dougherty, T. J. (1981) : Photoradiation therapy for cutaneous and subcutaneous malignancies. *J.Invest dermatol.*, 77 : 122.
- Douglas, J .M., Eron, L.J., Judson, F. N., Rogers, M., Alder, M.B., Taylor, E., Tanner, D. and Peets, E. (1990) : A randomized trial of combination therapy with intralesional interferon α 2b and podophyllin versus podophyllin alone for the therapy of anogenital warts. *J.Infect.Dis.*, 162 : 52.
- Dover, J.S. and Johnson, R. A. (1991) : Cutaneous manifestations of human immunodeficiency virus infection. *Arch.Dermatol.*, 127 : 1383.
- Drake, L.A., Ceilley, R.I., Cornelison, R.L., Dobes, W.A., Dorner, W., Goltz, R.W., Lewis, C.W., Salasche, S.J. and Turner, M.L.C. (1992) : Guidelines of care for basal cell carcinoma. *J.Am.Acad.Dermatol.*, 26 : 117.
- Dreiding, P., Staeheli, P. and Haller, O. (1985) : Interferon-induced protein Mx accumulates in nuclei of mouse cells expressing resistance to influenza viruses. *Virology*, 140 : 191.
- Dubin, N. and Kopf, A.W. (1983) : Multivariate risk score for recurrence of cutaneous basal cell carcinomas. *Arch.Dermatol.*, 119 : 373.
- Duncan, M.R. and Berman, B. (1985) : Gamma interferon is the lymphokine and beta interferon the monokine responsible for inhibition of fibroblast collagen production and late but not early fibroblast proliferations. *J.Exp.Med.*, 162:516.

- Duncan, M.R. and Berman, B.(1987) : Persistence of a reduced - collagen producing phenotype cultured scleroderma fibroblasts after short term exposure to interferons. *J.Clin.Invest.*, 79 : 1318.
- Duncan,M.R. and Berman, B.(1989) : Differential regulation of glycosaminoglycan, fibronectin and collagenase production in cultured dermal fibroblasts by interferon-alpha, -beta, and - gamma . *Arch.Dermatol.Res.*, 281:11.
- Dunnick, J.K. and Galasso, G.J. (1979) : Clinical trials with exogenous interferon : Summary of a meeting. *J. Infect. Dis.*, 139 : 109.
- Edwards , L. (1987) : Interferon. *Arch.Dermatol.*, 123 : 743.
- Edwards , L., Levine, N., Weidner, M., Piepkorn, M. and Smiles, K.A. (1986) : Effect of intralesional α_2 -interferon on actinic keratoses. *Arch.Dermatol.*, 122:779.
- Edwards, L., Tucker, S. B., Perednia, D., Smiles, K. .A., Taylor, E. L., Tanner, D. J. and Peets, E. (1990 a) : The Effect of an intralesional sustained-release formulation of interferon alpha-2b on basal cell carcinomas. *Arch Dermatol.*, 126 : 1029.
- Edwards, L.,Whiting, D., Rogers, D., Luck, K. and Smiles, K. A. (1990 b) : The effect of intralesional interferon gamma on basal cell carcinomas. *J.Am.Acad.Dermatol.*, 22(3) : 496.
- El-Komy, M., El-Nasr, H.S., El-Ansari, M., Sobeih, S., Saleh, N.F. and Zaher, H. A. (1992) : Immunology of basal cell epithelioma. *Egypt. J. Dermatol. & Venereol.*, 12(1) : 37.
- Epstein, E. (1992) : Intralesional interferon therapy for basal cell carcinoma. *J. Am. Acad.Dermatol.*, 26(1):142.
- Eron, L.J., Judson, F., Tucker, S., Prawer, S., Mills, J. ,Murphy, K., Hickey, M., Rogers, M., Flannigan,S., Hien, N., Katz, H.I., Goldman, S., Gottlieb, A., Adams, K., Burton, P., Tanner,D., Taylor, E. and Peets, E. (1986) : Interferon therapy for condylomata acuminata. *N. Engl.J. Med.*, 315(17) : 1059.

- Ezekowitz, R. A. B., Dinanuer, M.C., Jaffe, H.C., Orkin, S.H. and Newburger, P. E. (1988) : Partial correction of the phagocyte defect in patients with X-linked chronic granulomatous disease by subcutaneous IFN- γ . *N.Engl.J.Med.*, 319 : 146.
- Faltynek, C.R., Mc Candless, S., Chebath, J. and Baglioni, C. (1985) : Different mechanisms for activation of gene transcription by interferons alpha and gamma. *Virology* , 144 : 173.
- Farmer, E. R. and Helwig, E. B. (1980) : Metastatic basal cell carcinoma : A clinicopathologic study of seventeen cases. *Cancer*, 46 : 748.
- Farrar, W.L., Johnson, H.M. and Farrar, J.J. (1981) : Regulation of the production of immune interferon and cytotoxic T lymphocytes by interleukin - 2. *J.Immunol.*, 126:1120
- Fassio, A., Gariglio, M., Cofano, F., Cavallo, G. and Landolfo, S. (1988) : Functional characterization of murine cell lines expressing high, intermediate, or negative levels of surface receptors for interferon- γ . *J.Interferon Res.*, 8 : 333.
- Fierlbeck, G. and Rassner, G. (1990) : Treatment of psoriasis and psoriatic arthritis with interferon gamma. *J.Invest.Dermatol.*, 95 : 138s.
- Finbloom, D.S., Hoover, D.L. and Wahl, L.M. (1985) : The characteristics of binding of human recombinant interferon- γ to its receptor on human monocytes and human monocyte- like cell lines. *J.Immunol.*, 135:300.
- Fitzgerald, P.A., Von Wusson, P. and Lopez, C. (1982) : Role of interferon in natural killing of HSV-1 infected fibroblasts. *J.Immunol.* , 129 : 819.
- Fleischmajer, R. and Nedwich , A. (1972) : Generalized morphea .I.Histology of the dermis and subcutaneous tissue. *Arch.Dermatol.*, 106 : 509.
- Fleischmajer, R., Perlish, J.J. and Duncan, M. (1983) : Scleroderma : A model for fibrosis. *Arch.Dermatol.*, 119 : 957.
- Flenniken, A. M., Galabru, J., Rutherford,M. N., Hovanessian, A.G. and Williams, B. R. G. (1988) : Expression of interferon-induced genes in different tissues of mice. *J.Virol.*, 62 : 3077.

- Fossa, S.D., Gunderson, R. and Moe, B. (1990) : Recombinant interferon - alpha combined with prednisone in metastatic renal cell carcinoma : Reduced toxicity without reduction of response rate-a phase II study . *Cancer*, 65 : 2451.
- Friedland, G.H. , Klein, R.S. and Saltzman, B.R. (1988) : The interferon alpha study group. A randomized placebo- controlled trial of recombinant human interferon alpha 2a in patients with AIDS. *J.Acquir Immune Defic.Sydr.*, 1 : 111.
- Friedman-Kien, A.E., Eron, L.J., Conant, M., Growdon, W., Badiak, H., Bradstreet, P.W., Fedorczyk, D., Trout, R. and Plasse, T.F. (1988): Natural Interferon alfa for treatment of condylomata acuminata. *JAMA.*, 259 (4) : 533.
- Friedman-Kien, A.E., Klein, R.J., Glaser, R.D. and Czelusniak, S.M. (1986): Treatment of recurrent genital herpes with topical alpha interferon gel combined with nonoxynol-9. *J.Am. Acad. Dermatol.*, 15 : 989.
- Friedman, R.M. and Sonnabend, J.A. (1965) : Inhibition of interferon action by puromycin. *J.Immunol.*, 95 : 696. Quoted from: Richtsmeier, W.J. and Grossberg, S.E. (1989).
- Friesel, R., Komorly, A. and MacLag, T. (1987) : Inhibition of endothelial cell proliferation by gamma-interferon. *J.Cell Biol.*, 104 : 689.
- Fukui, Y., Iseki, R. and Ohashi, M. (1988): Lymphokine-Activated killer (LAK) activity against autologous malignant tumors of the skin . *J.Invest Dermatol.*, 919 : 319.
- Galabru, J., Robert, N., Buffer-Janvresse, C., Riviere, Y. and Hovanessian, A.G. (1985) : Continuous production of interferon in normal mice : Effect of anti-interferon globulin, sex, age, strain, and environment on the levels of 2'-5' synthetase and p 67K kinase *J.Gen.Virol.*, 66 : 711.
- Galand, P. and Heenen, M. (1988) : DNA replication fork progression rate and temporal organization of S phase in normal epidermis and in basal cell carcinoma. *J.Cell.Physiol.*, 137 : 374.

- Galbraith, G. M.P. (1988) : Therapeutic immunomodulation. *Dermatol.Clinic.*, 6(4) : 561.
- Garbe, C., Krasagakis, K., Zouboulis, Ch.C., Schröder, K., Krüger, S., Stadler, R. and Orfanos, C.E. (1990): Antitumor activities of interferon alpha, beta, and gamma and their combinations on human melanoma cells in vitro : Changes of proliferation melanin synthesis, and immuno-phenotype . *J. Invest. Dermatol.*, 95 : 231s.
- Garbe, C., Schröder, K., Krasagakis, K., Zouboulis, Ch.C., Krüger, S. and Schwartingl, R. (1989) : Presence of ICAM-1 on melanoma cells in vitro and enhancement of its expression by interferons. *J Invest. Dermatol.*, 93 : 550.
- Geffen, J.R., Klein, R.J. and Friedman-kien, A.E. (1984) : Intralesional administration of large doses of human leukocyte interferon for the treatment of condylomata acuminata. *J. Infect. Dis.*, 150 : 612.
- Gelfant, S.(1976) : The cell cycle in psoriasis : A reappraisal. *Br J Dermatol.*, 95 : 577.
- Geronemus, R. and Ashinoff, R. (1991) : Lasers in the treatment of skin cancer. *Dermatol.Clin* ., 9(4) : 765.
- Glashan, R.W .(1990) : A randomized controlled study of intravesical alpha 2b interferon in carcinoma in situ of the bladder. *J.Urol* ., 144 : 658.
- Glezerman, M., Cohen, V. and Movshovitz, M. (1988) : Placebo controlled trial of topical interferon in labial and genital herpes. *Lancet*, i : 150.
- Goldfarb, M.T., Gupta, A.K., Gupta, M. A . and Sawchuk , W.S. (1991) : Office therapy for human papilloma virus infection in nongenital sites. *Dermatol. Clin.*, 9 : 287.
- Goldring, M.B., Sandell, L.J., Stephenson, M.L. and Krane, S.M. (1986) : Immune interferon suppresses levels of procollagen mRNA and type II collagen synthesis in cultured human articular and costal chondocytes. *J.Biol.Chem.*, 261 : 9049.
- Goldstein, D. and Laszlo, J. (1986) : Interferon therapy in cancer. From imagination to interferon. *Cancer Res.*, 46 : 4315.

- Gooding, C.A., White,G. and Yatsuhashi, M., (1965): Significance of marginal extension in excised basal cell carcinoma . *N. Engl.J Med.* 273 : 923. Quoted from: Cornell, R.C. (1992).
- Goudsmit,J.,De Wolf,F. and Paul, D.A.(1986):Expression of human immuno deficiency virus antigen (HIV -Ag) in serum and cerebrospinal fluid during acute and chronic infection. *Lancet*, 2 : 177.
- Graham-Brown , R. (1991) : Benign and malignant skin tumours. *Pract. East. Med .*,2 : 161.
- Granstein, R.D., Deak, M.R. , Jacques, S.L., Margolis, R.J., Flotte , T.J. Whitaker, D., Long, F.H. and Amento, E.P. (1989) : The systemic administration of gamma interferon inhibits collagen synthesis and acute inflammation in a murine skin wounding model. *J.Invest.Dermatol.*, 93 : 18.
- Granstein, R.D., Flotte, T.J. and Amento, E.P. (1990 a) : Interferons and collagen production *J.Invest.Dermatol.*, 95 : 75s.
- Granstein, R.D., Rook, A., Flotte, T.J., Haas, A., Callo, R.L., Jaffe., H.S. and Amento E.P. (1990 b) : A controlled trial of intralesional recombinant interferon γ in treatment of keloidal scarring. *Arch.Dermatol.*, 126 : 1295.
- Grau, G. E., Taylor, T.E., Trop, M., Molyneux, M.E., Wirima, J.J., Vassalli, P., Hommel, M. and Lambert, P -H. (1989) : Tumor necrosis factor and disease severity in children with falciparum malaria *N.Eng.J Med.*, 320 : 1586.
- Gray, P.W. and Goddel, D.V.(1982) : Structure of the human immune interferon gene. *Nature*, 298 : 859.
- Greenway, H.T. and Cornell, RC. (1990) : Interferon. Coming of age . *Arch . Dermatol.*, 126 : 1080.
- Greenway, H.T., Cornell, R.C., Tanner, D.J., Peets, E., Bordin, G.M. and Nagi, C. (1986) : Treatment of basal cell carcinoma with intralesional interferon *J.Am.Acad. Dermatol.*, 15 : 437.

- Gresser, I. (1990) : Biologic effects of interferons. *J.Invest.Dermatol.*, 95 : 66s.
- Gresser, I., Maury, C., Woodrow, D., Moss, J., Grütter, M.G., Vignaux, F., Belardelli, F. and Maunoury, M.T., (1988) : Interferon treatment markedly, inhibits the development of tumor metastases in the liver and spleen and increases survival time of mice after intravenous inoculation of Friend erythroleukemia. *Int. J.Cancer*, 41:135.
- Gresser, I., Vignaux, F, Belardelli, F., Tovey, M.G. and Maunoury, M.T. (1985) : Injection of mice with antibody to mouse interferon α / β decreases the level of 2'-5'-oligoadenylate synthetase in peritoneal macrophages *J. Virol.*, 53 : 221 .
- Gribaudo, G., Lembo, D., Landolfo, S. and Lengyel,P. (1991) : Interferon action : Binding of viral RNA to the 40 -kilodalton 2'-5'- oligoadenylate synthetase in interferon - treated hela cells infected with encephalo-myocarditis virus (EMCV). *J. Virol.*, 65(4) : 1748.
- Grimwood, R.E., Ferris, C.F. and Mercill, D.B. (1986) : Proliferating cells of human basal cell carcinoma are located on the nodules. *J. Invest. Dermatol.*, 86: 191.
- Grimwood, R.E., Glanz, S.M. and Siegle, R.J. (1988) : Transplantation. of human basal cell carcinoma. to C 57 / BALB / Cbg. J /bg J-nu / nu (beige - nude) mouse. *J. Dermatol. Surg. Oncol.*, 14 : 59.
- Grob, J.J. , Collet, A.M. , Munoz , M.H. and Bonerandi, J. J. (1988) : Treatment of large basal - cell carcinomas with intralesional interferon alpha 2a.*Lancet*, 16:878.
- Groopman, J.E., Gottlieb, M.S., Goodman, J., Mitsuyasu, R.T. , Conant, M.A., Prince, H. , Fahey, J.L. , Derezin, M. , Weinstein, W.M., Casavante, C., Rothman, J. , Rudnick, S.A. and Volberding, P.A. (1984) : Recombinant alpha - 2 interferon therapy for kaposi's sarcoma associated with the acquired immunodeficiency syndrome. *Ann.Int. Med.*, 100 : 671.
- Groopman, J.E. and Scadden, D.T. (1989) : Interferon therapy for kaposi's sarcoma associated with the acquired immunodeficiency syndrome (AIDS).*Ann.Intern Med.*, 110 : 335.

- Gross, G., Roussaki, A. and Papendick, U. (1990) : Efficacy of interferons on bowenoid papulosis and other precancerous lesions. *J. Invest. Dermatol.*, 95: 152s.
- Gross, G., Roussaki, A. and Schöpf, E. (1986) : Successful treatment of condylomata acuminata and bowenoid papulosis with subcutaneous injections of low - dose recombinant interferon - α . *Arch. Dermatol.*, 122:749.
- Guillen, F.J., Day, C.L. and Murphy, G.F. (1985) : Expression of activation antigens by T- cells infiltrating basal cell carcinomas. *J. Invest. Dermatol.*, 85 : 203.
- Guthrie, T.H., Me Elveen, L.J. and Porubsky, E.S (1985): Cisplatin and doxorubicin. An effective chemotherapy in the treatment of advanced basal cell and squamous carcinoma of skin. *Cancer*, 55 : 1629.
- Gutterman, J.U., Blumenschein, G.R., Alexanian, R., Yap, H-Y., Buzdar, A.U., Cabanillas, F., Hortobagyi, G.N., Hersh, E.M. , Rasmussen, S.L., Harmon M., Kramer, M. and Pestka, S. (1981) : Leukocyte interferon - induced tumor regression in human metastatic breast cancer, multiple myeloma, and malignant lymphoma. *Ann.Intern.Med.*, 93 : 399.
- Gutterman, J.U., Fein, S. and Quesada, J. (1982) : Recombinant leukocyte A interferon : Pharmacokinetics single - dose tolerance and biologic effects in cancer patients. *Ann. Intern.Med.*, 96 : 546.
- Guyre, P.M., Morganelli, P.M. and Miller, R.(1983) : Recombinant immune interferon increases immunoglobulin G Fc receptors on cultured human mononuclear phagocytes. *J. Clin.Invest.*, 72 : 393.
- Habets, J.M., Tank, B. and Vuzevski, V.D., (1988) : Characterization of mononuclear infiltrate in basal cell carcinoma : A predominantly T-cell-mediated immune response with minor participation of Leu-7 ⁺ (natural Killer) cells and Leu - 14 ⁺ (B) cells. *J. Invest. Dermatol.*, 9: 289.
- Hales, S.A., Stamp, G., Evans, M. and Fleming, K.A.(1989) : Identification of the origin of cells in human basal cell carcinoma xenografts in mice using *in situ* hybridization. *Br.J. Dermatol.*, 120 : 351.

- Härfast, B., Huddleston, J.R., Casali, P., Merigan, T.C. and Oldstone, M.B.A.(1981): Interferon acts directly on human B lymphocytes to modulate immunoglobulin synthesis. *J. Immunol*, 127 : 2146.
- Harms, G., Zwingenberger, K., Chehade, A.K., Talhari, S., Racz, P., Mouakeh, A., Douba, M., Nakel, L., Naiff, R.D. , Kremsner, P.G., Feldmeier, H. and Bienzle, U. (1989) : Effects of intradermal gamma- interferon in cutaneous leishmaniasis. *Lancet*, 10 : 1287.
- Hawkins, M.J., Krown,S.E., Krim, E.C.B.M., Real, F.X., Edwards,B.S., Anderson, S.A., Cunningham-Rundles, S.C. and Oettgen, H.F (1984) : American cancer society phase I trial of naturally produced β - interferon. *Cancer Res.* , 44 : 5934.
- Heenen, M., Achten,G. and Ganland, P.(1973) : Autoradiographic analysis of cell kinetics in human normal epidermis and basal cell carcinoma. *Cancer Res.*, 33 : 123.
- Heenen, M. and Galand, P. (1980) : Decreased rate of DNA-chain growth in human basal cell carcinoma. *Nature*, 285 : 265.
- Herberman, R.B., Ortaldo, J.R. , Mantovani, A., Hobbs, D.S., Kung, H-F. and Pestka, S.(1982) : Effect of human recombinant interferon on cytotoxic activity of natural killer (NK) cells and monocytes. *Cell. Immunol.*, 67 : 160.
- Hershman, M.J., Sonnenfeld, G., Logan,W.A., Pietsch, J.D., Wellhausen,S.R. and Polk,H.C.(1988) : Effect of interferon-gamma treatment on the course of a burn wound infection . *J.Interferon.Res.*, 8 : 367.
- Ho, D.D., Hartshorn, K.L., Rota , T.R., Andrews, C.A., Kaplan,J.C., Schooley,R.T. and Hirsch, M.S. (1985) : Recombinsnt human interferon alpha - A suppresses HTLV-III replication in vitro. *Lancet*, 16 : 602.
- Ho,M.(1982) : Recent advances in the study of interferon . *Pharmacol. Rev.*, 34 : 119.
- Ho,M. (1990) : Interferon as an agent against herpes simplex virus . *J. Invest. Dermatol.*, 95 : 158s.

- Hochster,H., Dieterich,D., Bozzette, S., Reichman, C., Connor, J.D., Liebes, L., Sonke,R.L., Spector, S.A., Valentine, F., Pettinelli,C. and Richman, D.D. (1990) : Toxicity of combined Ganciclovir and Zidovudine for cytomegalovirus disease associated with AIDS. *Ann. Int.Med.*, 113 : 111.
- Hodak, K., Ginzburg, A. and David, M. (1987) : Etreinate treatment of nevoid basal cell carcinoma syndrome. Theraputic and chemopreventative effect. *Int.J.Dermatol.*, 26 : 606.
- Hokland, P. and Berg, K.(1981) : Interferon enhances the antibody - dependent cellular cytotoxicity (ADCC) of human polymorphonuclear leukocytes. *J. Immunol.*, 127 : 1585.
- Horisberger, M.A. and De Staritzky, K. (1989) : Expression and stability of the Mx protein in different tissues of mice, in response to interferon inducers or to influenza virus infection . *J. Interferon Res.*, 9 : 583.
- Houghton, A.N., Thomson ,T.M., Gross, D., Oettgen , H.F. and Old, L.J. (1984): Surface antigens of melanoma and melanocytes : Specificity of induction of Ia antigens by human γ - interferon . *J Exp. Med.*, 160 : 255.
- Hughes, T.K. and Baron.S. (1987) : Do the IFNs act singly or in combination ? *J Interferon Res.*, 7 : 603.
- Iric, D., Padovan, I., Brodarec, I., Knezevic, M.and Soos, E. (1981) : Application of human leucocyte interferon in patients with tumours of head and neck. *Lancet*, 1 : 1025.
- Isaacs, A., Klemperer, H.G. and Hitchcock , G. (1961): Studies on the mechanism of action of interferon . *Virology*, 13 : 191. Quoted from : Richtsmeier, W.J. and Grossberg, S.E. (1989).
- Isaacs, A. and Lindenmann, J. (1957): Viral interference . 1. The interferon. *Proc . R. Soc Lond . [Biol].*, 147 : 258. Quoted from : Trofatter, K.F. (1991).

- Jackson , G.G., Paul,D.A., Falk, A., Rubenis, M., Despotes, J.C., Mack, D., Knigge , M. and Emeson,E.E. (1988): Human immunodeficiency virus (HIV) antigenemia (p24) in the acquired immunodeficiency syndrome (AIDS) and the effect of treatment with zidovudine (AZT). *Ann. Int . Med.*, 108 : 175.
- Jacobs, B.L., Miyamoto, N.G. and Samuel , C.E. (1988) : Mechanism of interferon action : Studies on the activation of protein phosphorylation and the inhibition of translation in cell free systems . *J.Interferon Res.*, 8 : 617.
- Jacobsen, H. (1986) : Interferons and antiviral activity . *Arzeim Forsch Drug Res.*, 36 :512. Quoted from : Baron , S., Tying, S.K., Fleischmann , W.R, Coppenhaver , D.H., Niesal, D.W., Kimpel, G.R.,Stanton , G.J. and Hughes T.K.(1991).
- Jaffe, H.S., Chen, A.B., Kramer, S. and Sherwin, S.A. (1987) : The absence of interferon antibody formation in patients receiving recombinant human interferon - gamma. *J. Biol. Res . Modif.* 6:576. Quoted from : Mahrle , G. and Schultz, H. (1990) .
- Jakschies, D., Hochkeppel, H-K., Horisberger, M. A., Deicher, H. and Von Wussow , P. (1990) : Correlation of the antiproliferative effect and the Mx - homologous protein induction by IFN in patients with malignant melanoma. *J. Invest. Dermatol.*, 95 : 238s.
- Jimenez, S.A., Freundlich, B. and Rosenbloom, J. (1984) : Selective inhibition of human diploid fibroblast collagen synthesis by interferons . *J. Clin. Invest.*, 74 : 1112.
- Johnson, V. A., Merrill, D.P., Videler, J.A., Chou, T-C., Byington , R.E., Eron , J.J. ; D'Aquila , R.T. and Hirsch, M.S.(1991) : Two- Drug combinations of zidovudine , didanosine, and recombinant interferon – α . A inhibit replication of zidovudine - resistant human immunodeficiency virus type I synergistically. *J. Infect. Dis.* , 164 : 646.

- Jones, G.J. and Itri, L.M. (1986) : Safety and tolerance of recombinant interferon alpha - 2a. (*Roferon-A*) in cancer patients. *Cancer Res.*, 57:1709.
- Kahan, A., Amor, B., Menkes., C.J. and Strauch, G. (1989): Recombinant interferon- γ in the treatment for systemic sclerosis. *Am. J. Med.*, 87 : 273 .
- Kaplan, G. , Luster, A.D. , Hancock, G. and Cohn, Z. A. (1987 a) : The Expression of a γ - interferon-induced protein (IP - 10)in delayed immune responses in human skin. *J. Exp. Med.*, 166 : 1098.
- Kaplan, G., Nusrat, A. , Sarno, E.N., Job, C.K., Mc Elrath, J., Porto, J.A., Nathan, C.F. and Cohn, Z.A. (1987 b) : Cellular responses to the intradermal injection of recombinant human γ - interferon in lepromatous leprosy patients. *Am. J. Pathol.*, 128 (2) : 345.
- Kaplan, J.E., Spira, T.J., Fishbein, D.B., Bozeman, L.H., Pinsky, P.F. and Schonberger, L.B. (1988) : A six - year follow-up of HIV-infected homosexual men with lymphadenopathy. Evidence for an increased risk for developing AIDS after the third year of lymphadenopathy. *JAMA.*, 260 : 2694.
- Kapp, A. and Zeck-Kapp, G. (1990) : Activation of the oxidative metabolism in human polymorphonuclear neutrophilic granulocytes; The role of immuno-modulating cytokines. *J. Invest. Dermatol.*, 95 : 94s.
- Kawase, I., Urdal. D.L., Newman, W. and Henney, C.S. (1983) : The mechanism of augmentation of natural killer cell activity by syngeneic tumor cells : Role of macrophage-derived factor in NK boosting. *Int. J. Cancer*, 31: 365.
- Kerr, I. M. and Martin, E.M. (1971) : Virus protein synthesis in animal cell-free systems : Nature of the products synthesized in response to ribonucleic acid of encephalomyocarditis virus. *J. Virol.*, 7:438. Quoted from : Richtsmeier, W.J. and Grossberg, S.E. (1989).

- Kerr, M.I., Cayley, P.J., Roberts, W.K., Rice, A., Reid, A., Hersh, C., Stark, G.R., Laurence, L., Cailla, H., Marti, J., Welles, V. and Mallucci, L. (1983) : Molecular mechanisms of interferon action and the possible wider significance of the 2-5 A system. In: *The Biology of The Interferon System 1983*. De Maeyer, E. and Schellekens, H. (eds). Elsevier science Publishers, Amsterdam. New York .Oxford. p 213.
- Kirkwood, J.M. and Ernstoff, M.S. (1990) : Role of interferons in the therapy of melanoma. *J. Invest. Dermatol.*, 95 : 180s.
- Kirkwood, J.M. , Ernstoff, M.S., Davis, C.A., Reiss, M., Ferraresi, R. and Rudnick, S.A. (1985) : Comparison of intramuscular and intravenous recombinant alpha -2 interferon in melanoma and other cancers. *Ann. Int. Med.*, 103 : 32..
- Knobler, R.M., Trautinger, F., Radaszkiewicz, T., Kokoschka, E.M. and Micksche, M. (1991) : Treatment of cutaneous T - cell lymphoma with a combination of low - dose interferon alfa-2b and retinoids. *J.Am. Acad. Dermatol.*, 24 (2) : 247.
- Knop, J. (1990) : Immunologic effects of interferon. *J. Invest. Dermatol.*, 95 : 72 s.
- Knop, J., Riechmann. R. and Macher. E. (1981):Modulation of suppressor mechanisms in allergic contact dermatitis. I. Effect of *C. parvum* on the induction phase of contact allergy. *J. Invest Dermatol.*, 76 : 193.
- Knop. J., Stremmer. R., Neumann, Ch., De Maeyer, E. and Macher, E.(1982) : Interferon inhibits the suppressor T - cell response of delayed - type hypersensitivity. *Nature*, 296 : 775.
- Kohase, M., Zhang. Y., Lin, J - X., Yamazaki, S., Sehgal, P.B. and Vilcek, J. (1988) : Interleukin - 1 can inhibit interferon - β synthesis and its antiviral action : Comparison with tumor necrosis factor *J. Interferon Res.*, 8 : 559.
- Kohn, E.C., Steis, R. and Sausville E. (1990) : Phase II trial of intermittent high dose recombinant interferon alpha 2a in mycosis fungoides and sezary syndrome. *J. Clin. Oncol.*, 8:155.
- Kokoschka, E-M., Trautinger, F., Knobler, R.M., Pohl-Markl, H. and Micksche, M. (1990) : Long-term adjuvant therapy of high - risk malignant melanoma with interferon - α 2b. *J. Invest. Dermatol.*, 95 : 193s.

- Kowalzick, L., Weyer, U., Lange, P. and Breitbart, E.W. (1990) :Systemic therapy of advanced metastatic malignant melanoma with a combination of fibroblast interferon- β and recombinant interferon - γ . *Dermatologica*, 181 : 298 .
- Kronenberg, L.H., Rosenblatt, H.M. Bryson, Y. and Merigan, T.C. (1982) Interferon : immunobiology and Clinical significance. *Ann. Intern. Med.*, 96 : 80.
- Krown, S.E., Real, F.X. and Cunningham - Rundles, S.(1983): Preliminary observations on the effect of recombinant leukocyte A-interferon homosexual men with kaposi's sarcoma. *N. Engl.J.Med.*, 308 : 1071.
- Krust, B., Galabru, J. and Hovanessian, A.G. (1984) : Further characterization of the protein kinase activity by interferon in mouse and human cells. *J. Biol. Chem.*, 259:8494.
- Kurzrock, R., Feinberg, B., Talpaz, M., Saks,S. and Gutterman, J.U.(1989) : Phase I study of a combination of recombinant tumor necrosis factor - α and recombinant interferon - γ in cancer patients. *J. Interferon Res.*, 9:435.
- Kurzrock, R., Quesada, J.R., Talpaz,M., Hersh, E.M., Reuben, J.M., Sherwin, S.A. and Gutterman, J.U. (1986) : Phase I study of multiple dose interamuscularly administered recombinant gamma interferon. *J.Clin. Oncol.* , 4: 1101.
- Kurzrock, R., Rosenblum, M.G., Sherwin, S.A., Rios, A., Talpaz, M., Quesada, J.R. and Gutterman, J.U.(1985): Pharmacokinetics, single - dose tolerance, and biological activity of recombinant γ - interferon in cancer patients. *Cancer Res.*, 45 : 2866 .
- Kusari, J. and Sen, G.C. (1986) : Regulation of synthesis and turnover of an interferon - induced mRNA. *Mol.Cell. Biol.*, 6:2062.
- Kusari, J. and Sen, G.C. (1987) : Transcriptional analysis of interferon - induced mRNAs. *Mol.Cell. Biol.*, 7:528.
- Landow , R.K. (1988) :The interferons:A clinician's view. *Dermatol.Clin.*, 6 (4) : 569.
- Lane, H.C., Davey, V. and Kovacs, J.A. (1990) : Interferon alpha in patients with asymptomatic human immunodeficiency virus (HIV) infection : A randomized, placebo-controlled trial. *Ann. Intern. Med.*, 112 : 805.

- Lane, H.C., Kovacs, J.A., Feinberg, J., Herin, B., Davey, V., Walker, R., Deyton, L., Metcalf, J.A., Baseler, M., Salzman, N., Manischewitz, J., Quinnan, G., Masur, H. and Fauci, A.S. (1988) : Anti-retroviral effects of interferon α in AIDS - associated kaposi's sarcoma. *Lancet*, 26 : 1218.
- Lang, M.H., Lodemann, E. and Holzmann, H. (1988) : Serum interferon level and (2'-5') - oligo (A) synthetase activity. *Z. Hautkr*, 63 : 1028. Quoted from: Miller, S.J. (1991) .
- Laskin, O.L., De Miranda, P. and Blum, MR (1989) : Azidothymidine steady - state pharmacokinetics in patients with AIDS and AIDS-related complex. *J. Infect. Dis.*, 159:745.
- Legha, S.S. (1986) : Interferons in the treatment of malignant melanoma. A review of recent trials. *Cancer.*, 57 : 1675.
- Leigh, I.M., Eady, R.A.J. and Heagerty, A.H.M. (1988) : Type VII collagen is a normal component of epidermal basement membrane, which shows altered expression in recessive dystrophic epidermolysis bullosa. *J. Invest. Dermatol.*, 90 : 639.
- Lengyel, P. (1982) : Biochemistry of interferons and their actions. *Ann . Rev. Biochem.*, 51:251.
- Lengyel, P. (1987) : Double - stranded RNA and interferon action. *J. Interferon Res.*, 7:511.
- Lever, A.M.L, Brook, M.G., Yap, I. and Thomas, H.C. (1987) : Treatment of thrombocytopenia with alpha interferon. *Br. Med. J.*, 295 : 1519.
- Levine, N., Miller, R.C. and Meyskens, F.L. Jr. (1984) : Oral isotretinoin therapy : Use in a patient with multiple cutaneous squamous cell carcinomas and Keratoacanthomas. *Arch Dermatol.*, 120 : 1215.
- Levine, S. (1964) : Effect of actinomycin D and puromycin dihydrochloride on action of interferon. *Virology*, 24: 586. Quoted from : Richtsmeier, W.J. and Grossberg, S.E. (1989) .
- Lober, C.W. and Fenske, N.A. (1991) : Basal cell, squamous cell, and sebaceous gland carcinoma of the periorbital region. *J. Am. Acad. Dermatol.*, 25 (4) : 685.

- Luger, T.A. and Schwarz, T. (1990) : Evidence for an epidermal cytokine network. *J. Invest Dermatol.*, 95 : 100s.
- Luster, A.D., Unkeless, J. and Ravetch, J.V. (1985) : γ -interferon transcriptionally regulates an early response gene containing homology to platelet proteins . *Nature*, 315 : 672.
- Mahrle, G. and Schulze, H-J. (1990) : Recombinant interferon - gamma (r IFN $-\gamma$) in dermatology. *J. Invest. Dermatol.*, 95 : 132s.
- Maluish, A.E. , Ortaldo, J.R., Conlon, J.C., Sherwin, S.A., Leavitt, R., Strong, D.M., Weirnik, P., Oldham. R.K. and Herberman, R.B. (1983) : Depression of natural killer cytotoxicity after *in vivo* administration of recombinant leukocyte interferon. *J. Immunol.*, 131 : 503.
- Manara, G.C., Ferrari, C. and Torresani, C. (1989) : Dermal infiltrates in basal cell carcinoma (letter). *J. Am. Acad. Dermatol.*, 20 : 701.
- Markey, A.C., Tidman, M.J., Churchill, L.J., Aplin, J.D., Leigh, I.M., Purkis, P. and Macdonald, D.M. (1991) : The epidermal basement membrane in basal cell carcinoma : An immunohistochemical study. *Br. J. Dermatol.*, 125 : 21 .
- Maroteau, L., Chen, L., Mitrani - Rosenbaum, S., Howley, P. M. and Revel, M. (1983) : Cycloheximide induces expression of the human interferon B, gene in mouse cells transformed by bovine papillomavirus - interferon B, recombinants. *J. Virol.*, 47 : 89
- Marziali, G., Fiorucci, G., Coccia, E.M., Percario, Z., Raber, J., Battistini, A., Rossi, G.B., Affabris, E. and Romeo, G. (1991) : Posttranscriptional regulation of Beta - interferon expression in erythroid friend cells treated with gamma interferon. *J. Virol.*, 65 : 4130.
- Mc Daniel, W.E. (1986) : Adequate follow up for basal cell carcinoma. *Arch. Dermatol.*, 122 : 243.
- Mc Innes, B., Chambers, P.J., Cheetham, B.F., Beilharz, M.W. and Tymms, M.J. (1989) : Structure - Function studies of interferon - α : Amino acid substitutions at the conserved residue tyrosine 123 in human interferon - α_1 . *J. Interferon Res.*, 9 : 305.

- Mc Kenzie, R.C. and Sauder, D.N. (1990) : The Role of Keratinocyte cytokines in inflammation. *J. Invest. Dermatol.*, 95 : 105s.
- McLeod, G.R., Thomson, D.B. and Hersey, P. (1990): Clinical evaluation of interferons in malignant melanoma. *J. Invest. Dermatol.*, 95 : 185s.
- Meigel, W., Kern, P., Plettenberg, A. and Dietrich, M. (1989) : Experience with interferon therapy of HIV-related kaposi's sarcoma in hamburg (Abstr). *J. Invest. Dermatol.*, 93 : 564.
- Merigan, T.C., Gallagher, J.G., Pollard, R.B. and Arvin, A.M. (1981) : Short - course human leukocyte interferon in treatment of herpes zoster in patients with cancer. *Antimicrob. Agent. Chemother.*, 19 : 193. Quoted from : Ho, M. (1990) .
- Meyskens, F.L. Jr., Gilmartin, E. and Alberts, D.S. (1982) :Activity of isotretinoin against SCC., and preneoplastic lesion. *Cancer. Treat. Rep.*, 66 : 1315. Quoted from : Robinson, J.K. (1991).
- Miles, S., Marcus, S. and Mitsuyasu, R. (1989) : Antiviral activity and low incidence of opportunistic infections in patients with poor prognosis AIDS - related kaposi's sarcoma treated with betaser interferon (Abster.). *J. Invest. Dermatol.*, 93 : 565.
- Miller, S.J. (1991 I) :Biology of basal cell carcinoma (Part I). *J. Am. Acad. Dermatol.*, 24 (1) : 1 .
- Miller, S. J. (1991 II) : Biology of basal cell carcinoma (Part II). *J. Am. Acad. Dermatol.*, 24 (2) : 161 .
- Miyakoshi, J., Dobler, K.D., Allalunis -Turner, J., Mc Kean, J.D.S., Petruk, K., Allen, P.B.R., Aronyk, K.N., Weir, B., Huyser - Wierenga, D., Fulton, D., Urtasun, R.C. and Day, R.S. (1990): Absence of IFN A and IFN B genes from human malignant glioma cell lines and lack of correlation with cellular sensitivity to interferons. *Cancer Res.*, 50:278.
- Mohs, F.E. (1986) : Micrographic surgery for the microscopically controlled excision of eyelid cancers. *Arch. Ophthalmol.*, 104 : 901.

- Morel, P., Rybcjad, M. and Borradori, L. (1989) : Therapy of classic kaposi sarcoma with recombinant interferon - alpha 2b. (Abstr.). *J. Invest. Dermatol.*, 93 : 566.
- Morfeldt-Månson, L, Böttiger, B., Nilsson, B. and Von Stedingk, L.V. (1991) : Clinical signs and laboratory markers in predicting progression to AIDS in HIV - 1 infected patients. *J. Infect. Dis.*, 23 : 443 .
- Morhenn, V.B., Pregerson - Rodan, K., Henzel - Mullen, R., Wood, G.S., Nickoloff, B.J., Sherwin, S.A. and Farber, E.M. (1987) : Use of recombinant interferon gamma administered intramuscularly for the treatment of psoriasis. *Arch. Dermatol.*, 123 : 1633.
- Morimoto, Y. , Saga, K., Bando, M. and Takahashi, M. (1991) : In - vitro DNA synthesis of keratinocytes in normal human skin, psoriasis, seborrhoeic keratosis, bowen's disease and basal cell carcinoma. *Br. J. Dermatol.*, 125 : 9 .
- Morley, S.M., Rennison, A. and Macadam, R.F. (1983) : Membrane - bound structures at the interface between tumor and in basal cell carcinoma. An ultrastructural study. *J. Pathol.*, 140 : 267.
- Moss, AR. (1988) : Predicting who will progress to AIDS. *Br. Med. J.*, 297 : 1067.
- Murray, H.W. (1988) : Interferon – γ , the activated macrophage, and host defense against microbial challenge. *Ann. Intern. Med.*, 108 : 595.
- Murray, J.C., Pollack, S.V. and Pinnell, S.R. (1981) : Keloids : A review. *J. Am. Acad. Dermatol.*, 4 : 461.
- Myskowski, P.L., Wattenberg, D. and Flomenberg, N. (1988) : Inhibition of natural killer cell activity by basal cell carcinoma extract [Abstr.]. *J. Invest Dermatol.*, 90 : 591.
- Nathan, C.F., Kaplan, G., Levis, W.R., Nusrat, A., Witmer M.D., Sherwin, S.A., Job, G.K., Horowitz, C.R., Steinman, R.M. and Cohn, Z.A. (1986) : Local and systemic of intradermal recombinant interferon – γ in patients with lepromatous leprosy. *N.Engl. J. Med.*, 315 (1) : 6.

- Nathan, C.F., Murray, H.W., Wiebe, M.E. and Rubin, B.Y. (1983) : Identification of interferon- γ as the lymphokine that activates human macrophage oxidative metabolism and antimicrobial activity. *J. Exp. Med.*, 158 : 670.
- Nickoloff, B.J. (1987) : Binding of 125 I - gamma interferon to cultured human keratinocytes *J. Invest. Dermatol.*, 89 ; 132 .
- Nickoloff, B.J. (1989) : Role of interferon - γ in cutaneous trafficking of lymphocytes with emphasis on molecular and cellular adhesion events. *Arch. Dermatol.*, 124 : 1835.
- Nickoloff, B.J., Basham, T.Y. and Merigan, T.C. (1985) : Immunomodulatory and antiproliferative effect of recombinant alpha, beta, gamma interferons on cultured human malignant squamous cell line, SCL-1 and SW - 1271. *J. Invest. Dermatol.*, 84 : 487.
- Nickoloff, B.J. and Griffiths, C.E.M. (1989) : T-lymphocytes and monocytes bind to keratinocytes in frozen sections of biopsy specimens of normal skin treated with gamma interferon. *J. Am Acad. Dermatol.*, 20 : 736.
- Nickoloff, B.J., Mahrle, G. and Morhenn, G.B. (1986) : Ultrastructural effects of recombinant gamma interferon on cultured human keratinocytes. *Ultrastruct. Pathol.*, 10 : 17. Quoted from : Mahrle, G. and Schulze, H-J. (1990).
- Nickoloff, B.J., Mitra, R.S., Elder, J.T., Fisher, G.J. and Voorhees, J.J. (1989) : Decreased growth inhibition by recombinant gamma interferon is associated with increased transforming growth factor - alpha production in keratinocytes cultured from psoriatic lesions. *Br. J. Dermatol.*, 121 : 161.
- Nicolas, J-F., Thivolet, J. , Kanitakis, J. and Lyonnet, S. (1990) : Response of discoid and subacute cutaneous lupus erythematosus to recombinant interferon alpha 2A. *J. Invest. Dermatol.*, 95 : 142s.
- Niimura, M. (1990) : Application of Beta - interferon in virus - induced papillomas. *J. Invest. Dermatol.*, 95 : 149s.

- Nilsson, H., Johansson, C., Sandberg, K., Funa, K., Alm, G.V. and Scheynius, A. (1989) : Induction of mRNA for HLA-DR beta in human keratinocytes cocultured with interferon - gamma. *Arch. Dermatol. Res.* , 281 : 260.
- Norris, D.A. (1990) : Cytokine modulation of adhesion molecules in the regulation of immunologic cytotoxicity of epidermal targets. *J. Invest. Dermatol.*, 95:111s. -
- Novick, D., Fischer, D.G., Reiter. Z., Eshhar, Z. and Rubinstein, M. (1989) : Monoclonal antibodies to the human interferon - γ receptor : Blocking of the biological activities of interferon - γ and purification of the receptor. *J. Interferon Res.*, 9 : 315.
- Novick, D., Orchansky, P., Revel, M. and Rubinstein, M. (1987) : The human interferon- γ receptor. *J. Biol. Chem.* , 262 : 8483.
- Oberman, F. and Panet, A. (1989) : Characterization of the early steps of herpes simplex virus replication in interferon - treated human cells. *J. Interferon Res.* , 9 : 563.
- Olsen, E.A., Rosen, S.T. and Vollmer, R.T. (1989) : Interferon alpha 2a in the treatment of cutaneous T cell lymphoma. *J. Am. Acad. Dermatol.*, 20 : 395.
- Oratz, R. (1986) : Interferons. *Melanoma Lett.*, 4 : 1.
- Oratz, R. and Bystryn. J-C. (1991) : Immunotherapy of malignant melanoma. *Dermatol .Clin.* , 9 : 669.
- Orchansky, P., Goren, T. and Rubinstein, M. (1983) : Isolated subtypes of human interferon - α : Synergistic action and affinity for the receptor. *In : The Biology of The Interferon System 1983*. De Maeyer, E. and Schellekens, H (eds). Elsevier Science Publishers, Amsterdam. NewYork. Oxford. p. 183.
- Orholm, M ., Pedersen, C., Mathiesen, L., Dowd, P. and Nielsen, J.O. (1989) : Suppression of p 24 antigen in sera from HIV - infected individuals with low - dose alpha - interferon and zidovudine : A pilot study. *AIDS.*, 3 : 97.
- Ozzello, L., Habif, D.V., Rosa, C.M.D. and Cantell, K. (1988) : Effect of intralesional injections of interferons - α on xenografts of human mammary carcinoma cells (BT 20 and MCF - 7). *J. Interferon Res.*, 8 : 207 .

- Pandey, R.; Ziboh, V.A.; Berman, B. and Duncan, M. (1985) : Gamma - interferon induced inhibition of eicosanoid synthesis in human skin. *Clin. Res.* , 33 : 155 A.
Quoted from : Mahrle, G., and Schultz, H. J. (1990) .
- Pazin, G.J., Armstrong, J.A., Lam, M.T., Tarr, G.C., Jannetta, P.J. and Ho, M. (1979) : Prevention of reactivated herpes simplex infection by human leukocyte interferon after operation on the trigeminal root. *N. Engl. J. Med.*, 301 : 225.
- Pearlman, D.L. (1991) : Weekly pulse dosing : Effective and comfortable topical 5-fluorouracil treatment of multiple facial actinic keratoses. *J. Am. Acad. Dermatol.*, 25 (4) : 665.
- Pennica, D., Nedwin, G.E., Hayflick, J.S., Seeburg, P.H., Derynck, R., Palladino, M.A., Koher, W.J., Aggarwal, B.B. and Goeddel, D.V. (1984) : Human tumor necrosis factor : Precursor structure, expression and homology to lymphotoxin. *Nature*, 312 : 724.
- Pestka, S. (1983) : The human interferons from protein purification and sequence to cloning and expression in bacteria: Before, between, and beyond . *Arch. Biochem. Biophys.*, 221 : 1. Quoted from : Stadler, R., Mayer-da- Silva, A., Bratzke, B., Gorbe, C. and Orfanos, C. (1989) .
- Pestka, S., Langer J.A., Zoon, K.C. and Samuel, C.E. (1987) : Interferons and their actions. *Ann . Rev. Biochem.*, 56 : 727.
- Petersen, C.S., Albrechtsen, J., Larsen, J., Sindrup, J., Tikjøb, G., Ottevanger, V., Karlsmark, T. , Fogn, H. , Mogensen, A. M. and Wolff- Sneedorff, A. (1991) : Subclinical human papilloma virus infection in condylomata acuminata, patients attending a VD clinic. *Acta . Dermatol. Venerol.*, 71:252.
- Philip, P. and Epstein, L.B. (1986) : Tumor necrosis factor as immunomodulator and mediator of monocyte cytotoxicity induced by itself, γ -IFN and interleukin-1. *Nature*, 323:86.

- Piguet, P-F., Grau, G.E. , Allet, B. and Vassalli, P. (1987) : Tumor necrosis factor / Cachectin is an effector of skin and gut lesions of the acute phase of graft- vs - host disease. *J.Exp. Med.*, 166: 1280.
- Pinnell, S.R.(1981): Regulation of collagen synthesis. *J.Invest. Dermatol* , 79:73s -
- Pinocy, J. and Hopt, U.T. (1989): Interferon-gamma. A potent inducer of local inflammation. *J. Invest. Dermatol.*, 93:569 .
- Poher, J.S., Gimbrone, M.A. Jr., Lapierre, L.A., Mendrick, D.L., Fiers, W., Rotherlein, R. and Springer, T.A. (1986) Overlapping patterns of activation of human endothelial cells by interleuking I, tumor necrosis factor and immune interferon , *J. Immunol.*, 137: 1893
- Poumay, Y. , Boucher, F. , Degen, A. and Leloup, R . (1991) : Inhibition of basal cell proliferation during storage of detached cultured epidermal keratinocyte sheets. *Acta. Dermatol .Venereol.*, 71: 195.
- Proctor, S.J. , Jackson, G. , Garey, P. and Stark,A. (1988) : Short - course alpha - interferon therapy in severe unresponsive immune thrombocytopenic purpura . *Lancet.* , 23 : 947 .
- Quesada, J.R. and Gutterman J .U. (1986) : Psoriasis and alpha - interferon. *Lancet*, 28: 1466.
- Rachlis, A.R. (1989) : Antiviral therapy . *Clin. Dermatol.* , 7 (3) : 69.
- Rapp, F. and Wrzos, H. (1985) : Synergistic effect of human leukocyte interferon and nonoxynal-9 against herpes simplex type2. *Antimicrob. Agents. Chemother.*, 28 : 449. Quoted from : Shupack, J., Stiller, M., Knobler, E.,Ackerman, C., Jpndreau, L. and Kenny,C. (1990).
- Rashidbaigi, A. , Stefanos, S. , Jung, V. and Langer, J.A. (1988) : Immune interferon receptor : Chemical and enzymatic sensitivity . *J. Interferon Res.*, 8:641 .
- Ratain, M.J., Golomb, H.M., Vardiman, J.W., Westbrook, C.A., Barker, Ch., Hooberman, A., Bitter, M.A. and Daly, K. (1988) : Relapse after interferon alpha -

- 2b therapy for hairy-cell leukemia : Analysis of prognostic variables. *J. Clin. Oncol.*, 6:1714.
- Real, F.X., Oettgen, H.F. and Krown, S.E. (1986) : Kaposi's sarcoma and the acquired immunodeficiency syndrome treatment with high and low doses of recombinant leukocyte A interferon. *J. Clin. Oncol.*, 4: 544.
 - Rehberg, E., Kelder, B., Hoal, E.G. and Pestka, S. (1982) : Specific molecular activities of recombinant and hybrid leukocyte interferons. *J. Biol. Chem.*, 257:11497.
 - Reinhold, U., Wehrmann, W., Kukel, S. and Kreysel, H.W. (1989) : Defective IFN- γ production in hyper-IgE atopic dermatitis patients : Evidence for an intrinsic abnormality. (Abstr.) *J. Invest. Dermatol.*, 93:571.
 - Reitamo, S., Komulainen, M., Iiliius, P. and Grohn, P. (1989) : Treatment of basal cell carcinoma with intralesional interferon alpha 2b (Abstr.) *J. Invest. Dermatol.*, 93 : 572.
 - Reymann, F. (1985) : 15 years experience with treatment of basal cell carcinomas of the skin with curettage. *Acta. Dermatol. Venereol.*, 120: 56 s.
 - Reynolds, J.E.F., Parfit, K., Parsons, A.V. and Sweetman, S.C. (1989) : Interferons
In: *Martindale. The Extra Pharmacopeia*. Reynolds, J.E.F., Parfit, K., Parsons, A. V. and Sweetman, S.C. (eds) . The Pharmaceutical Press, London. p.696 .
 - Richtsmeier, W.J. and Grossberg, S.E. (1989) : Inhibitory effect of mitochondrial metabolic inhibitors on interferon action. *J. Interferon Res.*, 9: 87
 - Ringenberg, Q.S. and Anderson, P.C. (1986) : Interferons in the treatment of skin disease. *Int. J. Dermatol.*, 25 (5) : 273 .
 - Ritz, J. (1989) : The role of natural killer cells in immune surveillance [Editorial] . *N.Engl. J. Med.*, 320: 1748 .
 - Robbins, P. (1978): Chemosurgery : My 15 years of experience. *J. Dermatol. Surg. Oncol.*, 4 : 41.

- Roberts, R.M. (1989) : A novel group of interferons associated with the early ovine and bovine embryo. *J. Interferon Res.*, 9: 373.
- Robinson, J.K. (1987) : Use of a combination of chemotherapy and radiation therapy in the management of advanced basal cell carcinoma of the head and neck. *J. Am. Acad. Dermatol.*, 17:770
- Robinson, J.K.(1991) : Advances in the treatment of nonmelanoma skin cancer. *Dermatol. Clin.*, 9 (4) : 757 .
- Roenigk, H.H. , Kuzel, T.M. , Skoutelis, A.P. , Springer, E. , Yu, G. , Caro, W. , Gilyon, K. , Variakojis, D. , Kaul, K. , Bunn, P.A. , Evans, L. and Rosen, S.T. (1990) : photochemotherapy alone or combined with interferon alpha - 2 a in the treatment of cutaneous T- cell lymphoma. *J. Invest. Dermatol.*, 95: 198s .
- Roenigk, R.K., Ratz, J.L. and Bailin, P.L. (1986): Trends in the presentation and treatment of basal cell carcinoma. *J Dermatol. Surg. Oncol.*, 12: 860 .
- Romeo, G., Affabris, E , Frederico, M., Jemma, C., Belardelli, F., Mechti, N., Gresser, I. and Rossi, G. B. (1983) : 2' - 5' A synthetase activity does not increase in IFN- resistant friend leukemia cell. Variants treated with α / β IFN despite the presence of high affinity IFN receptor sites In : *The Biology of The Interferon System 1983* . De Maeyer. E., and Schellekens, H. (eds) . Elsevier Science Publishers., Amsterdam . NewYork . Oxford. p. 207
- Rönblom, L.E. , Alm, G.V. and Öberg, K.E. (1991) : Autoimmunity after alpha - interferon therapy for malignant carcinoid tumors. *Ann. Int. Med.* , 115: 178.
- Rowe, D.E., Carroll, R.J. and Day, CL. (1989): Long-term recurrence rates in previously untreated (primary) basal cell carcinoma : implications for patient follow-up. *J. Dermatol Surg. Oncol.*, 15: 315 .
- Rozenbaum, W., Gharakhanian, S. , Nevarette, S-M. , De Sahb, R., Cardon, B. and Rouzioux, C. (1990) : Long-term follow-up of 120 patients with AIDS-related kaposi's sarcoma treated with interferon alpha - 2a. *J. Invest. Dermatol.*, 95 : 161.

- Ruszczak, Z., Detmar, M. , Imcke, E. and Orfanos, C.E. (1990) : Effects of rIFN alpha, Beta , and gamma on the morphology, proliferation, and cell surface antigen expression of human dermal microvascular endothelial cells *in vitro* . *J. Invest . Dermatol .*, 95: 693.
- Rybojad, M. , Borradori, L. , Verola,O. , Zeller, J. , Puissant, A. and Morel, P. (1990) : Non-AIDS - associated kaposi's sarcoma (classical and endemic African types) : Treatment with low doses of recombinant interferon- alpha . *J . Invest. Dermatol.*, 95: 176s .
- Sacchini, V., Melloni, E. and Marchesini, R . (1987) : Preliminary clinical studies with PDT by topical TPPS in neoplastic skin lesions. *Lasers. Surg . Med.*, 7:6. Quoted from: Miller, S.J. (1991 II)
- Safai, B., Bason, M. , Friedman-Birnbaum, R. and Nisce, L. (1990) : Interferon in the treatment of AIDS - associated kaposi' s sarcoma : The American experience. *J. Invest. Dermatol .* , 95 : 166 s .
- Samuel, C.E. (1979) : Mechanism of interferon action : Kinetics of interferon action in mouse L 929 cells : Phosphorylation of protein synthesis factor eIF-2 and ribosome-associated P.I. *Virology*, 93 : 2 81 .
- Samuel , C.E. (1988) : Mechanisms of the antiviral action of interferons .*Prog . Nucl. Acid Res. Mol. Biol.* 35 : 27. Quoted from : Baron, S., Tyring,S. K., Fleischmann, W.R.; Copenhagen, D.H., Niesel, D.W., Klimpel, G.R., Stanton, G.J.and Hughes T.K. (1991) .
- Sankowski, A, Janik, P. and Jeziorska, M. (1987) : The results of topical application of 13-cis- - retinoic acid on basal cell carcinoma A correlation of the clinical effect with histopathological examination and serum retinol level. *Neo. Plasma*, 34 : 138. Quoted from : Miller, S. J. (1991 II).
- Sato , N., Nariuchi , H., Tsuruoka, N., Nishihara,T. , Beitz, J.G., Calabresi, P . and Frackelton, A.R. (1990) : Action of TNF and IFN - γ on angiogenesis *in vitro*. *J. Invest. Dermatol.* , 95 : 85 s .

- Schiller, J.H. ,Bushmeyer. S.M. , Ruzicka, F.J. , Princler,G.L., Faltynek, C.R. and Borden, E.C. (1990) : Modulation of interferon receptor expression during combination B_{ser} – interferon and γ – interferon treatment of human colon carcinoma cells. *Cancer Res.* , 50 : 26.
- Schiller, J.H. , Willson, J.K.V., Bittner, G., Wolberg, W.H. , Hawkins, M.J. and Borden , E.C. (1986) : Antiproliferative effects of interferons on human melanoma cells in the human tumor colony forming assay .*J . Interferon Res.*, 6:615.
- Schonfeld, A. , Nitke, S, Schattner, A. , Wallach, D., Crespi, M. , Hahn, T. , Levavi H., , Yarden, O. , Schoham, J. , Doerner, T . and Revel, M. (1984) : Intramuscular human interferon- β injections in treatment of condylomata acuminata. *Lancet* , 12: 1038 .
- Schrieber, R.D., Hicks, L. J., Celada, A., Buchmeier, N. and Gray, P.W.(1985) : Monoclonal antibodies to murine γ – interferon which differentially modulate activation and antiviral activity .*J. Immunol.*, 134 : 1609 .
- Schopf, R.E., Keller, R. , Rehder, M. , Benes, P., Kallinowski, F. and Vaupel , P. (1990) : TNF α primes polymorphonuclear leukocytes for an enhanced respiratory burst to a similar extent as bacterial lipopolysaccharide. *J. Invest. Dermatol.*, 95 : 216s.
- Schulze , H. J. and Mahrle, G. (1986) : Effect of interferon (rIFN-alpha 2, rIFN -gamma) on DNA - synthesis and HLA - DR expression in psoriasis. *Arch. Dermatol. Res.*, 278 : 416 .
- Sehgal, P.B., May, L. T. and Sagar A.D. (1983) : Multiple human β – interferon genes. In : *The Biology of The Interferon System 1983* . De Maeyer, E., and Schellekens, H. (eds). Elsevier Science Publishers, Amsterdam. NewYork . Oxford ., pp.15 .
- Sherwin, S.A., Mayer, D., Ochs, J., Abrams, P.G. , Knost, J.A. , Foon, K .A. , Fein, S. and Oldham, R . K . (1983) : Recombinant leukocyte A interferon in advanced breast cancer . *Ann. Int. Med.* , 98 : 598 .
- Shupack, J. , Stiller, M. , Knobler, E. , Ackerman, C. , Jondreau, L . and Kenny, G . (1990) : Topical alpha - interferon in recurrent genital herpes simplex infection. *Dermatologica*, 181: 134.

- Sidkey, Y.A. and Borden , E.C. (1987) : Inhibition of angiogenesis by interferons : Effects on tumor and lymphocyte - induced vascular responses. *Cancer Res.*, .47:5155.
- Silva, P.D., Micha, J.P. and Silva, D.G. (1985) : Management of condyloma acuminatum. *J . Am. Acad. Dermatol.*, 13: 457 .
- Silver, H.K.B. (1986) : Preface. In : *Interferons In Cancer Treatment*. Silver, H.K.B. (ed). Medical Education Services, Ontario, Canada. p. ix .
- Silver , H.K.B., Salinas, F.A. and Kong, S. (1986) : Immune effects: Role of interferons in immune regulation.. In : *Interferons In Cancer Treatment*. Silver , H.K.B.(ed). Medical Education Services, Ontario, Conada., p.1.
- Silverman, R.H., Krause, D., Jacobsen, H. , Leisy, S. A. , Barlow, D.P. and Friedman, R.M.(1983) : 2-5A- dependent RNase levels vary with interferon treatment , growth rate and cell differentiation . In :*The Biology of The Interferon System 1983* . De Maeyer, E. and Schellekens, H. (eds) . Elsevier Science Publishers. Amsterdam. NewYork. Oxford., p.189 .
- Smoller, B.R. and Krueger,J. (1991) : Detection of cytokine - induced protein γ -immune protein-10 (γ -IP10) in atypical melanocytic proliferations. *J. Am . Acad. Dermatol.*, 25 (4) : 627 .
- Spiegel .R.J. (1986) : Dosage and toxicity, Alpha interferon dosage, toxicity and antibody formation. In : *Interferons In Cancer Treatment* . Silver, H.K.B. (ed) . Medical Education Services. Ontario, Canada., p. 17 .
- Spiegel , R . J . (1989) : Alpha interferons : A clinical overview. *Urology* .34:75s .
- Spiegel, R.J., Jacobs, S.L. and Treuhaft, M.W. (1989) : Anti-interferon Antibodies to interferon - α 2b: Results of compartive assays and clinical perspective . *J.Interferon Res.*, 9 : 17s.
- Spiegel, R.J., Spicehandler, J.R. , Jacobs , S.L. and Oden, E.M. (1986) : Low incidence of serum neutralizing factors in patients receiving recombinant alfa- 2b interferon, (Intron A) . *Am . J. Med.*, 80 : 223 .

- Stadler,R., Bratzke, B. and Baumann, I. (1987) : Morbus Behcet and exogenous interferon. *Hautarzt*, 38 : 97 . Quoted from : Stadler, R. , Mayer - da- Silva, A.M., Bratzke, B. , Garbe, C. and Orfanos, C.(1989) .
- Stadler, R. , Bratzke, B., Schaart, F . and Orfanos, C.E. (1990) : Long- term combined rIFN alpha - 2a and zidovudine therapy for HIV- associated kaposi's sarcoma : Clinical consequences and side effects . *J. Invest. Dermatol.* , 95 :170s.
- Stadler, R. , Mayer - da- Silva, A.M. , Bratzke, B. , Garbe, C. and Orfanos, C.(1989): Interferons in dermatology. *J. Am. Acad. Dermatol.*, 20 (4) : 650 .
- Stadler,R.,Müller, R. and Orfanos, C.E. (1986) : Effect of recombinant alpha - A – interferon on DNA synthesis and differentiation of human keratinocytes *in vitro* *Br.J.Dermatol.* ,114: 273.
- Strander, H. and Cantell , K. (1966) : Production of interferon by human leukocytes *in vitro* . *Ann. Med. Exp. Fenn.* , 44: 265. Quoted from: Trofatter, K. F. (1991) .
- Suhge d' Aubermont,T. and Bennett, R. G.(1984): Failure of curettage and electro-desiccation for removal of basal cell carcinoma . *Arch Dermatol.*, 120 : 1456 .
- Sundmacher, R., Cantell, K., Skoda, R., Hallermann, C. and Neumann- Haeflin, D. (1978) : Human leukocyte and fibroblast interferon in a combination therapy of dendritic keratitis. *Arch. Ophthalmol.*, 208 : 229.
- Symington, F.W. (1989) : Lymphotoxin, tumor necrosis factor and gamma interferon are cytostatic for normal human keratinocytes. *J Invest. Dermatol.*, 92: 798 .
- Taniguchi, T. (1989) : Regulation of interferon - β gene : Structure and function of *cis* - elements and *trans*- acting factors . *J.Interferon Res.* , 9 : 633.
- Taniguchi, T., Ohno,S., Fujii - Kuriyama, Y. and Muramatsu, M. (1980) : The nucleotide sequence of human fibroblast interferon cDNA. *Gene*, 20 : 11 . Quoted from : Goldstein, D. and Laszlo, J. (1986) .
- Tank, B., Habets, W., Naafs, B., Stolz, E. and Van Joost, T. (1989) : Intralesional treatment of basal cell carcinoma with low - dose recombinant interferon gamma . *J. Am. Acad. Dermatol.* , 21(4) ; 734 .

- Taylor, R.S. , Griffiths C.E.M. and Brown, M.D. (1989) : Constitutive absence and cytokine - induced expression of adhesion molecules in basal cell carcinoma. (Abstr.). *J. Invest . Dermatol.*, 92: 530 A .
- Taylor, R.S. , Griffiths, C.E.M. , Brown, M.D. , Swanson, N.A. and Nickoloff, B.J.(1990) : Gonitutive absence and interferon- γ induced expression of adhesion molecules in basal cell carcinoma. *J. Am . Acad. Dermatol.* , 22 (5) : 721.
- Teichmann, J.V., Sieber, G., Ludwig, W.D. and Ruehl, H. (1988) : Modulation of immune function by long - term treatment with recombinant interferon- α 2 in a patient with hairy - cell leukemia . *J. Interferon Res.*, 8:15 - 24 .
- Thestrup - Pedersen, K., Hammer, R. and Kaltoft,K. (1988) : Treatment of mycosis fungoides with recombinant interferon alpha 2a alone and in combination with etretinate. *Br. J. Dermatol* ., 118 : 811
- Thestrup - Pedersen, K. , Jacobsen, I.E., and Frentz, G. (1990) : Intralesional interferon - α 2b treatment of basal cell carcinoma . *Acta. Dermatol . Venereol .* , 70 : 512 .
- Thivolet, J.,Nicolas. J.F., Kanitakis, J., Lyonnet, S.L . and Chouvet, B. (1990) : Recombinant interferon α 2a is effective in the treatment of discoid and subacute cutaneous lupus erythematosus. *Br. J. Dermatol.* 122:405
- Tominaga, S.I., Tominaga, K. and Lengyel, P. (1985) : Characteristics of 30-, 63 - and 89 - Kilodalton proteins whose secretion from mouse fibroblasts in altered by beta - interferon. *J.Biol. Chem.*, 260 : 16406 .
- Tovey, M.G. (1988) : The expression of cytokines in the organs of normal individuals : Role in homeostasis . A review . *J.Biol . Reg. Homeost. Agent.* , 2 : 87. Quoted from : Gresser, I (1990) .
- Treuner, J., Niethammer, D., Dannecker, G., Hagman, R., Neef, V. and Hofschneider, P.H. (1980) : Successful treatment of naso-pharyngeal carcinoma. with interferon. *Lancet* , 1:817 .

- Trofatter, K.F. (1991) : Interferon treatment of anogenital human papillomavirus- related diseases . *Dermatol . Clin.*, 9 (2) : 343 .
- Tura, S., Mazza,P. and Ainzani, P.L. (1987) : Alpha recombinant interferon in the treatment of mycosis fungoides. *Hematologica*, 72 : 337 .
- Tyring, S.K. (1987) : Antitumor actions of interferon : direct, indirect, and synergy with other treatment modalities . *Int . J. Dermatol.*, 26 : 549
- Tyrrell,D.A.J.(1976) : The discovery of interferon. In : *Interferon And Its Clinical Potential* . Tyrrell, D.A.J . (ed). William Heinemann Medical Books Ltd., London, p.1.
- Vance, J.C. , Bart, B.J., Hansen, R.C. , Reichman, R.C. , McEwen, C. , Hatch, K.D. , Berman,B. and Tanner, D.J. (1986) : Intralesional recombinant Alpha - 2 interferon for the treatment of patients with condyloma acuminatum or verruca plantaris. *Arch . Dermatol .*, 122 : 272 .
- Vance, J.C. and Davis, D. (1990) : Interferon alpha 2-b injections used as an adjuvant therapy to carbon dioxide laser vaporization of recalcitrant ano - genital condylomata acuminata. *J. Invest . Dermatol .*, 95 : 146s .
- Van Damme, J., De Ley, M., Van Snick, J., Dinarello,C . and Billiau, A. (1987) : The role of interferon - β_1 and the 26 kDa protein (interferon - β_2) as mediators of the antiviral effect of interleukin 1 and tumor necrosis factor. *J Immunol.*, 139 :1867
- Van Heuvel , M., Bosveld, I.J., Klaassen, P., Zwarthoff, E.C. and Trapman, J. (1988): Structure-function analysis of murine interferon - α : antiviral properties of novel hybrid interferons . *J.Interferon Res.*, 8 : 5 .
- Veomett, G.E. (1988) : Effect of interferon on secretion of proteins by various murine cell lines *J. Interferon Res .* , 8 : 217.
- Viac , J ., Bustamante , R. and Thivolet, J. (1977) : Characterization of mononuclear cells in the inflammatory infiltrates of cutaneous tumours . *Br.J. Dermatol .*, 97: 1
- Visco, G. , Boumis, E. , Noto, P. and Comandini, U.V. (1991) : Prevention of side - effects of interferon . *Lancet*, 337 : 741 .

- Vogel, S.N., Finbloom, D.S., English, K.E., Rosensterich, D.L., and Langrath, S.G. (1983) : Interferon - induced enhancement of macrophage Fc receptor expression : β - interferon treatment of C3H / He J macrophages results in increased numbers and density of Fc receptors *J. Immunol.*, 130 : 1210 .
- Vogetseder , W. , Feichtinger, H. and .Schulz, T.F. (1989) : Expression of 7F7 - antigen, a human adhesion molecule identical to intercellular adhesion molecule - 1 (ICAM - 1) in human carcinomas and their stromal fibroblasts, *Int. J. Cancer* 43 : 768 .
- Vonderheid , E.C . , Thompson , R., Smiles , K.A . and Lattanand, A . (1987) : Recombinant interferon alpha - 2b in plaque - phase mycosis fungoides . *Arch. Dermatol.* , 123 : 757 .
- Von Wussow , P., Freund, M. and Block B . (1987) : Clinical significance of anti - interferon - α antibody titres during interferon therapy . *Lancet* , ii: 635 .
- Von Wussow, P., Jakschies, D. , Freund , M. and Deicher, H. (1989) : Humoral response to recombinant interferon- α 2b in patients receiving recombinant interferon - α 2b therapy . *J. Interferon Res.* , 9 : 25s. .
- Wagner, W.H. (1983) : Interferons. *Pharm. Ind.* , 45 (11) : 1141 . Quoted from : Stadler R., Mayer -da- Silva . A., Bratzke, B., Garbe,C. and Orfanos,C. (1989) :
- Wallberg, P . and Skog, E. (1991) : The incidence of basal cell carcinoma in an area of Stockholm county during the period 1971 - 1980 . *Acta . Dermatol. Venereol.*, 71 : 134 .
- Weck, P.K. , Leventhal, B. G. , Brand, C. and Finter , N. B . (1989) : Detection and incidence of neutralizing antibodies to interferon- α - nl *J. Interferon Res.* , 9: 37s.
- Weil., J., Epstein . C.J., Epstein, L.B., Sedmak , J.J. , Sabran, J.L. and Grossberg, S.E. (1983) : A unique set of polypeptides is induced by gamma interferon in addition to those induced in common with alpha and beta interferons *Nature*, 301: 437.
- Weinstein, G.D. and Frost,P . (1970) : Cell proliferation in human basal cell carcinoma. *Cancer Res.* , 30 : 724. Quoted from : Miller,S.J. (1991 I)

- Weltfriend, S. , Friedman -Birnbaum, R. and Pollack, S. (1990) : Decreased natural killer cell function in patients with classical kaposi's sarcoma. *Dermatologica* , 181: 207.
- Wetzel , R., Johnston, P.D. and Czarniecki, C.W. (1983) : Roles of the disulfide bonds in a human alpha interferon. In : *The Biology of The Interferon System 1983* . De Maeyer, E. and schellekens, H. (eds). Elsevier Science Publishers., Amsterdam. NewYork. Oxford . p. 101 .
- Wheelock, E.F. (1965) : Interferon like virus-inhibitor induced in human leukocytes by phytohemagglutinin. *Science* , 149 :310.
- White, C.W., Sondheimer, H .M. , Crouch, E.C. , Wilaon, H. and Fan , L.L . (1989) : Treatment of pulmonary hemangiomatosis with recombinant interferon alpha 2a . *N . Engl. J. Med.* , 320 (18) : 1197 .
- Wickramasinghe, L. , Hindson, T.C. and Wacks, H. (1989) : Treatment of neoplastic skin lesions with intralesional interferon. *J. Am . Acad . Dermatol.*, 20 (1) : 71.
- Wilder, R.L. (1988) : Treatment of the patient with rheumatoid arthritis refractory to standard therapy. *J. Am. Acad. Dermatol.*, 259 : 2446.
- Winston, D.J., Eron, L.J. and Ho, M. (1988) : Recombinant interferon alpha -2a for treatment of herpes zoster in immunosuppressed patients with cancer. *Am. J. Med*, 85 : 147.
- Wolff, J.M., Zitelli, J.A . , Rabin, B.S. , Smiles , K.A. and Abell, E. (1985) : Intralesional interferon in the treatment of early mycosis fungoides. *J.Am . Acad . Dermatol .* , 13 (4) : 604 .
- Yancey , K. B. and Smith, J.G.(1980) : Interferon : Status in treatment of skin disease . *J. Am.Acad Dermatol .* 3: 585 .
- Yarbov, A. (1982) : Anticancer agent having multifarious activity . *Prog. Clin. Cancer* 9:99. Quoted from : Ringenberg Q.S. and Anderson. P.C. (1986) .
- Yarchoan , R., Mitsuya, H., Myers, C.E. and Broder,S. (1989) : Clinical pharmacology of 3'-azido - 2' . 3' - dideoxythymidine (zidovudine) and related dideoxynucleosides *N. Engl. J. Med.*, 321: 726 .

- Yonehara, S., Yonehara - Takahashi, M., Ishii, A. and Nagata, S. (1983) : Different binding of human interferon α 1 and α 2 to common receptors on human and bovine cells. Studies with recombinant interferons produced in *Escherichia coli*. *J. Biol. Chem.*, 258 : 9046.
- Zacarian, S.A. (1970) : The cryogenic approach to treatment of lid tumors. *Ann. . Ophthalmol.* , 2 : 706.
- Zachariae, H. and Thestrup- Pedersen , K. (1990) : Interferon Alpha and etretinate combination treatment of cutaneous T - cell lymphoma . *J. Invest. Dermatol.* , 95 : 206s .
- Ziegler-Heitbrock, H.W.L and Thiel, E. (1990) : Recombinant IFN- α in lymphomas. *J. Invest. Dermatol.* , 95 : 213s.
- Zouboulis, C.C., Schröder, K. , Garbe , C. , Krasagakis, K. , Krüger, S . and Orfanos, C.E. (1990) : Cytostatic and cytotoxic effects of recombinant tumor necrosis factor - α on sensitive human melanoma cells in *vitro* may result in selection of cells with enhanced markers of malignancy . *J. Invest. Dermatol.* , 95 : 223s .