

Chapter VIII.

REFERENCES

VIII. REFERENCES

- Ahsan, S.N. (1966). Cyclical changes in the testicular activity of the lake chub, *Couesius plumbeus* (Agassiz). Can. J. Zool. 44: 149-160.
- Al-Elyani, R.A. (1983). Biological spawning of family Siganidae *Siganus rivulatus* Forsk. in the Red Sea. M.Sc. Thesis, Girls College, Jeddah, 141 p.
- Al-Hussaini, A.H. (1949). On the functional morphology of the alimentary tract of some fish in relation to differences in their feeding habits: anatomy and histology. Quart. J. Microscop. Sci. 90 (2): 109-141.
- Alexeeva, E.V. (1964). Maturation and resorption of ova in *Vimba vimba natio* Corinata (Pallas). Vopr. Iktyol, 4, 2, (31): 304-314.
- Allam, S.M. (1976). Biological studies of the fish stocks belonging to genus *Trachurus* *rafinesque* in the Egyptian Mediterranean waters. M.Sc. Thesis, Faculty of Science, Alexandria Univ., 208 p.
- Allam, S.M. (1989). Revision of order Hypotremata along the Mediterranean coast off Alexandria with special reference to family Dasyatidae. Ph.D. Thesis, Faculty of Science, Alexandria Univ. 284 p.

- Allen, K.R. (1935). The food and migration of the perch (*Perca fluviatilis*) in Windermere. J. Anim. Ecol. 4: 264-273.
- Allen, K.R. (1967). Computer programs available at St. Andrews biological station. F. Res. Bd. Can. Tech. Rept. (20) 32 p. and Appendix.
- Amin, E.M. (1985 a). Seasonal developmental changes in the ovaries of *Siganus rivulatus* from the Red Sea. Bull. Ins. Oceanogr. & Fish. ARE, 11: 131-147.
- Amin, E.M. (1985 b). Reproductive cycle of male *Siganus rivulatus* Forsk. with indication to gonadosomatic and hepatosomatic indices. Bull. Ins. Oceanogr. & Fish. ARE, 11: 149-164.
- Angelesen, V., Gneri, F.S. and Nani, A. (1958). La Mer luza del mar Argentina (biologia y taxonomia). Argent. Secr. Mar. Serv. Hidrog. Naval Public Health 1004, 224 p.
- Anokhina, L.E. (1963). Some aspects of the fecundity of the herring in the white sea. Râpp. P.V. Réun Cons. Permivint. Explor. Mer. 154: 123-127.
- Appelget, J. and Smith, L.L. (1951). The determination of age and rate of growth from vertebrae of the channel catfish, *Ictalurus lacustris punctatus*. Trans. Am. Fish. Soc. 80: 119-139.

- Apte, D. and Rao, M. (1992). Electrophoretic studies on lens and muscle proteins of some flat fishes of Bombay coast. *Fishery Technology* 29 (2): 147-150.
- Avtalion, R.R., Duczminer, M., Wojdani, A. and Pruginin, Y. (1976). Determination of allogenic and xenogenic markers in the genus *Tilapia*. II. Identification of *T. aurea*, *T. vulcania* and *T. nilotica* by electrophoretic analysis of their serum proteins. *Aquaculture*, 7: 255-265.
- Bagenal, T.B. (1966). The ecological and geographical aspects of the fecundity of Plaice. *J. Mar. Biol. Ass., U.K.*, 46: 161-186.
- Bagenal, T.B. (1967). A short review of fish fecundity. In: *The biological basis of freshwater fish production* (Ed. Gerking, S.D.) Pp 29-111, Blackwell, Oxford.
- Bagenal, T.B. and Braum, E. (1968). Eggs and early life history. In: *IBP Handbook No.3. Methods of assessment of fish production in freshwater* (Ed. Ricker, W.E.) p.159-181. Blackwell, Scientific, Oxford and Edinburg
- Bagenal, T.B. (1971). The inter-relation of the size of fish eggs, the date of spawning and the production cycle. *J. Fish. Biol.*, 3: 207-219.

- Bagenal, T.B. (1978). Aspects of fish fecundity. In: methods of assessment of ecology of fresh water fish production. p.75-101. (Ed. Gerking, S.D. Gerking) Blackwell, Oxford.
- Bailey, K. (1954). Structure proteins. II. Muscle. In: "The protein" (Eds. Neurath, H. and Bailey, K.) 2, p.951-1055. Academic Press, New York.
- Basaglia, F. (1989). The study of the soluble lens proteins from fifteen Sparidae species. Comparative Biochem. & Physiol. B-Comparative Biochemistry, 94 (4): 625-630.
- Basaglia, F. and Marchetti, M.G. (1990). A comparative study of some soluble proteins of the genus *Diplodus* (Sparidae L. Teleostei). Comparative Biochem. & Physiol. B-Comparative Biochemistry, 95 (4): 653-656.
- Basaglia, F. and Marchetti, A.G. (1991). Study of the soluble white muscle tissue proteins from fifteen sparidae species. J. Fish. Biol., 38: 763 - 772.
- Batts, B.S. (1972). Sexual maturity, fecundity and sex ratio of the Skipjack tuna, *Katsuwonus pelamis* (Linnaeus) in north Carolina waters. Trans. Amer. Fish. Soc., 101 (4): 626-637.

- Beckman, W.M.C. (1942). Length-weight relationship, age, sex-ratio and food habits of the smelt (*Osmerus mordax*) from Crystal Lake Benzie County, Michigan Copeia (2): 120-124.
- Beckman, W.M.C. (1945). The length-weight relationship, factors for conversions between standard and total lengths, and coefficients of condition for seven Michigan fishes. Trans. Amer. Fish. Soc. 75: 237-256.
- Ben-Tuvia, A. (1964). Two Siganid fishes of Red Sea origin in the eastern Mediterranean. Bull. Sea. Fish. Res. Sta. Haifa, 37: 3-9.
- Ben-Tuvia, A. (1966). Red Sea fishes recently found in the Mediterranean. Copeia (2): 254-275.
- Ben-Tuvia, A. (1972). Immigration of fishes through the Suez Canal. Proc. Inter. Congr. of Zool. ILOR, Haifa, 7 p.
- Ben-Tuvia, A. (1985). The impact of the Lessepsin (Suez Canal) fish migration on the Eastern Mediterranean ecosystem. In: Mediterranean Marine Ecosystem. p. 367-375 (Eds. Moraitou, M. et al.) Nato Conference Series I: Ecology (8) Plenum Press, New York.
- Ben-Tuvia, A. (1986). Siganide. In: Fish of the North-eastern Atlantic and the Mediterranean. II 1473 p. Unesco, Paris.

- Ben-Tuvia, A., Kissil, G.W. and Popper, D. (1973). Experiments in rearing rabbit fish *Siganus rivulatus* in sea water. *Aquaculture* 1: 359-364.
- Berg, J. (1979). Discussion of methods of investigating the food of fishes with reference to a preliminary study of the prey of *Gobiusculus flavescens*. *Mar. Biol.* 50: 263-273.
- Berhaut, J.A. (1973). Biologie des stades Juvenile de Teleosteens Mugilidae, *Mugil auratus* Risso 1810, *Mugil capito* Curvier 1829 et *Mugil saliens* Risso 1810. *Aquaculture*, 2: 251-266.
- Berry, F.H.; Lee, D.W. and Bertolino, A.R. (1977). Progress in Atlantic blue fin tuna aging attempts. In: *Internat. Comm. for the Conserv. of Atlantic Tunas. Collect. Vol. of Scienti. Pap.*, 6 (67): 305-319.
- Beverton, R.J. and Holt, S.J. (1957). On the dynamics of exploited fish populations. *Fish. Invest. Lond. Ser.2* (19): 533 p.
- Bhattacharya, C.G. (1966). Fitting a class of growth curves. *Sankhya, Ser.* 13 (28) 1 & 2: 1-10.
- Blackburn, M. (1960). Study of condition (weight or length) of Australian Baracounts, *Thyrsites atus* (Euphresen). *Aust. J. Mar. Fresh Res.* 11 (1): 14-41.
-

- Blake, B.F. (1977) Food and feeding of the mummyrid fishes of Lake Kainji, with special reference to seasonal variation and interspecific difference. J. Fish. Biol. II: 315-328.
- Blegvad, H. (1917) On the food of fish in the Danish waters within the skaw. Rep. Dan Biol. Stn. 24: 17-72.
- Bohlert, G.W. (1985) Using objective criteria and multiple regression models for age determination in fishes. Fish. Bull. 83: 103-117.
- Boothby, R.N. and James, W.A. Vault, Jr. (1971) Food habits, length-weight relationship, and condition factor of the Red Drum (*Sciaenops ocellata*) in southeastern Louisiana. Trans. Amer. Fish. Soc. 100 (2): 290-295.
- Brown, M.E. (1957) The physiology of fishes. Vol.I. Academic Press Inc., New York, 447 p.
- Bryan, P.G. (1975) Food habits, functional digestive morphology, and assimilation efficiency of rabbit fish *Siganus spinus* (Pisces, Siganidae) on Guam. Pac. Sci., 29 (3): 269-277.
- Buchannan-Wallaston, H.J. and Hodgson, W.C. (1929): A new method of treating frequency curves in fishery statistics, with some results. J. Cons. Perm. Int. Explor. Mer. 4: 207-225.
- Bulkly, R.V. (1970). Changes in yellow bass reproduction associated with environmental conditions. Iowa State.J. Sci., 45 (2): 137-180.
-

- Burkett, R.D. and Jackson, W.B. (1971). The eye lens as an indicator in fresh water drum. *Am. Midl. Nat.* 85: 22-25.
- Carlton, W.G. and Jackson, W.B. (1968). The eye lens as an age indicator in carp. *Copeia* 1968: 633-636.
- Cassie, R.M. (1954). Some uses of probability papers in the analyses of size frequency distribution. *Aust. J. Mar. Fresh Nat. Res.* 5: 513-522.
- Chadwick, E.M.P. (1976). A comparison of growth and abundance for tidal pool fishes in California and British Columbia. *J. Fish Biol.* 8 (1): 27-34.
- Chitravadivelu, K. and Sivapalan, A. (1984). Food and feeding of *Siganus lineatus* from waters around Northern Srilanka. *J. Natn. Sci. Coun. Srilanka*, 12 (1): 129-139.
- Chugunova, N.I. (1963). Age and growth studies in fish. English Text, National Science Foundation, Washington, 132 p.
- Clark, F.N. (1928). The weight-length relationship of California sardine (*Sardina caerulea*) at San Pedro. *Calif. Fish. & Game Bull.*, 12: 58 p.
- Clark, F.N. (1934). Maturity of the California sardine (*Sardina caerulea*) determination by ova diameter measurements. *Calif. Fish. & Game Bull.* 42: 1-11.
-

- Concon, J.M. (1975). Rapid and simple method for the determination of tryptophan in cereal grains. *Analytical Biochem.* 67: 206-219.
- Connell, J.J. (1953). Studies on the proteins of fish skeletal muscle. 2- Electrophoretic analysis of low ionic strength extracts of several species of fish. *Biochem. J.* 55: 378-88.
- Connell, J.J. and Howgate, P.F. (1959). The amino acid composition of some British food fishes. *J. Sci. Food Agr.*, 10: 241-244.
- Cowey, C.B. (1965). Amino acids and related substances in fish. In: "Studies in comparative biochemistry" (Ed. Munday, K.A.) p.41-61, Pergamon Press, Oxford.
- Cowie, W.P. (1968). Identification of fish species by thin-slab polyacrylamide gel electrophoresis of the muscle myogens. *J. Sci. Food Agric.*, 19: 226-228.
- Crivelli, A. (1980). The eye lens weight and age in the common carp, *Cyprinus carpio* L. *J. Fish Biol.* 16: 469-473.
- Crozier, W.J. and Hecht, S. (1914). Correction of weight, length and other body measurements in the weak fish *Cynoscion regalis*. *Bull. U.S. Bur. Fish.*, 33: 139-147.
- D'Ancona, M.V. (1936). La croissance chez les animaux de la mediterrance. *Râpp. Comm. Int. Mer Medit*, 10: 163-224.
-

- Dannevig, E.H. (1956). Chemical composition of the zones in Cod otoliths. *J. Cons. Int. Explor. Mer.*, 21,(2): 156-159.
- Das, S.M. and Moitra, S.K. (1955). Studies on the food of some common fishes of Uttar Pradesh, India. 1. The surface feeders, the mid-feeders and the bottom-feeders. *Proc. Nat. Acad. Sci. India*, 25 B (1 & 2): 1-6.
- Dawson, C.E. (1965). Length-weight relationships of some Gulf of Mexico fishes. *Trans. Amer. Fish. Soc.* 94 (3): 279-280.
- Day, F. (1878). *Fishes of India*. Bernard Quaritch, London, 202 p.
- De Bont, A.F. (1967). Some aspects of age and growth of fish in temperate and tropical waters. *The biological basis of fresh water fish production shel* (Ed. Gerking, D.) 495 p. Blackwell, Oxford.
- Douglas, R.H. (1987). Ocular lens diameter as an indicator of age in brown trout, *Salmo trutta*. *J. Fish. Biol.*, 31 (6): 835-836.
- Drew, A.W. (1971). Preliminary report on Klsebuul and Meyas, two fish of Palau Islands, unpublished report, January, 1971, 20 p.
- Drysdale, J.W. (1975). Isoelectric focusing in polyacrylamide gel. In: *Methods of protein separation, I*. Plenum Publ. Co. New York.

- Dunn, M.S. (1947) Amino acids in food and analytical methods for their determination. Food Technol., 1: 269-286.
- Durant. M. and Cerletti. P. (1979) Amino acid composition of seed proteins of *Lupinus albus*. J. Agric. Food Chem., Vol. 27 No 5: 977-1190.
- El-Deeb, S.I., El-Serafy S.S, Zawail M.E.M. and Saied H.E. (1995) Genetic differntiation of sarcoplasmic protein in family Mugillidae at different habitats. Bull. Nat. Inst. Of Oceanogr. & Fish., ARE 21 (I): 263 - 277.
- El-Gammal, F.I. (1988) Age, growth and mortality of rabbit fish, *Siganus rivulatus* (Forsk, 1775) from the Red Sea. Bull. Inst. Oceanogr. & Fish., ARE, 14: 13-21.
- El-Sayed, A.M. (1979) Seasonal biochemical changes in (*Pelamys* spp.) from Egyptian Mediterranean waters. M.Sc. Thesis, Faculty of Science, Alexandria Univ., 186 p.
- El-Sayed, A.M. and Bary, K.A. (1994). Life cycle and fecundity of rabbit fish, *Siganus canaliculatus* (Teleostei : Siganidae) in the Arabian Gulf. Oebalia., Vol.XX: 79-88.
- Elserafy, S.S. (1981): Studies on some biological aspects of *Clarias batrachus*. Ph.D. Thesis, Calcutta Univ., 291 p.
- Emlen, J.M. (1966) The role of time and energy in food preference. Am. Nat. 100, 611-617.
-

FAO (1984). Species identification sheets for fishery purposes. Fishing area SW Indian Ocean, FAO, Rome.

- Forbes, S.A. (1883). The food of the smaller fresh water fishes. Bull. III. St. Lab. Nat. Hist., 1, 61-86.

Ford, E. (1933). Herring Investigations. J. Mar. Biol. Ass. UK. 19: 305-384.

- Forskal, P. (1775). Desoz. anim.: X, 25 (Red Sea) Lectotype: Universitetes Zoologiske Museum, Kobenhagen.

- Fowler, H.W. and Bean, B.A. (1929). Fishes of the Philippines and adjacent seas. Bull. US Natl. Mus. US Gov. Printing Office, Washington, 8: 100 p.

Fowler, H.W. and Steinitz, H. (1956). Fishes from Cyprus, Iran, Iraq, Israel and Oman. Bull. Res. Counc. of Israel, 5 B (3-4): 260-292.

Fujita, S. and Ueno, M. (1954). On the development of the egg and prelarval stages of *Siganus fuscescens* (Houttuyn) by artificial insemination. Jap. J. Ichthyol., 3: 129-132.

- Fulton, F. (1902). Rate of growth of sea fishes. Scient. Invest. Fish. Div., Scot. Rep. 20.

- GAFRD (1996). Statistical yearbook. General Authority for Fish Resources Development, Cairo, Egypt, 27 p. (in Arabic).
- Gamal, S.M. (1993). Reproductive biology and induced spawning of one species of family Siganidae. Ph.D. Thesis, Faculty of Science, Suez Canal Univ., 235 p.
- Gamee, F.M. (1996). Biological studies of Family Mugilidae from the vicinity off Alexandria. M.Sc. Thesis, Faculty of Science, Alexandria Univ., 207 p.
- Geiser, S.W. (1924). Sex-ratios and spermatogenesis in the top-minnow, *Gambusia holbrooki*. Ord. Biol. Bull., 47 (3): 175-212.
- George, C.J. (1972). Notes on the breeding and movements of the rabbit fish *Siganus rivulatus* (Forsk.) and *Siganus luridus* (Ruppl) in the coastal water of Lebanon. Ann. Mus. Stor. Nat. Genova, 79: 32-44.
- George, C.J. and Athanassiou, V. (1967). A two years study of the fishes appearing in the Seine fishery of St. George Bay, Lebanon. Ann. Mus. Stor. Nat. Genova, 76: 237-294.
- George, E.L. and Hadley, W.F. (1979). Food and habitat partitioning between rock bass (*Ambloplites rupestris*) and small mouth bass (*Micropterus dolomieu*) young of the year. Trans. Amer. Fish. Soc., 108: 253 - 261.
-

- Ghosh, K.K. (1974). On the choice of a growth curve for Indian Major Carps Von Bertalanffy or Gompertz. J. Ind. Soc. Agr. New Delhi 26 (2): 57-70.
- Golani and Baranes, A. (1990). Adaption of Red Sea fish to the eastern Mediterranean environment. Rapp. Comm. Int. Mer. Medit. 32, 1 (1990).
- Graham, M. (1924). The annual cycle in the life of the mature cod in North Sea. Fish. Invest. Lond. Ser. 2 (6): 1-77.
- Graham, M. (1929). Studies of age determination in fish. Fish. Invest. Lond. Ser. 11 (2): 1-50.
- Graham, M. (1956). Sea fisheries. Their investigation in the United Kingdom. Edward Arnold (Publ.) Ltd. London, 487 p.
- Gulland, H.A. (1969). Manual for fish stock assessments. Part I. Fish population analysis. FAO Mar. Fish. Sci. 154 p.
- Günther, A. (1861). Catalogus of Acanthepterygian fishes in the collection of the British Museum. London, 313-324.
 - Hajort, J. (1910). Report on Herring Investigation untill January 1910. Publ. Circons. Cons. Perm. Int. Explor. Mer.
-

- Hamza, A.K. (1980). Studies on certain aspects of biology and reproduction of genus *Notopterus*. Ph. D. Thesis Submitted to Calcutta Univ. 252 p.
- Hamza, A-K., Ezzat, A.A., Allam, S.M. and Hashem, M.T. (1988 a). Food and feeding habits of Stingray *Dasyatis pastinaca* in the Egyptian Mediterranean waters off Alexandria. Bull. Nat. Inst. Oceanogr. & Fish. ARE, 14 (3): 77-85.
- Hamza, A. K., Ezzat, A. A., Allam, S.M. and Hashem, M. T. (1988 b). Age and growth of stingray *Dasyatis pastinaca* in the Egyptian Mediterranean water off Alexandria. Bull. Nat. Inst. Oceanogr. & Fish., ARE, 14 (3): 227 - 237.
- Hara, S.; Duray, M.N.; Parazo, M. and Taki, Y. (1986) .Year round spawning and seed production of the rabbit fish, *Siganus guttatus*. Aquaculture, 59: 259-272.
- Harding, J. P. (1949). The use of probability paper for the graphical analysis of polymodal frequency distributions. J. mar. Biol. Assoc. UK 28: 141 - 153.
- Hashem, M.T. (1983) . Biological studies on *Siganus rivulatus* (Forsk.) in the Red Sea. J. Fac. Mar. Sci. Jeddah, 3: 119-127.
-

- Hederstorm, Hans (1959). Observations on the age of fishes. Rept. Inst. Freshwater Res. Drottningholm, 40: 161-164. (Original Version Published in 1759).
- Heincke, F. (1905). Das Vorkommen and die Verbreitung der, Eier, derlarven and der verschiedenen. Altersstufen der Nutzfische in der Nordsee. Rapp. P.V. Reun. Cons. Permunt. Explor. Mer., 3: Append E 41 p.
- Herre, A.W. and Montalban, H.R. (1928). The philippine siganids. Philipp. J. Sci. 35 (2): 151 - 185.
- Hiatt, R.W. and Strasburg, D.W. (1960). Ecological relationships of the fish fauna on coral reefs of the Marshall Island Ecol. Monogr, 30: 65 - 127.
- Hickling, C.F. (1930). The natural history of the Hake, III: Seasonal changes in the condition of the hake. Fish. Invest., Ser. 2 (12): 1-78 .
- Hickling, C.F. (1933) . The natural history of the Hake. IV: Age determination and the growth rate. Fish Invest., Ser. 1 (8) 2: 1-84.
- Hile, R. (1936). Age and growth of cisco, *Leucichthys arteda* (Le Suerr) in the lakes of north eastern highlands Wisconsin. Bull. Bur. Fish. U.S. 48 (19): 211-317.

- Hill, W.S., Learson, R.J. and Lane, J.P. (1966). Identification of fish species by Agar Gel electrophoresis. J. Assoc. Offic. Agr. Chemists. (49): 1245.
- Hoar, W.S. (1965). Comparative physiology: hormones and reproduction in fishes. A. Rev. Physiol, 27: 51 - 70.
- Hodder, V.M. (1963). Fecundity of Grand Bank Haddock J. Fish Res. Bd. Can. 20: 1465 - 1487.
- Hoffbauer, C. (1898). Die Altersbestimmung des Karp Fens an Seiner Schuppe. Allgemeine Fischereizeitung, 23 (19).
- Hussein, Kh.A. (1986 a). Timing of Spawing and Fecundity of Mediterranean *Siganus rivulatus* Forsk. Bull. Inst. Oceanogr & Fish. ARE, 12 : 175-186.
- Hussein, Kh.A. (1986 b). The use of minute embeded scales for age determination of the Mediterrenean sea Siganid: *Siganus rivulatus* Forsk. Bull. Inst. Oceanogr & Fish. ARE, 12: 187-198.
- Hynes, H.B.N. (1950) . The food of freshwater stickelbacks *Gasterosteus aculeatus* and *Pygosteus pungitius*) with a review of methods used in studies of the food of fishes. J. Anim. Ecol, 19: 36-58.
- Hyslop, E.J. (1980). Stomach content analysis - a review of methods and their application. J. Fish. Biol. 17, 411 - 429.

- Ivlev, V.S. (1961). Experimental ecology of feeding of fishes. Yale Univ. Press, New Haven, 302 p.
- Jamison, A. and Teixeira, A.S. (1991). Eye lens proteins as genetic markers in anguillids. *Animal Genetics* 22 (1): 45 - 52.
- Jayasankar, P. (1990). Some aspects of the white spotted spine-foot *Siganus canaliculatus* (Park, 1797) from the Gulf of Manner. *Indian J. Fish.* 37 (1): 9-14.
- Jhingran, A.G. (1966). The Biology and fisheries of the Clupeoids, *Setipinna phasa* (Hamilton) and *Gudusia chapra* (Hamilton) in the Ganga river system. Thesis, Ph.D., Allahabad Univ. 1966, 335 p.
- Johnson, J.L., Hepfer, C.F. and Turchi, S.L. (1989). Quantitative analysis of eye lens proteins in sunfish. *J. of the Pennsylvania Acad. of Scien.* 462 (3): 138 - 141.
- Jones, N.R. (1959). The free amino acids of fish. II. Fresh skeletal muscle from lemon sole (*Pleuronectes microcephalus*). *J. Sci. Fel. Agric.*, 10: 282 - 286.
- Jones, R.S. (1968). Preliminary study of the food habits of the rabbit fish *Siganus spinus* (Siganidae) with notes on the use of this fish in pisciculture. Univ. Guam Mar. Lab. Lab. 8 p., Mimographed.

- Kamal, M. Yusuf. (1971). Length-weight relation of *Cirrhina mirigal* (Hamilton) from commercial catch at Allahabad. Proc. Natn. Inst. Sci. India (B), 41 (4): 419 - 422.
- Kamel, S.A. (1987) -Identification of some fish species from the Egyptian Mediterranean waters by isoelectric focusing. Proc. Zool. Soc. Egypt. 14: 169 - 177.
- Kamel, S. A. (1989) -Identification of *Oreochromis niloticus* in Egypt by thin layer polyacrylamide isoelectric focusing. Bull. Nat. Inst. Oceanogr. & Fish. ARE, 15 (2): 31 - 37.
- Karagitson, E., Stergiou, K. and Papaconstantinou, C. (1986). Preliminary study on the diet of *Pagrus pagrus*, *phycis phycis* and *Siganus rivulatus* in Kastellorize waters. Eastern Mediterranean Greece. FAO. Fisheries Report No. 361.
- Kesteven, G. L. (1960). Manual of field methods in fisheries biology. FAO. Man. Fish. Sci. No. 1 : 1 - 152.
- Klunzinger, C.B. (1884). Die Fische des Rothen Meeres. E. Schweizerbartische verlagshandlung. Stuttgart, 133 p.
- Kohno, H., Hara, S., Duray, M. and Gallego, A. (1988). Transition from endogenous to exogenous nutrition sources in larval rabbit fish *Siganus guttatus*. Bull. Jap. Soc. Sci. Fish, 54 (7): 1083-1091.

- Konosu, S., S. Katori; R. Ota, S. Eguchi and T. Mori (1956). Amino acid composition of fish muscle protein. Bull. Jap. Soc. Sci. Fish, 21 (11): 1163 - 1166. (in Japanese with English Summary).
- Ktari, F. and Bouhlal, M. (1971). Capture de *Siganus luridus* (Ruppel) dans Le Galf de Tunis. Bull. Inst. Oceangr. Pêche, Salammbô, 2 (I): 49 - 52.
- Lagler, K.F. (1956). Fresh water fishery biology. WMC Brown Co., Dubuque. Iowa, 421 p.
- Lam, T.J. (1974). Siganids, their biology and mariculture potential. Aquaculture (3): 235 - 254.
- Lane, J.P., Hill, W.S. and Learson, R.J. (1966). Identification of fish species by cellulose acetate electrophoresis. Commer. Fish Rev., 28 (3): 10
- Le Cren, E.D. (1951). The length - weight relationship and seasonal cycle in gonad weight and condition in the Perch *Perca fluviatilis*. J. Am. Ecol., 20 (2): 201 - 219.
- Lee, R.M. (1920). A review of methods of age and growth determination in fishes by means of scales. J. Cons. 4 (2): 1-32.
- Liley, N. and Donaldson, R. (1969). Reproduction, fish physiology. Acad. Press, New York, Part II, 76 - 85.

- Linnaeus, C. (1758). *Systema naturae*, 10th Ed.

Love, R.M. (1970). *The chemical biology of fishes*. Academic Press, London, New York, 547 p.

Lundberg, B. and Lipkin, Y. (1979). Natural food of the herbivorous rabbit fish (*Siganus* spp.) in north Red sea. *Bot. Mar.*, (22): 173 - 181.

Lundberg, B. and Lipkin, Y. (1992). Seasonal grazing site and fish size effects on patterns of local consumption by the herbivorous fish, *Siganus rivulatus* at Mikhmoret (Mediterranean, Israel). *Envir. Qualit. & Ecosyst. Stabil.* 27 (7-8): 413-419.

Lundstorm, R.C. (1981). Fish species identification by isoelectric focusing sarcoplasmic protein polymorphism in monk fish (*Lophius americanus*). *J. Assoc. Anal. Chem.* 64 (1): 32 - 37.

Lundstorm, R.C. (1983). Identification of Pacific Rockfish (*Sebastes*) species by isoelectric focusing. *J. Assoc. Anal. Chem.* 66 (4): 974 - 980.

Man, H.S.H. and Hodgkiss, I.J. (1977). Studies on the ichthyofauna in Plover Cove Reservoir, Hong Kong: feeding and food relations. *J. Fish. Biol.* 11: 1-3.

- Mansour, H.A.M. (1984). Comparative biochemical studies on the protein and amino acids contents in fresh water fishes and marine fishes. M.Sc. Thesis, Faculty of Science, Alexandria Univ. 234 p.
- Mather, F.J., and Skuck, H.A. (1960). Growth of bluefin tuna of the western north Atlantic. Fish Bull. Fish and Wildl. Ser. 61 (179): 39-52.
- Mc Kearn, J.L. and Horton, H.F. (1974). Development of steelhead trout (*Salmo gairdneri*) otoliths and their use for age analysis and for separating summer from winter races and wild from hatchery stocks. J. Fish Res. Bd. Can., 31 (8): 1420-1426.
- McArthur, R.H. and Pinaka, E.R. (1966). On optimal use of a patchy environment. Am. Nat. 100: 603-609.
- Medford, B.A. and Mackay, W.C. (1978). Protein and lipid content of gonads, liver and muscle of Northern Pike (*Esox lucius*) in relation to gonad growth. J. Fish Res. Bd. Can., 35 (2): 213-219.
- Mendies, A.S. (1954). Fishes of Ceylon (A Catalogue, Key and Bibliography). 2. Bull. Fish. Res. Stn. Ceylon.
- Menon, M.D. (1950). The use of bones, other than otoliths, in determining the age and growth rate of fishes. J. Cons. Perm. Int. Explor. Mer., 16 (3): 311-335.
-

- Menzies, W.J.H. (1924). The relation of the weight to length in Scottish salmon. *Salm. Trout. Mag.*, 36: 133-136.
- Mertz, E.T. (1968). Amino acid and protein requirements of fish. In: *Fish Research* (Eds. Neuhaus, O.W. and Halver, J.E.) p.233-244.
- Mohamed, N.I. (1991). Biological and Biochemical studies of some siganid fishes from the Mediterranean waters off Alexandria. M.Sc. Thesis, Faculty of Sciences, Alexandria Univ. 215 p.
- Moore, E. (1947). Studies of the marine resources of New England. VI. The sand flounder *Lophsetta aquasa* (Mitchill) a general study of the species with special emphasis on age determination by means of scales and otoliths. *Bull. Bingham. Oceanogr. Coll* II (3): 1-29.
- Moreau. J., Bambino, G. and Pauly, D. (1986). Indices of overall growth performance of 100 tilapia (Cichlidae) populations, P-201 - 206. In : *Forum. Asian Fisheries Societ.* (Eds. Maclean, J.L. and Fisheries, L.B.) Manila, Philippines.
- Mouneimine, N. (1978)-Poisson elescotes duliban. These de Doct. D'Etat Universite Pierreet Mariecurie, Paris VI, 490 p.
- Natrajan, A.V. and A. G. Jhingran, A.G. (1961)-Index of preponderance A method of grading, the food elements in the stomach analysis of fishes. *Indian. J. Fish.*, 8: 54 - 59.

- Natrajan, A.V. and Jhingran, A.G. (1962). On the biology of *Catla catla* (Ham.) from the River Jamuna. Proc. Nat. Inst. Sci. India, 29 (3): 326-355.
- Ng, C.S., Low, L.K., Lam, C.P. and Yamada, J. (1986). Thin layer polyacrylamid isoelectric focusing as a method for species identification of tropical fishes. Singapore J. Pri. Ind., 14 (1): 36-45.
- Nikolsky, G.V. (1963). The Ecology of fishes. Academic press, London and New York, 352 p.
- Nirenberg, M.W., Matthaei, J.H., Jones, O.W., Martin, R.G. and Barondes, S.H. (1963). Approximation of genetic code via cell - free protein synthesis directed by template RNA. Fed. Proc., 22: 55 - 61.
- Ntiba, M.J. and Jaccarini, V. (1988). Age and growth parameters of *Siganus sutor* in Kenyan marine inshore water derived from numbers of otolith microbands and fish lengths. J. Fish Biol. 33 (3): 465-470.
- Ntiba, M.J. and Jaccarini, V. (1990). Gonad maturation and spawning times of *Siganus sutor* off the Kenya Coast: evidence for definite spawning seas on in a tropical fish. J. Fish. Biol., 37 (2): 315 - 325.
- Ochoa, S. (1963). Synthetic polynucleotides and genetic code. Fed. Proc., 22: 62 - 74.

- O'Donoghue, C.H. and Boyd, E.M. (1930). A preliminary investigation of the food of the sea trout (*Salmo trutta*). Fish Bd. Scotland. Salmon Fish. III: 15 p.
- Okada, Y. (1966). Fishes of Japan. Maruzen, Tokyo, 434 p.
- O' Maoileidight, N., Cawdery, S., Bracken, J.J. and A. Ferguson, A. (1988). Morphometric, meristic character and electrophoretic analysis of two Irish populations of white shad, *Alosa fallax* (Lacepede). J. Fish. Biol., 32: 355 - 366.
- Pal, J. N. and Basu, S. (1991). Isolation, Purification and preliminary studies on fish, eye lens glycoprotein. Indian. J. of Biochem. & Biophys. 28 (1): 63 - 64.
- Paloheimo, J.E. and Dickie, L.M. (1965). Food and growth of fishes. 1- A growth curve derived from experimental data. J. Fish Res. Bd. Can., 22 (2): 521-542.
- Pantulu R. V. (1963). Studies on the age and growth, fecundity and spawning of *Osteogeneiosus militaris* (Linn.). J. Cons. Perm. Int. Explor. Mer 28: 295 - 315.
- Papaconstantinou, C. (1987). Distribution of the Leseptian fish migrants in the Aegean Sea. Biologia Gallo - Hellenica (13): 15 - 20.
-

- Paul, V. J.; Nelson, S. G. and Sanger, H. R. (1990). Feeding preference of adult and Juvenile rabbit fish *Siganus argenteus* in relation to chemical defenses of tropical sea weeds. Mar. Ecol. Prog. Ser. (60): 23 - 34.
- Pauly, D. (1983). Some simple methods for the assessment of tropical fish stocks. FAO. Fish. Tech. Pap. (234): 52 p.
- Payne Jr., W.R. (1963). Protein typing of fish, pork, and beef by disc electrophoresis" J. Assoc. Offic. Agr. Chemist. 46, 1003.
- Pears, A.S. (1915). On the food of the small shore fishes in the waters near Madison, Wisconsin. Bull. Wis. Nat. Hist. Soc. Milwaukee, 13: 1122.
- Peiris, T.S.S. and Grero, J. (1972). Chemical analysis of some Ceylon fishes. Bull. Fish. Res. Stn. Srilanka, 23 (1 & 2): 1-7.
- Perutz, M.F. (1962). Proteins and nucleic acids : Structure and function. Elsevier, New York. 211 p .
- Petersen, C.G.J. (1892). Fiskensbiologiske forholdi Holboek Fjord 1890 - 91. Beretning Fra: den. Dansk. Biologiske. Station for 1890 - 91 (1): 121 - 183.
- Pillay, T.V.R. (1952). A critique of the methods of study of food of fishes. J. Zool. Soc. India, 4: 185-200.

- Pinkas, L., Oliphant, M.S. and Iverson, I.L.K. (1971). Food habits of albacore, blue fin tuna and bonito in California waters. Fish. Bull. Calif. 152: 1-105.
- Polder, J.J. (1961). Cyclical changes in testis and ovary related to maturity stages in the north sea herring *Clupea herengus* L. Archs. Neerl. Zool. 14: 45 - 60.
- Pollard, D.A. (1973). The biology of a land-locked form of the normally catadromous Salmon form fish *Galaxias maculatus* (Jenyns) V. Composition of the diet. Aust. J. Mar. Freshwater Res. 24: 281-295.
- Popper, D. and Gundermann, N. (1975). Some ecological and behavioural aspects of Siganid populations in the Red sea and Mediterranean Coasts of Israel in relation to their suitability for aquaculture. Aquaculture (6): 127 - 141.
- Popper, D.; Gordin, H. and Kissil, G.W. (1973). Fertilization and hatching of the rabbit fish *Siganus rivulatus*. Aquaculture, 2 (1): 37-44.
- Popper, D.; Pitt, R. and Zohar, Y. (1979). Experiments on the propagation of Red Sea Siganids and some notes on their reproduction in nature. Aquaculture, 16: 177-181.
- Por, F.D. (1978). Lessepsian migration. Ecol. Stud. 23, 228 p.

- Powels, P.M. (1965) Life history and ecology of American plaice (*Hippoglossoides platessoides* Fab.) in the Magdalen Shallows. J. Fish Res. Bd. Can., 22 (2): 565-598.
- Qasim, S.Z. (1973). Some implications of the problem of age and growth in marine fishes from Indian waters. Curr. Sci. 20 (2): 351-371.
- Rabaey, M. (1965) Comparative study of tissue proteins (lens and muscle) in fish. In: Proteins of the Biological fluids". 12th Colloquium, 1964 (Ed. Peters, H.). p.547.
- Reay, P.J. (1972) The seasonal pattern of otolith growth and its application to back calculation studies in *Amnodytes tobianus* L. J. Cons. Int. Explor. Mer. 34 (3): 485-504.
- Reibisch, I. (1899) Upwe die Eizahl bei *Pleurinectes platessa* and die Altersbestimmung dieser form aus den otolithen. Wiss Meeresunteres. 4 : 231 - 248.
- Ricker, W.E. (1958) Hand book of computation for biological statistics of fish populations. Bull. Fish. Res. Bd. Can., (119), 300 p.
- Ricker, W.E. (1975) Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Bd. Can., 191-382p.
- Riggs, C.D. (1955) Reproduction of the white bass, *Morone chrysops*. Invest. Indiana lake streams 4 (3): 87 - 110.

- RizKallah, W., Amir, F.I., El-Gammal, F. and Shiekh El-Din, M.Y. (1989). Studies on the digestive system of *Siganus rivulatus* with special reference to its food and feeding habits, Bull. Inst. Oceanogr. & Fish., ARE, 15: 1 - 30.
- Robertson, J.A. (1933). Notes on the optical appearance of winter rings in some clupoid scales. J. Cons. CIEM, 8: 357-370.
- Rounsfall, G.A. and Evarhart, W.H. (1953). Fishery Science: its methods and applications. J. Wiley & Sons, New York, 444 p.
- Ruell, R. (1977). Reproductive cycle and fecundity of white bass in levis and clark lake. Trans. Am. Fish. Soc. 106 (1): 67 - 76.
- Rüppel (1828). Fische Rothen Meeres: 45 (Tor, Red Sea) Lectotype: Senckenberg Museum, Frankfort No. 5153-5154.
- Schaefer, M.B. and Orange, C.J. (1956). Studies on the sexual development and spawning of yellow fin tuna (*Neothunnus macropterus*) and Skipjack (*Katsuwonus pelamis*) in three areas of the eastern Pacific Ocean by examination of gonads. Bull. Inter. Amr. Trop. Tuna Comm., 1 (6): 281-349.
- Sibley, C. G. (1962). The comparative morphology of protein molecules as data for classification. Systematic Zoology, 11: 108 - 118.

- Siezen, R.J. (1989). Eye lens aging in the spiny dogfish (*Squalus acanthias*): 1. Age determination from lens weight. Curr. Eye Res., 8 (7): 707-712.
- Sinha, M.R. (1981). Biology and Fishery of the canine cat fish *Plotosus canius* (Hamilton). Ph.D Thesis, Calcutta Univ. 225 p.
- Smithies, V. (1955). Zone electrophoresis in starch gel group variation in the serum proteins of normal human adults. Bioche. J., 61 : 629 - 641.
- Smith, A.C. (1989). Tracing phylogeny in a fish group (groupers) through proteins of the layers of the eye lens core. Comparative Biochem & physiol. B - Comparative Biochemistry, 93 (3). 523 - 528.
- Snedecor, G.W. (1961). Statistical Methods. Allied. Pacific Pvt. Ltd., Bombay, 534 p.
- Southern, R. (1935). Reports from the Limnological Laboratory. III. The food and growth of brown trout from Lough Derg and River Shannon. Proc. R. Irish Acad. 42 B: 87-172.
- Steinitz, W. (1967). A tentative list of immigrants via the Suez Canal. Israel. J. of Zool., 16: 166-169.

- Steinitz, W. and Ben-Tuvia, A. (1972). Contribution to the knowledge of Suez Canal migration (Fishes of the Suez Canal). Israel J. of Zool., 21: 285-289.
- Stergiou, K.I. (1988). Feeding habits of the Lessepsian migrant *Siganus luridus* in the eastern Mediterranean, its new environment J. Fish. Biol, 33 : 531 - 543.
- Stickney, R. R. (1976). Food habits of Georgia estuary fishes. II *Symphurus plagiusa* (Pluronecti formes: Cynoglossidae). Trans. Am. Fish. Soc. 105: 202 - 207.
- Suyehiro, Y. (1942). A study of the digestive system and feeding habits of fish. Jap. J. Zool. 10 (1), 303 p.
- Suzuki, H. and Phan, V.N. (1990). Electrophoretic study on intraspecific variations and interspecific relation of marine cat fishes (*Siluriformes, Ariidae*) of Cananeria (Sao Paulo, Brazil). 1 - General proteins of eye lens and skeletal muscle. Do Instituto Oceanografica, 38 (1): 31 - 42.
- Swynnerton, G.H. and Worthington, E.B. (1940). Note on the food of fish in Hawes water. J. Anim. Ecol. 9: 183-187.
- Tanaka, S. (1962). A method of analysing a polymodal frequency distribution and its application to the length distribution of the porgy, *Taeniumiforms* (T. and S.). J. Fish. Res. Bd. Can., 19 (6): 1143-1159.
-

- Tech, F.W. (1971). Age and growth in fish production in fresh water. (Ed. Ricker, W.F.) Blackwell, Oxford. 98 - 130 .
- Thompson, R.B. (1959). Food of squawfish *Ptychocheilus oregonensis* (Richardson) of the lower Columbia River. Fish Bull. Fish & Wildl. Serv. 158 (60): 43 - 58.
- Thompson, R.R. (1960). Species identification by starch gel zone electrophoresis of protein extracts 1. Fish. J. A. Assoc. offic. Agr. Chemists (43): 763 - 764.
- Tillier, J.B. (1902). Le canal de Suez et sa faune ichthyologique. Mem. Soc. Zool. Fr., 14: 279-318.
 - Tjurin, P.V. (1927). About the relation between the length of fish and its weight. Rep. Ichthyol. Lab. Siberia, 2 (3): 3-21.
- Tobias, W. J. (1976). Ecology of *Siganus argenteus* (Pisces : Siganidae) in relation to its mariculture potential in Guam. Univ. Guam Mar. Lab. Tech. Rep., 29: 58 - 93.
- Tortonese, E. (1972). Facts and perspectives related to spreading of red Sea organisms into the Mediterranean. Proc. Int. Congr of Zool 7th No. (3). Haifa, 7 p.
- Tsuda, K. T. and Bryan, P. G. (1973). Food preference of Juvenile *Siganus rostratus* and *Siganus spinus* in Guam. Copeia (3) : 604 - 606.
-

- Tsuyuki, H. and Roberts, E. (1961). Muscle proteins of Pacific Salmon (*Oncorhynchus*) I. A note on the separation of muscle proteins soluble in low ionic strength salt solutions. J. Fish Res. Bd. Can., 18: 638-640.
- Tsuyuki, H., Roberts, E. and Gadd, R.E.A. (1962). Muscle protein of Pacific Salmon (*Oncorhynchus*). III. The separation of muscle proteins soluble in low ionic strength salt solutions by starch gel electrophoresis. Canadian J. Biochem. Physiol., 40: 929 - 936.
- Ukishima, Y., Naritan, H., Masui, T. and Naser, M. (1984). Studies on the application of thin layer isoelectric focusing to food analysis. I. Identification of species of shiced escolar and castor of oil fish. Eisei. Kagaku. Vol. 30 (4): 189 - 193.
- Ursin, E. (1967). A mathematical model of some aspects of fish growth, respiration and mortality. J. Fish. Res. Bd. Can. (24): 2355-253 .
- Utter, F.M., Hodgins, H.O. and Allendorf, F.W. (1974). Biochemical genetic studies of fishes: potentialities and limitations. In: Malins and Sargent (Eds.) Biochemical and Biophysical Prespectives in Marine Biology, Vol.1, Academic Press, New York.
- Van Oosten, J. (1929). Life history of the lake herring (*Leucichthys artedi* Le Sueur) of lake Huron as revealed by its scales, with a critique of the scale method. Bull. Bur. Fish Wash., 44: 265-428.
-

- Verheyen, E., Van-Rompaey, J., Blust, R. and Decleir, W. (1991). The eye lens protein pattern of eight Lake Tanganyika cichlids studied by isoelectric focusing. Koninklijk Museum Voor Midden-Afrika Tervuren België Annalen Zoologische Wetenschappen 263 (0): 29-32.
- Vinogradov, A. P. (1953). The elementary chemical composition of marine organisms. Sears Foundation, New Haven. 647 p.
- Von Bertalanffy, A. (1938). A quantitative theory of organic growth. Hum. Biol., 10 (2): 181 - 182.
- Walford, L.A. (1946). A new graphic method of describing the growth of animals. Biol. Bull. Mar. Biol. Lab. Woodstole, 90: 141-147.
- Wassef, E.A. and Abdel-Hady, H.A. (1997). Breeding biology of rabbit fish *Siganus canaliculatus* (Siganidae) in mid Arabian Gulf. Fisheries Research, 33: 159-166.
- Weatherly, A. H. (1972). Growth and ecology of fish population. Academic press. London and New York, 291 p.
- Westernhagen, H. von (1973 a). A preliminary study on the food preferences of *Siganus concatenata* (Cuvier and Valenciennes). Philipp. Scient. 10: 61-73.

- Westernhagen, H. von (1973 b). The natural food of the rabbit fish *Siganus oramin* and *S. striolata*. Mar. Biol. 22: 367-370.
- Whitmore, D. W. (1986). Identification of Sunfish species by muscle protein isoelectric focusing. Comp. Biochem. Physiol., 84B: 177 - 180.
- Willer, A. (1929). Untersuchungen über das Wachstum von Fischen. Verh. Int. Ver. Limnol., 4: 668-684.
- Windell, J. T and Bowen, S. H. (1978). Methods for study of fish diets based on analysis of stomach contents. In: Methods for the assessment of fish production in fresh waters, p. 219-226 (Ed. Bagenel) Blackwell Scientific Publications, Oxford..
- Wood, H. (1930). Scottish herring shoals prespawning and spawning movements. Scotland. Fish. Bd. Sci. Invest. (1): 1 - 71.
- Wootton, R. J. (1990). Ecology of Teleost fishes (Eds. Chapman and Hall), Fish and Fisheries series I, 403 p.
- Yamada, J. and A. Suzuki. (1982). Identification of fish species by thin layer isoelectric focusing of sarcoplasmic proteins. Bull. Jap. Soc. Sci. Fish., 48 (1): 73 - 77.
-
- Not consulted in origin.