

REFERENCES

= = = = =

(A)

Addison , A. ; Fixsen , J. and Lioyed , R. (1983) :

A review of the Dillwyn Evans type collateral operation in sever clubfoot.

J. of Bone and Joint Surgery, 65 B : 12 .

(B)

Bensahel ,H. ; Catterall , A. and Chir , M. (1990) :

Practical application in idiopathic clubfoot.

J. of Pediatric orthop. , 10 : 186 .

Berman , A. ; Arnold, S. and Gartland , J.J. (1971) :

Metatarsal osteotomy for correction of adduction of the forepart of the foot in children.

J. of Bone and Joint Surg., 53-A : 498

(C)

Carroll , N.C. ; McMurty, R. and Leete, S.F. (1978) :

The pathoanatomy of the congenital clubfoot.

Orthopaedic Clinic Of North America. , 9 : 225.

Carroll , N.C. (1987) :

Congenital clubfoot , pathoanatomy and treatment.
Orthopaedic Clinic Of North America., 6 : 245.

Carroll , N.C. (1988) :

Pathoanatomy and surgical treatment of resistant clubfoot.
Orthopaedic Clinic Of North America., 8 : 116.

Carroll , N.C. (1990) :

Clubfoot . In : Raymond , T. and Morrissy , M.D. (Eds.) : Lovell and Winter's Pediatric Orthopedic , Third Edition.
Philadelphia and Lippincott. Pp : 927 .

Clark, C.E.;Cowell , H.R. and Telfer,M.A. et al ., (1980) :

Trisomy 6Q 25 - 6Q TER in two sisters resulting from maternal 6; 11 translocation.
American J. Med. Jenetic, 5 : 171.

Cowell , H.R. and Wein , B.K. (1980) :

Genetic aspects of clubfoot.
J. of Bone and Joint Surg. , 62 A : 1381 .

Crawford , A.H. ; Marxen,J.I. and Osterfeld,D.L. (1982) :

The cincinati incision ; A comperhensive approach for surgical procedures of the foot and ankle in childhood.
J. of Bone and Joint Surg., 64 A : 1355.

Cummings, R.J. (1987) :

The circumferential clubfoot release.
J. of Bone and Joint Surgery, 67-A :1044.

Cummings, R.J. and Lovell, W.W. (1988):

Operative treatment of congenital idiopathic clubfoot.
J. of Bone and Joint Surg. 70A:1108.

(D)

Depuy, J. and Drennan, J.C. (1989):

Correction of Idiopathic clubfoot : a comparison of results of early versus delayed posteromedial release.
J. Pediatric Orthop., 9:44.

Diepstraten, A.D. (1996) :

Congenital clubfoot.
Acta Orthop. Scand. , 67 : 305

Dravaric, D.M. (1989):

Congenital clubfoot.
Orthopaedic Clinic Of North America, 20:641.

Dwyer, F.C. (1963):

The treatment of relapsed clubfoot by the insertion of wedge into calcaneum.
J. of Bone and Joint Surg., 45B:67.

(E)

Ivans, A. and Dillwyn, A. (1961):

Relapsed clubfoot.

J of Bone and Joint Surg., 43B :722.

(F)

Line, R.N.; Gwinn, J.L. and Young, E.F. (1968):

Smith-Lemli-Opitz Syndrome.

American J. Dis. Child., 115:482.

Fukuhara, K.; Schdlmeier, G. and Uhthoff, H. (1994):

The pathogenesis of the clubfoot

J. of Bone and Joint surgery, 76 : 450.

(G)

Galdino, M.J.; Stifdf, S.J. and Butler, J.E. et al ., (1978):

*Triple arthrodesis in severely affected feet ; a salvage procedure
after failed release in severely affected feet.*

Foot and Ankle, 7:319.

Gorlin,R.J.;Cervenka,J. and Anderson,R.C. et al., (1970):

Robin's Syndrome.

American J.Dis. Child.,119:176.

Gray,H.(1966):

Anatomy of human body.

Gross,C.M. (Edin). Philadelphia , Lea and Febiger.

Grill,F. and Franke,J.(1987):

The ilizarov distractor for the correction of relapsed or neglected clubfoot.

J. of Bone and Joint Surg.,69B :593.

(H)

Hadidi, H.(1974):

Management of congenital talipes equino varus.

Orthopedic Clinic of North America,5:53.

Herzenberg,J.E.;Carroll,N.C. and Christoferson,M.R. et al .,(1988):

Clubfoot analysis with three-dimensional computer modeling.

J. of Pediatric Orthopaedic ,8:257.

Heyman, C.H.; Herndon, C.H. and Strong, J.M. (1958):

Mobilization of the tarsometatarsal and inter-metatarsal joints for the correction of resistant adduction of the forepart of the foot in congenital clubfoot or congenital metatarsus varus.

J. of Bone and Joint Surg., 40 :299.

Hjelmstedt, A. and Sahlstedt, B. (1978) :

Simultaneous arthrography of the talocrural and talo-navicular joints in children.

Acta Radiol., 19:223.

Hjelmstedt, A. and Sahlstedt, B. (1990):

Arthrography as a guide in treatment of congenital clubfoot.

Acta Orthop. Scand., 51:321.

Hoffman, A.A. (1984):

Osteotomy of the first cuneiform as treatment of residual adduction of the forepart of the foot in clubfoot.

J. of Bone and Joint surgery, 66 A:985.

(I)

Inclan, A. (1958):

Anomalous tendinous insertion in the pathogenesis of the clubfoot.

J. of Bone and Joint surgery, 40 :159.

Ippolito, E. and Ponseti, I. (1980) :

Congenital clubfoot in the human fetus : a histological study.
J. of Bone and Joint surgery, 62:8 .

Irani, R.N. and Sherman, S.S. (1963):

The pathological anatomy of the clubfoot.
J. of Bone and Joint surgery, 45:45.

Issac, H.; Handelsman, J.E.; Badenhort, M. and Pickering, A. (1977):

The muscle in clubfoot.
J. of Bone and Joint surgery, 59 :465.

(f)

Jago, R.H. (1970):

Arthrogryposis following treatment of maternal tetanus with muscle relaxants.
Arch. Dis. Child., 45 :277.

John, G. and Simons, W.G. (1993):

Deformity of the calcaneo-cuboid joint in patients who have Talipes Equino Varus.
J. of Bone and Joint surgery, 75 : 190.

(K)

Karski, T. and Wosko, I. (1989):

Experience in the conservative treatment of congenital clubfoot in newborn and infants.

J. Pediatric of Orthop., 9:134.

(L)

Last, R.J. (1994):

Anatomy of the foot and ankle. In : McMinn, R.M.H. (Ed.) : Last's Anatomy, Regional and Applied, Ninth Edition.

Edinburgh, London, Madrid, Melbourne, New York and Tokyo.

Churchill Livingstone.

Lichtblau, S. (1973) :

A medial and lateral release operation for clubfoot, a preliminary report.

J. of Bone and joint surgery, 55A : 1377.

Lundberg, B. J. (1981) :

Early Dwyer operation in talipes equino varus.

Clinical Orthopaedics and Related Research, 154 : 223.

(m)

Macleod, P. and Partriquin, H. (1974) :

The whistling face syndrome.

Clin. pediatr. 13 : 184-189.

Magone, J. B. ; Torch, M. A. and Clark, R. N. (1989) :

Comparitive review of surgical treatment of the idiopathic clubfoot by three different procedures.

J. pediatric orthop. vol. 9 No. 1, 49.

McKay, D. W. (1982) :

New concept of and approach to clubfoot treatment .

Section I .

J. pediatric of orthop. 2 : 374 .

McKay, D. W. (1983) :

New concept of and approach to clubfoot treatment .

Section II . Correction of the clubfoot .

J. pediatric orthop. , 3 : 10.

Michael, J. B. (1991) :

Combined split anterior tibial tendon transfere and intramuscular lengthening of posterior tibial tendon.

J. of Bone and joint surgery vol. 73 A, No. 5: 734.

(N)

Napiontic, M. and Nazar, J. (1994) :

Tibial steotomy as a salvage procedure in the treatment of congenital talipes equino varus.

J. pediatric orthop.; 14 (6) : 763.

Napiontic, M. (1995) :

Intraoperative ultrasound for evaluation of reduction in congenital talipes equinovarus.

J. of Pediatric of Orthopaedic , 4 (1) : 55 ..

Nather , A . and Bose , K. (1987) :

Conservative and surgical treatment of clubfoot.

J. of Pediatric of Orthopaedic , 7: 42.

Nichols,E.H. (1987):

Anatomy of congenital equino-varus.

Boston Med. J. of Surgery , 36: 150.

(O)

Olavar,R.(1989):

Torsion of the leg determined by C.T. scan.

Acta Orthop. Scand.,3:330.

(P)

Price, C.T.(1978):

Non-operative treatment of congenital clubfoot.

A long-term follow up study.

J. of Bone and Joint Surg., 53-A : 498 .

(R)

Reimann, I.(1967):

Congenital idiopathic clubfoot.

American J. Dis. Child., 121 : 366

Robertson, F.W.; Kozlowski, K. and Middleton, R.W. (1975) :

Larsen's Syndrom.

Clinic. Pediatric , 14: 53.

Ronald, M.C.(1992):

foot . In : Ronald McRae (Ed.) : Clinical orthopaedic examination.

Third Edition. Churchill Livingstone. 243..

(S)

Shaw, E.B. and Steinbach, H.L. (1968):

Aminopterin-induced fetal malformation.
American J. Dis. Child., 115 : 477.

Simons, G.W. (1985):

Complete subtalar release in clubfoot.
Part I. A preliminary report.
J. of Bone and Joint Surgery, 67-A : 1044.

Simons, G.W. (1987) :

The complete subtalar release in clubfoot.
Orthopaedic Clinics of North America, 18 : 43 .

Sodre, H. ; Bruschini, S. and Mestriner, A. et al ., (1990):

Arterial abnormalities talipes equinovarus as assessed by angiography and Doppler technique.
J. of Pediatric of Orthopaedic, 10 : 101.

(J)

Tachdjian, M.O. (1985):

Congenital deformities. In Tachdjian, M.O. (Ed.) : Pediatric Orthopaedic Philadelphia, W.B. Saunders Pp : 139.

Tachdjian,M.O.(1986):

Occult subluxation of the subtalar joint in clubfoot using computerized tomography.

J. of Pediatric of Orthopaedic ,6 : 334.

Thomas,D.;Grider,A.;Sherwainy,J. and Paul,G.(1982):

Arteriography in clubfoot.

J. of Bone and Joint Surgery, 64 A : 154.

Thomposon,G.H.;Richardson,A.B. and Wastin,G.W.(1982):

Surgical management of resistant congenital talipes equino varus deformity.

J. of Bone and Joint Surgery, 64 A :652.

Turco,V.J.(1971):

Surgical correction of the resistant clubfoot.

J. of Bone and Joint Surgery, 61 A :805.

Turco,V.J.(1979):

Resistant congenital clubfoot , One stage posteromedial release with internal fixation.

J. of Bone and Joint Surgery, 61 A , 805 .

Turco, V.J. (1981):

Clubfoot.

First Edition.

NewYork , London , Edinburgh and Melbourne.

Churchill Livingstone.

(V)

Vicente, R. G. (1996):

Clubfeet in congenital annular constricting bands.

Clinical Orthopaedics and Related Research, 323 : 155.

(W)

Weintroub, S. and Kherrmash, M.D. (1994):

Comparative evaluation of initial surgical procedure in clubfoot.

J. of Pediatric Orthopaedics, 3 : 171.

Williams, L.B.; Bannister, H.B.; Berry, M.M. and Collins, P. (1995):

Angiology In : Williams, L.B.; Bannister, H.B.; Berry, M.M. and Collins, P.

(Eds.) : Gray's Anatomy, Thirty - Eighth Edition.

ELBS, with Churchill, Livingstone. Pp : 432.

Wynne- Davies, R. (1972) :

Genetic and environmental factors in the aetiology of talipes equino-varus.

Clinical Orthopaedics and Related Research, 84 : 9.

(2)

Zimny, M.L.; Willig, S.J. and Roberts, J.M. et al ., (1985):

An electron microscopic study of the fascia from the medial and lateral sides of clubfoot.

J. of Pediatric of Orthopaedic , 5 : 577.