

REFERENCES

- Abou-Bakr, H. (1984). Studies on certain biological control agents of mosquitoes in Egypt. Ph.D. Thesis, Fac. Agric., Cairo Univ. 220 PP.
- Adams, J.B. (1960). Effect of spraying 2,4-D amine on coccinellid larvae. Can. J. Zool., 38 : 285-288.
- Akhurst, R.J. (1980). Morphological and functional dimorphism in Xenorhabdus spp., bacteria symbiotically associated with the insect pathogenic nematodes Neoaplectana and Heterorhabditis. J. Gen. Microbiol., 121 : 303-309.
- Altman, J. and Campbell, C.L. (1977^a). Ann. Rev. Phytopath. 15 : 361-385. (Cited by Robert, F.N. (1982) op.cit.).
- (1977^b). Phytopath., 67 : 1163-1165. (Cited by Robert, F.N. (1982) op.cit.).
- Anonymous (1972). Mass-produced nematodes down mosquito. Agric. Res., 20 (11) : 3-4.
- Anonymous (1979). Methazole. In "Herbicide Hand-book", weed. Sci. Soc. Amer. Champaign, III : 268-274.
- Azzam, K. (1987). Studies on the predator insects of Bilharziasis snails in Egypt. M.Sc. Thesis, Fac. Agric., Cairo Univ., p. 42-70.
- Baldwin, W.F.; James, H.G. and Welch, H.E. (1955). A study of mosquito larvae and pupae with a radio-active tracer. Canad. Ent., 87 : 350-356.
- Bates, M. (1965). "The Natural History of Mosquito". Harper Trochbooks. Harper & Row publishers, New York, pp. 157-162.

- Bay, E.C. (1974). Predator-prey relationships among aquatic insects. Ann. Rev. Entomol., 19 : 441-453.
- , and Anderson, L.D. (1965). Studies with the mosquito fish, Gambusia affinis as a chironomid control. Ann. Entomol. Soc. Amer., 59 : 150-135.
- Bishop, E.L. (1940). Cooperative investigations of the relation between mosquito control and wildlife conservation. Sci., 92 : 201-202.
- Bucher, G.E. (1960). potential bacterial pathogens of insects and their characteristic. J. Insect Pathol., 2 : 172-195.
- Burman, M. (1982). Neoaeplectana carpocapsae : toxin production by axenic insects parasitic nematodes. Nematologica, 28 : 62-70.
- Corbet, P.S. (1962). "A Biology of Dragonflies". Witherby, London, 247 pp.
- Corbett, J.R. (1974). "The Biochemical Mode of Action of Pesticides". Academic Press, London and New York, pp 96
- Cotz, P. (1969). Die Einkasierung von parasiten in der Hamolymphe von Chironomus-larven (Diptera). Zool. Anz. Suppl., 33, 610. (Cited by Poinar, G.O. (1979) op.cit.).
- Cremlyn, R. (1978). "Pesticides Preparation and Mode of Action". John Wiley & Sons, Ltd., pp. 140-172.
- Dadd, R.H. (1971). Effect on size and concentration of particles on rates of ingestion of latex particulates, by mosquito larva. Annals Entomol. Soc. Amer., 64 : 687-692.
- Darwazeh, H.A. and Mulla, M.S. (1982). Encapsulated formulations of pirimiphos-methyl against mosquito larvae and some nontarget organisms. Proc. Pap. 49th Ann. Conf. Calif. Mosq. & Vector Contr. Assoc., pp. 26-29.

- Dutcky, S.R. (1959). Insect Microbiology. Adv. appl. Microbiol., 1 : 175-200.
- Dutcky, S.R. (1974). Nematode parasite, p. 576-590. In : P.G. Maxwell and F.A. Harris (eds.). Proc. of the Summer Institute on Biological Control of Plant Insects and Diseases. Univ. Press of Mississippi, Jackson.
- Eisler, R. (1970). Acute toxicities of organochlorine and organophosphorus insecticides to Estuarine Fishes. U.S. Dept. Interior. Bur. Sport Fish Wildl. Tech. Paper, 46 : 12 pp.
- El-Kifl, T.A. (1980). Utilization of nematode Neosaplectana carpocapsae in The biological control of cotton leaf-worm Spodoptera littoralis. Ph.D. Thesis, Fac. Agric., Cairo Univ., p. 36-39.
- El-Kifl, T.A. and El-Sherif, M.A. (1986). Dispersal potentiality of the infective juveniles of the entomogenous nematodes, Neosaplectana carpocapsae. Bull. Fac. Agric. Cairo Univ., 37 (2) : 1085-1088.
- El-Rayah, El-Amin (1975). Dragonfly nymphs as active predators of mosquito larvae. Mosq. News, 35 (2) : 229-230.
- Eysell, A. (1905). "Die Stechmacken". In "Handbook der tropenkrankheiten" (Mense, C.ed.), Leipzig, Barth, 2 : 78-80.
- Fales, J.H.; Spangler, P.J.; Bodenstein, O.; Mills, G.D.Jr. Durbin, C.G.Jr. (1968). Laboratory and field evaluation of a bate against a backswimmer, Notonecta undulata Say (Hemiptera : Notonectidae). Mosq. News, 28 (1) : 77-81.
- Farghal, A.I.A. (1979). Recent trends in culicine mosquito control. Ph.D. Thesis, Fac. Agric., Assiut Univ., 181 pp.

Pedroko, A.; Kamionek, M.; Kozłowska, J. Mianowska, E. (1977^a).
The effect of some insecticides on nematodes of different
ecological groups. Pol. Ecol. Stud., 2 : 78-88.

----- (1977^b).

The effect of Vydate oxamyl on nematodes of different
ecological groups. Ibid, 2 : 89-93.

Frank, P.A. (1972). Herbicidal residues in aquatic environments. In "Fate of Organic Pesticides in the Aquatic Environment". Adv. Chem. Ser. III, pp. 135-148.

Gerberg, E.J.; Hopkins, T.M. and Gentry, J.W. (1969). Mass rearing of Culex pipiens L. Mosq. News, 29 (3) : 382-385.

Greekmur, G.D.; Russell, M.P. and Hazelrigg, J.E. (1982).
Field evaluation of the effect of slow release wettable powder formulation of Altosid on nontarget organisms. Proc. Pap. 49th Ann. Conf. Calif. Mosq. & Vector Contr. Assoc., 26-29.

Hara, A.H. and Kaya, H.K. (1982). Effect of selected insecticides and nematicides on the in vivo development of the entomogenous nematode, Neaplectana carpocapsae. J. Nematol., 14 (4) : 486-491.

----- (1983^a). Toxicity of selected organophosphate and carbamate pesticides to infective juveniles of the entomogenous nematode N. carpocapsae (Rhabditida : Steinernematidae). Environ. Entomol., 12 (2) : 496-501.

----- (1983^b). Development of the entomogenous nematode, Neaplectana carpocapsae (Rhabditidae : Steinernematidae) in insecticide-killed Beet armyworm (Lepidoptera : Noctuidae). J. Econ. Entomol., 76 (3) : 423-426.

- Harp, G.L. and Campbell, R.S. (1964). Effect of the herbicides silvex on benthos of a farm pond. J. Wildl. Manage., 28 : 308-317.
- Harvey, R.G.; Hagedorn, D.J. and Deloughery, R.L. (1975). Crop Sci., 15 ; 67-71. (Cited by Robert, F.N. (1982) op.cit.).
- Heitfuss, R. (1970). Z. Pflkrankh. Pflschutz. Sonderheft., 5 : 117-127. (Cited by Robert, F.N. (1982) op.cit.)
- (1972). Z. Pflkrankh. Pflschutz. Sonderheft., 6 : 79-87. (Cited by Robert, F.N. (1982) op.cit.).
- Hess, A.D. and Keener, G.G. (1947). Effect of airplane-distributed DDT thermal aerosols on fish and fish food organisms. J. Wild. Manage., 11 : 1-10.
- Hilsenhoff, W. (1966). Effect of diquat on aquatic insects and related animals. J. Econ. Entomol., 59 : 1520-1521.
- Hinman, E.H. (1934). Predators of Culicidae. I. The predators of larvae and pupae, exclusive of fish. J. Trop. Med. & Hyg., 37 : 129-134.
- Hooper, F.N. (1958). The effect of application of pelleted 2,4-D upon the bottom fauna of Kent Lake, Michigan. North Centre. Weed Control Conf., Proc., 15-41.
- Howard, L.O.; Dyar, H.G. and Knab, F. (1912). The mosquitoes of N. & C. America and the West Indies. Washington, Carnegie Inst., 1 : 156-179. (Cited by Abou-Bakr, H. (1984) op.cit.).
- Hurlbert, S.H. (1975). Secondary effects of pesticides on aquatic ecosystems. Residue Rev., 57 : 81-148.
- Hynes, H.B.N. (1961). The effect of sheep-dip containing the insecticides BHC on the fauna of a small stream. Ann. Trop. Med. Parasit., 55 : 192-196.

- Hynes, H.B.N. and Williams, T.R. (1962). The effect of DDT on the fauna of a Central African Stream. Ann. Trop. Med. Parasit., 56 : 78-91.
- Ignoffo, C.M.; Biever, K.D.; Johnson, W.W.; Sanders, H.O.; Chapman, H.C.; Petersen, J.J. and Woodward, D.B. (1973). Susceptibility of aquatic vertebrates and invertebrates to the infective stage of the mosquito nematode, Romanomermis culicivorax. Mosq. News, 33 (4) : 599-602.
- Isaeva, N.M. (1975). A study of Culicimermis schakhovii Rubtsov et Isaeva, 1975 (Mermithidae), a parasite of mosquitoes. Problemy parazitologu. Materialy VIII nauchnoi konferentsy parazitologov ukssr. chast'1, Kiev, USSR; Iz-datel's tvo : 197-198.
- (1977). Aedes caspius caspius Pall (Culicidae), a new host of Culicimermis scksknovii (Nematoda : Mermithidae). Venstnik Zoologii, 6 : 88-89.
- Jeen, R.F.; Roger, G.; William, J.C. and Thomson, N.R. (1977). Laboratory studies on the feasibility of integrated mosquito control using an insect growth regulator and a mermethid nematode. Mosq. News, 37 (1) : 6-11.
- Jenkins, D.W. (1964). Pathogens, parasites and predators of medically important arthropods. Bull. Who, 30 : 1-59.
- Kamionek, M. (1979). The influence of pesticides on the mortality and effectiveness of Neosaplectana carpocapsae Weiser, p. 87-88. In Proc. Inter. Colloq. Invertebr. Pathol. & 11th Ann. Meet. Soc. Invertebr. Pathol., Prague, Czech., pp. 87-88.
- Katan, J. and Eshel, Y. (1973). Res. Rev., 45 : 145-177. (Cited by Robert, F.N. (1982), op.cit.).
- Kavanagh, T. (1969). Sci. Proc. Royal Dublin Soc. Ser. B-2, 179-190.. (Cited by Robert, F.N. (1982), op.cit.).

- Kavonagh, T. (1974). Ibid. Ser. B-3 : 251-265. (Cited by Robert, F.N. (1982), op.cit.).
- Kearney, P.C. and Kaufmann, D.D. (1975). "Herbicides-Chemistry, Degradation and Mode of Action". 2nd ed., Vol. 1, Dekker, New York. (Cited, by Cremlyn R. (1978) op.cit.)
- Kennedy, H.D.; Eller, L.F. and Walsh, D.F. (1970). Chronic effects of Methoxychlor on Blugills and aquatic invertebrates. U.S. Dept. Int., Bur. Sport Fish Wild. Tech. Paper, 53, 18 pp.
- Laird, M. (1947). Some natural enemies of mosquitoes in the vicinity of palmalmal, New Britain. Trans. Roy. Soc. N.Z., 76 : 453-476.
- Lee, F.C. (1967). Laboratory observations on certain mosquito larval predators. Mosq. News, 27 (3) : 332-338.
- Levy, R. and Miller, T.W. (1977). Susceptibility of mosquito nematode Romanomermis culicivorax (Mermithidae) to pesticides and growth regulators. Environ. Entomol., 6 : 447-448.
- .; Powell, C.M. and Miller, T.W.Jr. (1985). Persistence of the mosquito larvicide and pupicide Arosury Msf in permanent and semi-permanent habitats. J. Florida Ant. Mosq. Assoc., 56 (1) : 32-36.
- Lewallen, L.L. (1962). Toxicity of certain insecticides to Hydrophilid larvae. Mosq. News, 22 : 112-113.
- Little, V.A. (1963). "General and Applied Entomology". 2nd edition, Oxford & IBH Publishing Co., 195 pp.
- Lutfallah, A.F. (1974). Studies on the aquatic insects in rice nurseries and fields. M.Sc. Thesis, Fac. Agric., Cairo Univ., 168 pp.

- Lysenko, Q. (1966). Bacterial toxins. In "Insect Pathology and Microbial Control". Van der Loan, P.A. ed., North Holland, Amsterdam, 214-237.
- Lysenko, O. and Weiser, J. (1974). Bacteria associated with the nematode Neosaplectana carpocapsae and the pathogenicity of this complex for Galleria mellonella larvae. J. Invertebr. Pathol., 35 : 256-259.
- Mather, T.H. (1984). "Environmental Management for vector Control in Rice Fields, pp. 14-17.
- McCraren, J.P.; Cope, O.B. and Eller, L. (1969). Some chronic effects of diuran on bluegills. Weed Sci., 17 : 497-504.
- McGarvey, B.D.; Potter, J.W. and Chiba, M. (1984). Nematostatic activity of oxamyl and N,N-Dimethyl-1-cyanofarmamide (DMCF) on Meloidogyne incognita juveniles. J. Nematol., 16 (3) : 328-331.
- Milstead, J.E. and Poinar, G.Q.Jr. (1978). A new entomogenous nematode for pest management systems. Calif. Agric., 32 : 12.
- Mitchell, C.J.; Chen, P.S. and Chapman, H.C. (1974). Exploratory trials utilized a mermithid nematode as a control agent for Culex mosquitoes in Taiwan. J. Formosan Med. Assoc., 73, 241.
- Miura, T. and Takahashi, R.M. (1973). Insect development inhibitors. 3. Effect on nontarget aquatic organisms. J. Econ. Entomol., 66 : 917-922.
- (1974). Insect developmental inhibitors : Effects of candidate mosquito control compounds on nontarget aquatic organisms. Environ. Entomol., 4 : 631-636.

Miura, T. and Takahashi, R.M. (1976). Effect of a synthetic pyrethroid, SD-3775, on nontarget organisms when utilized as a mosquito larvicides. Mosq. News, 36 (3) : 322-326.

----- (1987). Impact of fenoxycarb, a carbamate insect growth regulator on some aquatic invertebrate abundant in mosquito breeding habitat. J. Amer. Mosq. Contr. Assoc., 3 (3) : 476-480.

Miura, T.; Takahashi, R.M. and Mulligan, F.S., III (1978). Field evaluation of the effectiveness of predacious insects as a mosquito control agent. Proc. Pap., 46th Ann. Conf. Calif. Mosq. & Vector Contr. Assoc., pp. 80-81.

-----; Walker, W.H.; Grant, C.D.; Combs, J.C.; Coykendall, R.L.; Lusk, E.E. and Washino, R.K. (1983). The effect of BAY SIR 8514 against selected organisms associated with mosquito breeding habitats. Proc. Pap. 51th Ann. Conf. Calif. Mosq. & Vector Contr. Assoc., pp. 23-26.

Muirhead-Thompson, R.C. (1971). "Pesticides and Fresh-water Fauna". Academic Press, pp. 67-70.

Mulla, M.S.; Arias, J.R. and Darwazeh, H.A. (1971). Petroleum oil formulations against mosquitoes and their effects on some nontarget insects. Proc. Calif. Mosq. Control Assoc., 39 : 131-135.

----- and Darwazeh, H.A. (1971). Influence of aliphatic amines-petroleum oil formulations on aquatic non-target insects. Proc. Calif. Mosq. Assoc., 39 : 126-131.

----- (1982). Efficacy of petroleum larvicidal oils and their impact on some aquatic non-target organisms. Proc. Pap. 49th Ann. Conf. Calif. Mosq. & Vector Contr. Assoc., pp. 26-29.

- Mulla, M.S.; Darwazeh, H.A. and Ede, L. (1982). Evaluation of new pyrethroid against immature mosquitoes and their effects on nontarget organisms. Mosq. News, 42 (4) : 583-590.
- Mulla, M.S.; Sjogren, R.D. and Arias, J.R. (1972). Mosquito larvicides : Efficacy under field conditions and effects on non target organisms. Proc. Calif. Mosq. Control Assoc., 40 : 96-100.
- Muller, G. (1971). Laboruntersuchungen zur wirkung von Herbiziden auf carabiden. Arch. Pflanzenschutz, 7 : 351-367. (Cited by Brown, A. (1978) in "Ecology of Pesticides", John Wiley & Sons, Inc.).
- Mullison, W.R. (1970). Effect of herbicides on water and its inhabitants. Weed Sci., 18 : 738-750.
- Newman, J.F. and Way, J.M. (1966). Some ecological observations in the use of paraquat and diquat as aquatic herbicides. Br. Weed Contr. Conf. Proc., 583-585.
- Nicle, W.R. (1980). "Possible Commercial Formulations of Insect Parasitic Nematodes". Biotechnology and Bioengineering, Vol. XXII : 1407-1414.
- Notestine, M.K. (1971). Population densities of known invertebrate predators of mosquito larvae in Utah marshlands. Mosq. News, 31 (3) : 331-334.
- Oka, I.N. and Pimental, D. (1976). Science, 193 : 239-240. (Cited by Robert, F.N. (1982) op.cit.).
- Otieno, W.A. (1977). The pathology and host-parasite relationship of the entomogenous nematode Romanomermis culicivorax Ross and Smith (Mermithidae) in a mosquito host Culex pipiens Say (Culicidae) : Its application in integrated control. Ph.D. Thesis, Univ. of Calif., Berkley, 241 pp.

- Petersen, B.V. and Davis, D.M. (1960). Observations on some insect predators of black flies of Algonquim park, Ontario. Can. J. Zool., 38 : 9-18.
- Petersen, J.J. and Chapman, H.C. (1979). Checklist of mosquito species rested against the nematode parasite Romanomermis culicivorax. J. Med. Entomol., 15 (5) : 468-471.
- and Willis, Q.R. (1970). Some factor affecting parasitism by mermithid nematodes in southern house mosquito larvae. J. Econ. Entomol., 63 : 175.
- Pierce, M. (1960). Progress report of the effect of known on the biota of long pond, Dutchess county. N.E. Weed Contr. Conf. Proc., 14 : 472-475 and 338-343.
- Pimental, D. (1971). Ecological effect of pesticides on nontarget species. Office Sci. Tech., Exec. Office President, Govt Office, Washington, D.C., 220 pp.
- Poinar, G.Q.Jr. (1975). "Entomogenous nematodes". A manual and host list of insect nematode associations. Leiden E.J. brill, 317 pp.
- (1976). Insect immunity to parasitic nematodes. In : "Contemporary Topics in Immunobiology". (Cooper, E.L., ed.). Plenum press, New York, 1976 pp.
- (1979). "Nematode for Biological Control of Insects". CRC Press, Inc. Boca Raton, Florida. 249 pp.
- and Leutenegger, R. (1968). Anatomy of the infective and normal third stage juveniles of Neoaplectana carpocapsae Weiser (Steinernematidae - Nematoda). J. Parasitol., 54 : 340-346.
- (1971). Ultrastructural investigation of the melanization process in Culex pipiens (Culicidae) in response to a nematode. J. Ultrastruct. Res., 36 : 149-152.

- Poinar, G.O.Jr. and Thomas, G.M. (1966). Significance of Achromobacter nematophilus Poinar and Thomas (Achromobacteraceae : Eubacteriales) in the development of the nematode, DD-136 (Neosaplectana sp. Steinernematidae). Parasitology; 56 : 385-390.
- (1967). The nature of Achromobacter nematophilus as an insect pathogen. J. Invertebr. Pathol., 9 : 510-514.
- Pokasa Rao, P.S.; Das, P.K. and Padhi, G. (1975). Note on compatibility of DD-136 (Neosaplectana dutkyi), an insect parasitic nematode with some insecticides and fertilizers. Indian J. Agric. Sci., 45 : 275-277.
- Pritchard, G. (1964). The prey of the dragonfly nymphs in ponds in northern Alberta. Can. J. Zool., 42 : 785-800.
- Ramalingam, S. (1979). Prospects offered by the use of invertebrate predators for mosquito control with special emphasis on *Luzia* and *Toxorhynchites*. WHO/TDR/PCV/SWG, 79/WP/20.
- Reed, E.M. and Carne, P.B. (1967). The suitability of a nematode (DD-136) for the control of some pasture insects. J. Invertebr. Pathol., 9 : 196.
- Riedel, H.S. (1973). Cumulative Bibliography 1953-1972 parasites, predators, viruses and related agents Biological control of mosquitoes). Mosq. News, 33 (2) : 301.
- Robert, F.N. (1982). Interaction between weeds and other pests in the Agro-ecosystem. In "Biometeorology in integrated pest management" (Hatfield, J.L. and Thomson, I.J. eds.), pp. 343-397, Academic press, Inc.

- Romney, R.K.; Anderson, J.L. and Griffin, G.D. (1974). Weed Sci., 22 : 51-54. (Cited by Robert, F.N. (1982) Ibid.).
- Savage, K.E. and Petersen, J.J. (1971). Observations of mermithid nematodes in Florida mosquitoes. Mosq. News, 31 : 218-219.
- Sailer, R.I. and Lienk, S.E. (1954). Insect predators of mosquito larvae and pupae in Alaska. Mosq. News, 14 : 14-16.
- Schaefer, C.H.; Miura, T.; Stewart, R.J.; Dupras, E.F. Jr.; Grant, C.D.; Washino, R.K.; Lusk, E.E. and Coykendall, R.L. (1983). Studies on the potential environmental impact of the thiobencarb 'Bolero'. 38th Ann. Meet. Amer. Mosq. Contr. Assoc., p18-22.
- Schroeder, W.J. and Beavers, J.B. (1987). Movement of the entomogenous nematode of the families Heterorhabditidae and Steinernematidae in soil. J. Nematol., 19 (2) : 257-259.
- Scudder, G.G.E. (1976). Water-boatman of saline waters (Hemiptera : Corixidae). In "Marine Insects". (L. Cheng, ed.), pp. 263-289. North-Holland publishing Co.
- Serycznska, H. (1977). Defensive reactions of Galleria mellonella L. caterpillars to the effect of herbicides and the nematode N. carpocapsae Weiser. Bull. Acad. Pol. Sci., Ser. Sci. Biol., 25 : 749.
- Shalaby, F. (1958). Alphabetical list of Egyptian insects in the collection of the Ministry of Agriculture. Tech. Bull. No. 284.
- Smith, Y.B. (1904). Report of the New Jersey State Agricultural Experimenta Station upon the mosquitoes occurring within the state, their habitats, life history, etc. Trention, 482 pp.

- Smith, G.E. and Isom, B.G. (1967). Investigation of effect of large-scale applications of 2,4-D on aquatic fauna and water quality. Pestic. Monit. J., 1 (3) : 16-21.
- Stoffolano, J.G.Jr. and Streams, F.A. (1971). Host reactions of Musca domestica, Orthelia caesarion and Ravinia l'herminieri to the nematode Heterotylenchus autumnalis. Parasitology, 63 : 195.
- Takahashi, R.M.; Wilder, W.H. and Miura, T. (1984). Field evaluation of ISA-20 E for mosquito and effects on aquatic nontarget arthropods in experimental plots. Mosq. News, 44 (3) : 363-367.
- Tarzwell, C.M. (1947). The effects on surface organisms of the routine land application of DDT larvicides for mosquito control. U.S. Public Health Reports, 62 : 525-554.
- Tatum, W.M. and Blackburn, R.D. (1962). Preliminary study of the effects of diquat on the natural bottom fauna and plankton in two subtropical ponds. Proc. S.E. Assoc. Game Fish Comm., 16 : 301-307.
- Tawfik, M.F.S. (1953). Part I, Aquatic insects in swimming pools; Part II, The external morphology and life-history of the giant water-bug Lethocerus niloticus Stael (Hemiptera : Belostomatidae). M.Sc. Thesis, Fac. Agric., Cairo Univ., 184 pp.
- , and Awadallah, K.T. (1975). The biology of the water scorpion Ranatra vicia Signoret (Heteroptera : Nepidae). Dtsch. Ent. Z., 22 (1-3) : 229-240.
- Thomson, W.T. (1983). "Agricultural Chemicals" Book II, Herbicides. Thomson publications.

- Twinn, C.R.; Brown, A.W.A. and Hurtig, H. (1950). Area control of mosquito by air craft in sub-arctic Canada. Proc. N. J. Mosq. Extern. Assoc., p. 113-140.
- Walker, C.R. (1963). Endothall derivatives as aquatic herbicides in fishery habitats. Weeds, 11 : 226-232.
- (1965). Diuron, fenuron, monuron, neburon and TCA mixtures as aquatic herbicides in fish habitats. Weeds, 13 : 297-301.
- Walker, T.W.; Meek, C.L.; Wright, V.L. and Billodeaux (1985). Susceptibility of Romanomermis culicivorax (Nematode : Mermithidae) postparasites to agrichemicals used in Louisiana rice production. J. Amer. Mosq. Contr. Assoc., 1 (4) : 477.
- Warnick, S.L.; Gaufin, R.F. and Gaufin, A.R. (1966). Concentration and effects of pesticides on aquatic environments. J. Amer. Water Works Assoc., 58 : 601-608.
- Washino, R.K. (1969). Progress in biological control of mosquitoes invertebrate and vertebrate predators. Proc. Pap. 37th Ann. Conf. Calif. Mosq. Contr. Assoc., 37 : 16-19.
- and Kerwin, J.L. (1983). Establishment and overwintering of postparasitic Romanomermis culicivorax in rice fields. Ann. Rep. Univ. Calif. Mosq. Contr. Res., p.66-67.
- Webster, J.M. (1973). Manipulation of environment to facilitate use of nematodes in biocontrol of insects. Exp. Parasitol., 33 : 197-200.
- Weiser, J. (1966). NemociHmyzu, Academia, Prague, 554.

Welch, H.E. (1971). Various target species : attempts with DD-136. In : Biological control programmes against insect and weeds in Canada, 1959-1968. Tech. Comm., CIBC, 4 : 62-66.

----- and Bronskill, J.F. (1962). Parasitism of mosquito larvae by the nematode, DD-136 (Nematode : Neoplectanidae). Can. J. Zool., 40 : 1263.

Whitney, E.W. (1968). The toxicity of Silvex to aquatic fauna. Weed Soc. Amer. Abstr., 1968 : 63.

Winner, R.A. and Steelman, C.D. (1978). Effect of selected insecticides on Romanomermis culicivorax, a mermithid nematode parasite of mosquito larvae. Mosq. News, 38 : 546-553.

Woodard, D.B. and Fukuda, T. (1977). Laboratory resistance of the mosquito, Anopheles quadrimaculatus to the mermithid nematode Diximermis peterseni. Mosq. News, 73 (2) : 192-195.