

REFERENCES

- Acherman, L.V.; Sjut, H.J. and Abell, R.M. (1976):**
Osteoarthrosis (Ch. 8), Bones and Joints. The William and Wilkins Company, P. 308.
- Ahlgrant, T.; Berghem, L. and Jarstrand, T. (1985):**
Plasma fibronectin decrease during septicaemia. Scand. J. Infect. Dis., Vol. (17) (1), P. 107.
- Aitchenson, C.T.; Pebles, C.; Joslin, F. and Tan, E.M. (1982):**
Characteristics of antinuclear antibodies in rheumatoid arthritis. Arth. Rheum., Vol. 23, P. 528.
- Akiyama, S.K. and Yamada, K.M. (1983):**
Fibronectin in connective tissue diseases. (Eds.) : Wagner, B.M., Fleischmajor, R. Published by William and Wilkins. P. 55-96.
- Ali, I.U. and Hynes, R.O. (1978):**
Effects of LETS glycoprotein on cell motility. Cell, Vol. 14, P. 439-440.
- Armstrong, C. and Mow, V. (1982):**
Variation in the intrinsic mechanical properties of human articular cartilage with age, degeneration and water content. J. Bone Joint Surg., Vol. 64 A., P.88-94.
- Arnett, F.C.; Edworthy, S. and Block, D.A. (1988):**
The revised ARA criteria for classification of R.A. Arth. Rheum. 31: 315.
- Balian, G.; Chick, E. and Crouch, E. (1979):**
Isolation of a collagen binding fragment from fibronectin and cold insoluble globulin. J. Biol. Chem., Vol. 25 (4), P. 1429.

Ball, J. and Sharp, J. (1978):

Osteoarthritis, quoted from Copeman's Textbook of Rheumatic disease, 5th edition. by J.T. Scott (Editor), Churchill Livingstone, Edinburgh, London and New York, P. 595.

Barnes, G.C. and Mason, M. (1975):

Rheumatoid arthritis. In "An introduction to clinical rheumatology". Ed. by Mason, M. and Currey, H.L. 2nd. Ed., P. 24, Pitman.

Bayliss, M.T. and Venn, M. (1980):

Chemistry of human articular cartilage in: Maroudas, A.; Holborow, E.J., eds. Studies in Joint disease. Vol. 1. Tunbridge Wells: Pitman Medical, P. 2-58.

Bennett, J.C. (1981):

The aetiology of rheumatoid arthritis. In "Textbook of rheumatology". Edited by Kelly, W.N.; Harris, E.D.; Ruddy, S. and Sledge, C.B. 1st. Ed. P. 887-895. Philadelphia, London, Toronto, Saunders.

Bierther, M.F. and Wegner, K.W. (1971):

Elektronenmikroskopische untersuchungen synovialer Gefasueranderungen bie chronischer polyarthritits. Zeitsschrift fur Rheumaforschung: 30: 219-220. Quoted after Gardner, D.L. (1978) in (Copeman's Textbook of Rheumatic diseases).

Blake, D.R.; Hall, N.D.; Bacon, P.A. and Dieppe, P.A. (1981):

Articular cartilage destruction in rheumatoid disease Lancet; II : 1142.

Blumenstock, F.A.; Saba, T.M.; Weber, P. and Laffin, R. (1978):

Biochemical and immunological characterization of human opsonic alpha2-SB glycoprotein. Its identity with cold insoluble globulin. J. Biol. Chem., 253: 4287-91.

Bolarin and Adnuga (1986):

Plasma fibronectin in Sick cell disease.
Acta. Haematol., 75 (4): 209.

Boughton, B. and Simpson, A. (1982):

Plasma fibronectin in acute leukaemia.
Br. J. Haematol. Jul., 51 (3): 487.

Brewerton, D.A. (1957):

Hand deformities in rheumatoid disease.
Ann. Rheum. Dis., 16: 183.

Broom, N.D. (1982):

Abnormal softening in articular cartilage: its
relationship to the collagen framework.
Arth. Rheum., 25: 1209-1216.

**Brown, R.A.; Tomlinson, I.W.; Hill, C.R.; Weiss, J.B.;
Phillips, P. and Kumar, S. (1983):**

Relationship of angiogenesis factor in synovial fluid
to various joint diseases.
Ann. Rheum. Dis., 42: 301.

Buchanan, W.W. (1978):

Clinical feature of rheumatoid arthritis. In
"Copeman's textbook of the rheumatic diseases". Ed.
by Scott, J.J.; 5th ed., Livingstone, London and New
York, P. 318.

Buchanan, W.W. and Kean, W.F. (1986):

Articular and systemic manifestations of rheumatoid
arthritis. In "Copeman's textbook of the rheumatic
diseases", ed. by Scott, J.T., 6th.edition, Churchill
livingstone, London and New York, P. 653-705.

Burry, H.C. (1972):

Reduced glomerular function in rheumatoid arthritis.
Ann. Rheum. Dis. : 31: 65.

Caplan, A.; Payne, R.B. and Withey, J.L. (1962):

A broader concept of caplan's syndrome related to rheumatoid factors.
Thorax; 17: 205.

Carney, S.L.; Bilingham, M.E.; Muir, H. and Sandy, J.D. (1984):

Structure of newly synthesized (35-S) proteoglycans and (35-S) proteoglycans turnover products of cartilage explant cultures from dogs with experimental osteoarthritis.
J. Orthop. Res., 3: 140-147.

Carnemolla, B.; Cutolo, M.S.; Castellani, P.; Raffanti, S. and Zardi, L. (1984):

Characterization of synovial fluid fibronectin from patients with rheumatic inflammatory diseases and healthy subjects.
Arthr. Rheum., Vol. 27, No. 8, P. 913-920.

Carsons, S.; Mosesson, M.W. and Diamond, H.S. (1981):

Detection and quantitation of fibronectin in synovial fluid from patients with rheumatic disease.
Arthr. Rheum., Vol. 24, No. 10, P. 1261-1266.

Carsons, S.; Natarajan, C.; Drew, H. and Diamond, H. (1980):

Fibronectin in human synovial fluid.
Arthr. Rheum., Abstracts, Vol. (23), P. 661.

Carsons, S.; Parenti, D.; Laviates, B.B. and Diamond, H. (1985):

Plasma fibronectin in systemic lupus erythematosus, relationship to clinical activity.
J. Rheumatol. Dec., 12 (6): 1088-1092.

Chamberlain, M.A. and Bruckner, F.E. (1970):

Rheumatoid neuropathy.
Ann. Rheum. Dis., 29: 609.

Choate, M.; Mosher, M. and Jennifer, J. (1983):

Fibronectin concentration in plasma of patients with breast cancer, colon cancer and acute leukaemia.
Cancer, 51: 1142.

Clark, R.A.; Lanigan, J. and Dellapelle, P. (1982):

Fibronectin and fibrin provide a provisional matrix for epidermal cell migration during wound re-epithelialization.

J. of Investigative Dermatology, 79:264.

Clark, R.A.; Wilner, N.E. and Norris, D.A. (1988):

Chemotactic activity of fibronectin for human monocytes resides in the 120 K-D fibroblastic cell binding fragments.

J. Biol. Chem., Vol. 263, P. 12115-23.

Clemmensen, I. and Andersen, R.B. (1978):

Properties of fibrinogen antigenic material on the rheumatoid synovial membrane and in the rheumatoid synovial fluid.

J. Lab. Clin. Med., Vol.92, P.678-89.

Clemmensen, I. and Andersen, R.B. (1982):

Different molecular forms of fibronectin in rheumatoid synovial fluid.

Arthr. Rheum., Vol. 25, P. 25-31.

Clemmensen, I.; Holund, B. and Andersen, R.B. (1983):

Fibrin and fibronectin in rheumatoid synovial membrane and rheumatoid synovial fluid.

Arthr. Rheum., Vol. 26, No. 4, P. 479-84.

Cohen, A.S.; Brandt, K.D. and Krey, P. (1975):

Synovial fluid: In laboratory diagnostic procedures in the rheumatic diseases edited by Cohen, A.S. 2nd ed., Boston, Little, Brown and Co., P. 1-62.

Collins, W.J. and Jayson, M.I. (1972):

Measurement of pedal pressures. An illustration of a method.

Ann. Rheum. Dis., 31: 215.

Condemi, J.; Barnett, E.V. and Atwater, E.C. (1965):

The significance of anti-nuclear factors in rheumatoid arthritis.

Arth. Rheum., 8: 1080.

Cooper, N.S. (1986):

Pathology of rheumatoid arthritis.
Med. Clin. North. American, 52: 3.

Cosio, F.G.; Eddy, A.; Monster, M.I. and Bergstein, J.M. (1985):

Decreased plasma fibronectin levels in children with haemolytic uraemic syndrome.
Am. J. Med., 78:549-554.

Dacie, J.V. and Lewis, S.M. (1975):

Practical haematology, 3rd edition, New York, Grune and Stratton Inc.

De Jard, A.; Anderson, T. and Gluide, F. (1986):

The influence of insulin on the raised plasma fibronectin concentration in human obesity.
Acta. Med. Scand., 220 (3): 269.

Dessau, W.; Sasse, J.; Timpl, R. and Von der Mark, K. (1978):

Synthesis and extracellular deposition of fibronectin in chondrocyte cultures.
J. Cell. Biol., Vol. (79), P. 342-355.

Dixon, A.J. (1971):

The rheumatoid foot. In: "Modern trends in Rheumatology". Ed. by Hill, A.G. 2nd ed., P. 158-173.
London: Butterworths.

Domula, M.; Bykowska, K.; Lopaciuk, C.S. and Kopic, M. (1985):

Plasma fibronectin concentration in healthy and septic infants.
J. Pediatr., 144 (1): 49.

Duthie, R.B. and Bentley, G. (1983):

Murcer's orthopaedic surgery. Affection of the knee joint, 8th ed., Ch. 16.

Dutu, A.; Vlaicu-Rus, V.; Bolosiu, H.D. and Cristea, A. (1986):

Fibronectin in plasma and synovial fluid of patients with rheumatic diseases.
Med. Interne., Vol. 24 (1), P. 61-8.

Evans, C.H.; Mears, D.C. and McKnight, J.L. (1981):

A preliminary survey of the wear particles in human synovial fluid.
Arth. Rheum., 24: 912.

Fisher, R.A. (1964):

"Statistical methods of research workers".
10th ed., Edinburgh, Oliver and Boyd.

Forman, M.D.; Malamet, R. and Caplan, D. (1983):

A survey of osteoarthritis of the knee in the elderly
J. Rheum. 10: 282.

Fukuda, M. and Hakomori, S. (1979):

Carbohydrate structure of galactoprotein A.
J. Biol. Chem., 254: 5451.

Fye, K.H.; Moustsopoulos, H.M. and Talal, N. (1982):

Rheumatic diseases. In "Basic and clinical immunology"
Ed. by Fudenberg, H.; Stites, D.; Wells, J. 3rd ed.,
P. 422. California.

Fyrand, O.; Munthe, E. and Solum, N.O. (1978):

Studies on cold insoluble globulin..
Ann. Rheum. Dis., Vol. 37, P. 347-50.

Gahamberg, C.G.; Kiehn, D. and Hakomori, S. (1974):

Change in a surface labelled galactoprotein and in glycolipid concentration in cells transformed by a temperature-sensitive virus mutant.
Nature, 248:413.

Gardner, D.L. (1986):

Pathology of rheumatoid arthritis. In "Copeman's textbook of the rheumatic diseases", ed. by Scott, J.T., 6th edition, Churchill Livingstone London and New York, P. 604-652.

**Ginsberg, M.H.; Painter, R.G.; Forsyth, J.; Birdwell, C.
and Plow, E.F. (1980):**

Thrombin increases expression of fibronectin antigen
on the platelet surface.
Proc. Natl. Acad. Sci. USA., 77: 1049-1053.

**Goebel, K.M.; Storck, U.; Kruger, K. and Schattenkircher,
M. (1984):**

Different synovial fluid fibronectin level in
rheumatoid variants.
Klin Wochenschr., Vol. 62, No. 16, P. 768-72.

Goffin, Y.A.; Thoua, Y. and Potvliege, P.R. (1981):

Microdeposition of amyloid in the joints.
Ann. Rheum. Dis., 40: 27.

Golding, D.N. (1989):

Rheumatoid arthritis. In "Synopsis of Rheumatic
diseases". 5th ed., Wright, P.S.G., Bristol, London,
Boston, P. 52.

Gray, R.G. and Poppo, M.J. (1983):

Necrotising vasculitis as the initial manifestation
of rheumatoid arthritis.
J. Rheumatol., 10: 326.

Gressner, A.M. and Wallraff (1981):

Laser nephelometric quantitation of fibronectin in
human cerebrospinal fluid and seminal plasma.
Clin. Chem. Acta., 113: 207-214.

Grinnell, F. (1976):

The serum dependence of baby hamster kidney cell
attachment to substratum.
Exp. Cell. Res., 97: 265.

Grinnell, F. (1984):

Fibronectin and wound healing.
J. Cell Biochem., Vol. 26, P. 107-116.

Gutman, A.; Yamada, K.M. and Kornbihtt, A. (1986):

Human fibronectin is synthesis as a prepolypeptide.
Febs. Lett. 207 (1): 145.

Hamandberg, G.A.; Kramer, B.J.; Grant, D. and Eisenstein, R. (1986):

Heparin-binding fragments of fibronectin are potent inhibitors of endothelial cell growth.
Biochim. Biophys. Acta. Vol. 874, P. 61-71.

Hardwicke, J. and Squire, J.R. (1952):

The basis of the erythrocyte sedimentation rate.
Clin. Sci., 2: 333.

Harris, E.D. (1981a):

Pathogenesis of rheumatoid arthritis. In "textbook of rheumatology". edited by Kelly, W.N.; Harris, E.D.; Ruddy, S. and Sledge, C.B. 1st ed., P. 896-927. Philadelphia, London, Toronto, Saunders.

Harris, E.D. (1981b):

R.A. the clinical spectrum. In "textbook of Rheumatology" by Kelly, W.N.; Harris, E.D.; Ruddy, S.; and Sledge, C.B., 1st ed., P. 928-963 Philadelphia, London, Toronto, Saunders.

Hastings, D.E. and Evans, J.A. (1975):

Rheumatoid wrist deformities and their relation to ulnar drift.
Journal of Bone and Joint Surgery; 57A: 930.

Hayes, J.S. and Stinson, I.N. (1976):

Erythrocyte sedimentation rate and age.
Arch. Ophthalmol., 94: 939.

Heflet, A.J. (1974):

Disorders of the knee. J.B. Lippincott Company.
Philadelphia, Toronto, P. 165.

Hess, L.W.; William, F.O.; Jerry, A.H.; Craig, A.W. and Hemming, V.G. (1986):

Plasma and amniotic fluid concentrations of fibronectin during normal pregnancy.
Obstetrics and Gynaecology, 68 (1): 25-28.

Hickling, D. and Golding, J.R. (1984):

An outline of rheumatology. 1st edition,
P. 48-75. Bristol, John Wright and Sons.

Hofler, H. and Kingemann, H. (1984):

Fibronectin and factor VIII related antigen in liver
cirrhosis and acute liver failure.
J. Clin. Chem. Clin. Bioch., 22 (1): 14.

Holt, P.J. (1983):

Rheumatoid arthritis and related diseases. In
"Postgraduate textbook of clinical orthopedics". Ed.
by Harris, N.H. 3rd ed., P. 443-455. Bristol: John
Wright and Sons.

Hough, A.J. and Sokoloff, L. (1985):

Pathology. In "Rheumatoid arthritis, aetiology,
diagnosis and management" ed. by Utsinger, P.D.;
Zuawfler, N.J., P. 49-69, J.B. Lippincott,
Philadelphia.

Hughes, G.R. and Currey, H.L. (1975):

Laboratory investigations. In "An introduction to
clinical rheumatology". Ed. by Mason, M. and Currey,
H.L. 2nd ed., P. 348, Philadelphia.

Hughes, R.D. and William, R. (1987):

Fibronectin and liver disease.
Hepatology, 7: 800-801.

Humphries, M.J. and Ayad, S.R. (1983):

Stimulation of DNA synthesis by Cathepsin D digests
of fibronectin.
Nature, Vol. 305, P. 811-813.

Huskinson, E.C. (1982):

Drugs and rheumatoid disease. In "Drug treatment of
the Rheumatic diseases". Ed. by Hart, D.F.A. Press.
New York, Tokyo, P. 61.

Hynes, R.O. (1985):

Molecular biology of fibronectin.
Ann. Rev. Cell. Bio. 1: 67.

Hynes, R.O. (1986):

Fibronectins.
Scientific American, 254: 32-42.

Hynes, R.O. and Bye, J.M. (1974):

Density and cell cycle dependence of cell surface protein in hamster fibroblasts.
Cell, 5: 113.

Hynes, R.O. and Destree, A. (1977):

Extensive disulfide bonding at the mammalian cell surface.
Proc. Natl. Acad. Sci. USA. 74: 2855.

Jayson, M.I. and Dixon, A.S. (1970):

Valvular mechanisms in juxta. articular cysts.
Ann. Rheum. Dis., 29: 415.

Jitoku, M.; Koide, N. and Nagashima, H. (1986):

Decreased plasma fibronectin in liver diseases correlated to the severity of fibrotic, inflammatory and neurotic changes of liver tissues.
Acta. Med. Scandinavica, 404: 189.

Johansson, S.; Rubin, K.; Ahlgren, T. and Selijeld, R. (1979):

In vitro biosynthesis of CIG fibronectin by mouse peritoneal macrophages.
FEBS. LETT, 105: 313.

Jones, K.L.; Brown, M.; Ali, S.Y. and Brown, R.A. (1987):

An immunohistochemical study of fibronectin in human osteoarthritic and disease free articular cartilage.
Annals of the Rheumatic diseases.
Vol. 46, P.809-815.

Kay, J.; Austen, K.F. and Czop, J.K. (1991):

Identification and characterization of opsonic fibronectin in synovial fluids of patients with active rheumatoid arthritis.
Arthr. Rheum., Vol. 34, No. 6, P. 687-95.

Logue, G. (1976):

Felty's syndrome: granulocyte-bound immunoglobulin G and splenectomy.
Ann. Int. Med., 85: 437.

Lu-Steffes, M.; Iammartino, A.J.; Schmid, F.R.; Castor, C.W. and Anderson, B. (1982):

Fibronectin in rheumatoid and non-rheumatoid arthritic synovial fluids and in synovial fluid cryoproteins.
Ann. Clin. Lab. Sci., Vol. 12 (3), P. 178-185.

MacFarlane, J.D.; Franken, C.K. and Van Leeuwen, A.W. (1984):

Progressive cavitating pulmonary changes in rheumatoid arthritis: a case report.
Ann. Rheum. Dis., 43: 98.

MacFarlane, J.D. and Van Der Linden, S.J. (1981):

Leaking rheumatoid olecranon bursitis as a cause of forearm swelling.
Ann. Rheum. Dis., 40: 309-311.

Mackay, J.M.; Sim, A.K. and Gardner, D.L. (1983):

Aetiology of rheumatoid arthritis, 42: 443-447.

Magyar, E.; Talerman, A. and Wouters, H.W. (1973):

Histological abnormalities in muscle spindles in rheumatoid arthritis.
Ann. Rheum. Dis., 32: 143.

Mallya, R.D. and Mace, B.E. (1981):

The assessment of disease activity in rheumatoid arthritis using a multivariate analysis.
Rheumatism and Rehabilitation, 20: 14-17.

Mancini, G.; Carbonara, A.O. and Hereman, J.F. (1965):

Immunological quantitation of antigens by single radial immunodiffusion.
Int. J. Immunochem., 2: 235-254.

Maroudas, A. (1970):

Distribution and diffusion of solutes in articular cartilage.
Biophys. J. Vol. (10), P. 365-79.

Maroudas, A. (1976):

Balance between swelling pressure and collagen tension in normal and degenerated cartilage.
Nature, 260: 808.

Mastuda, M.; Yamanaka, K. and Mastuda, A. (1982):

Distribution of fibronectin in plasma and liver diseases.
Clin. Chem. Acta., 118: 141.

Matsuyama, T.; Yamada, A.; Kay, J.; Yamada, K.M. and Morimoto, C. (1989):

Activation of CD4 cells by fibronectin and anti-CD3 antibody.
J. Exp. Med., Vol. 170, P. 1133-48.

McCafferty, M.; Lepow, M.; Saba, T.; Cho, E. and White, J. (1983):

Normal fibronectin level as a function of age in the pediatric population.
Ped. Res. Jour., 17 (6): 482.

McCarty, D.J. (1975):

Calcium pyrophosphate dihydrate deposition disease.
In Holt, P.J. (ed.): Current Topics in connective tissue disease. New York, Churchill Livingstone, P. 181-197.

McCarty, D.J. (1979):

Clinical and laboratory findings in osteoarthritis.
In "Arthritis and Allied conditions", 9th ed., edited by D.J. Macarty, J.L. Hollander. Lea, Febiger, Philadelphia, P. 1168.

McDonald, J.A.; Bruce, J.B.; Rosenberg, D. and Crystal, R.G. (1979):

Destruction of a major extracellular fibronectin of human fibroblasts by neutral proteases from polymorphonuclear leucocyte granules.
Lab. Invest., Vol. 40, P. 350.

Menzel, E. and Dunky, A. (1986):

Plasma fibronectin in the aged.
J. Gerontol., 19 (3): 152-157.

Moore, C.P. and Willkens, R.F. (1977):

The subcutaneous nodules : its significance in the
diagnosis of rheumatic disease.
Seminars in Arth. Rheum., 7: 63.

Morrison, P.R.; Edsall, J.T. and Miller, S.G. (1948):

The separation of purified fibrinogen from fraction I
of human plasma.
J. Am. Chem. Soc., 70: 3103.

Mosesson, M.W. and Amrani, D.L. (1980):

The structure and biological activities of plasma
fibronectin.
J. American Society of Haematology, 56 (2): 145-158.

Mosesson, M.W.; Chen, A.B. and Huseby, R. (1975):

The cold-insoluble globulin of human plasma, studies
of its essential structural feature.
Biochem. Biophys. Acta., 386: 509.

Mosher, D.F. (1980):

Fibronectin.
Prog. Hemostasis Thromb., 5: 111-151.

Mosher, D.F. (1984):

Physiology of fibronectin.
Ann. Rev. Med., 35: 561.

Mosher, D.F. and Williams, E.M. (1978):

Fibronectin concentration is decreased in plasma of
severely ill patients with disseminated intravascular
coagulation.
J. Lab. Clin. Med., 91: 729-735.

Moskowitz, R.W. (1979):

In: "Arthritis and Allied conditions". 9th ed.,
edited by D.J. McCarty, J.L. Hollander. Lea, Febiger,
Philadelphia. P. 1161.

Moskowitz, R.W. and Goldberg, V.M. (1988):

Osteoarthritis. In "Primer on the Rheumatic diseases", 9th ed., by R. Schumacher, J.H. Klippel, D.R. Robinson (eds.). Arthritis foundation, Atlanta GA, P. 171-177.

Myers, D. and Broon, N.D. (1982):

Morphological and biomechanical studies of rheumatoid pannus and cartilage.
J. Rheumatol. 9: 502.

Nakata, H. and Russell, W. (1970):

Chest roentgenograms in rheumatoid arthritis.
American Journal of Roentgenology, 108: 819.

Narfolk, D.; Brown, M. and Roberts, B. (1983):

Plasma fibronectin in myeloproliferative disorders and chronic granulocytic leukaemia.
Br. J. Haematol., Oct., 55 (2): 319.

Nichols, P.J.R. (1976):

Rehabilitation medicine, the management of physical disabilities, Butter Worth, London, Boston, P. 149.

Norris, D.A.; Clark, R.A.; Swigart, L.M. and Howell, S.E. (1982):

Fibronectin fragments are chemotactic for human peripheral blood monocytes.
J. Immunol. Vol. 129, P. 1612-18.

Oh, E.; Pierschbacher, M. and Ruoslahti, E. (1981):

Deposition of plasma fibronectin in tissues.
Proc. Natl. Acad. Sci. USA., 78: 3218.

Oishi, M.; Yamamoto, M.; Yoshida, T.; Sade, M. and Matsumoto, K. (1986):

Opsonic activity of plasma fibronectin for staph. aureus by human alveolar macrophages.
J. Exp. Med., 149 (1): 112.

Omar, A. and Mowat, A.G. (1968):

Nature of anaemia in rheumatoid arthritis. IX folate metabolism in patients with rheumatoid arthritis. Ann. Rheum. Dis., 27: 414.

Panayi, G.S. (1986):

Aetiopathogenesis of rheumatoid arthritis. In "Copeman's textbook of the rheumatic diseases". ed. by Scott, J.T., 6th edition, Churchill livingstone, London, New York, P. 595-603.

Pearlstein, E. (1976):

Plasma glycoprotein which mediates adhesion of fibroblasts to collagen. Nature, 62: 497.

Pearlstein, E. and Gold, L.I. (1980):

Fibronectin: a review of its structure and biological activity. Mol. Cell. Biochem., 29: 103.

Pearson, C.M.; Paulus, H.E. and Machleder, H.I. (1975):

The role of the lymphocyte and its products in the propagation of joint disease. Ann. N.Y. Acad. Sci., 256: 150.

Perri, R.; K-e, N. and McCarthy, J. (1982):

Fibronectin enhances in vitro monocyte mediated tumouricidal activity. Blood, 60 (2): 430.

Pettipher, E.R.; Higgs, G.A. and Henderson, B. (1986):

Interleukin-1 induce leucocyte infiltration and cartilage proteoglycan degradation in the synovial joint. Proc. Natl. Acad. Sci., USA., 83.

Phan, T.M.; Foster, C.S.; Boruchoff, S.A. and Colvin, R.B. (1987):

Topical fibronectin in the treatment of persistent corneal epithelial defects and trophic ulcer. Am. J. Ophthalmol., 104 (5): 494-501.

Phillips, R.S. (1972):

Venous pathology in osteoarthrosis of the knee.
J. R. Col. Surg., Edinb., 17: 195.

Pickup, M.E.; Dixon, J.S.; Hallett, G.; Bird, H.A. and Wright, V. (1981):

Plasma viscosity. A new appraisal of its use as index of disease activity in R.A.,
Ann. Rheum. Dis., 40:272.

Plow, E. and Ginsberg, M. (1981):

Specific and saturable binding of plasma fibronectin to thrombin stimulated human platelets.
J. Biologic and Chem., 256: 9477.

Pommier, C.; Jnada, S.; Fries, L.; Frank, M. and Brown, E. (1983):

Plasma fibronectin enhance phagocytosis.
J. Exp. Med., 157: 1844.

Pope, R.M.; Mannik, M.; Gilli, B.G. and Teller, D.C. (1975):

The hyperviscosity syndrome in R.A.,
Arth. Rheum., 18: 97.

Pussell, B.A.; Peake, P.W.; Brown, M.A. and Charlesworth, J.A. (1985):

Human fibronectin metabolism.
J. Clin. Invest., Vol. 76, P. 143-148.

Radin, E.L.; Paul, I.L. and Rose, R.M. (1972):

Role of mechanical factors on pathogenesis of primary osteoarthritis.
Lancet, 11: 519.

Rajarman, R.; MacSween, J.M. and Murdock, C.A. (1983):

Exogenous fibronectin requirement for adhesion by neoplastic human cells. Exp. Cell. Biol., 51: 9-18.

Rees, J.A.; Ali, S.Y. and Brown, R.A. (1987):

Ultrastructural localisation of fibronectin in human osteoarthritic articular cartilage.
Ann. Rheum. Dis., Vol. 46, P. 816-22.

Rothermich, N.O. and Whisler, R.L. (1985):

Theories of causation and pathogenesis of rheumatoid synovitis. In "Rheumatoid arthritis". P. 3-23, Grune, Stratton, Inc. New York.

Ruoslahti, E.; Engvall, E. and Hayman, E.G. (1981):

Fibronectin current concepts of its structure and functions.
Coll. Res., 1: 95-128.

Ruoslahti, E. and Vaheri, A. (1974):

Novel human serum protein from fibroblast.
Nature, Vol. 248, P. 789-791.

Saba, T.M. (1970):

Physiology and physiopathology of reticuloendothelial system.
Arch. Int. Med., 126: 1031.

Saba, T.M.; Blumenstock, F.; Shah, D.M.; Hrinda, M.E. and Deno, D.C. (1986):

Reversal of opsonic deficiency in surgical trauma and burn by infusion of human plasma fibronectin.
Am. J. Med., 80 (2): 129.

Saba, T.M.; Gregory, T.J. and Blumenstock, F. (1980):

Human host defense mechanisms as they relate to surgery and trauma.
Surg. Annu., Vol. 21, P. 1-80.

Saba, T.M. and Jaffe, E. (1980):

Plasma fibronectin : its synthesis by vascular endothelial cells and role in cardio-pulmonary integrity after trauma as related to reticuloendothelial function.
Am. J. Med., 68: 577-594.

Saba, T.M.; Kiener, J. and Holman, J. (1982):

Fibronectin and critically ill patients.
Intensive care Medicine, 12 (5): 350.

Samson, D.; Holliday, D. and Gumpel, J.M. (1977):

Role of ineffective erythropoiesis in the anaemia of rheumatoid arthritis.
Ann. Rheum. Dis., 36: 181.

Saxne, T.; Heinegard, D.; Wollheim, F.A. (1986):

Therapeutic effects on cartilage metabolism in arthritis as measured by release of proteoglycan structures into the synovial fluid.
Ann. Rheum. Dis., 45: 491-497.

Scholmerich, J.; Volk, B.A.; Kottges, E.; Ehlers, S. and Gerok, W. (1984):

Fibronectin concentration in differentiation between malignant and non-malignant ascitis.
Gastroenterology, 87 (3): 160-164.

Schumacher, H.R.; Klippel, J.H. and Robinson, D.R. (1988):

9th. ed. by Schumacher, H.R.; Klippel, J.H. and Robinson, D.R. Arthritis Foundation, Atlanta G.A., P. 171.

Scott, D.L.; Delamere, J.P. and Walton, K.W. (1981):

The distribution of fibronectin in the pannus, in rheumatoid arthritis.
Br. J. Exp. Pathol., Vol. 62, P. 362-368.

Scott, D.L.; Farr, M.; Crockson, A.P. and Walton, K.W. (1982):

Synovial fluid and plasma fibronectin levels in rheumatoid arthritis.
Clin. Sci., Vol.62 (1), P.71-76.

Scott, D.L.; Grindulis, K.A.; Coulton, B.L.; Popert, A.I. and Bacon, P.A. (1984):

Progression of radiological changes in rheumatoid arthritis.
Ann. Rheum. Dis., 43: 8-17.

Scott, D.L.; Wainwright, A.C.; Walton, K.W. and Williamson, N. (1981):

Significance of fibronectin in rheumatoid arthritis and osteoarthritis.
Ann. Rheum. Dis., 40: 142-153.

Sekiguchi, K. and Hakomori, S. (1983):

Domain structure of human plasma fibronectin: differences and similarities between human and hamster fibronectins.

J. Biol. Chem., Vol. 258, P. 3967-73.

Shepeard, H. (1983):

An update on mechanisms of cartilage destruction in rheumatoid arthritis.

Australian and New Zealand J. Med., 13: 195.

Shiozawa, K.; Shiozawa, S.; Shimizu, S. and Fujita, T. (1984):

Fibronectin on the surface of articular cartilage in rheumatoid arthritis.

Arthr. Rheum., Vol. 27, No. 6, P. 615-621.

Shiozawa, S. and Ziff, M. (1983):

Immunoelectron microscopic demonstration of fibronectin in rheumatoid pannus and at the cartilage pannus junction.

Ann. Rheum. Dis., Vol. 42, P. 254-263

Sigal, L.H.; Steere, A.C. and Niderman, J.C. (1983):

Symmetrical polyarthrititis associated with heterophile negative infectious mononucleosis.

Arth. Rheum., 26: 553-556.

Silberbeg, R. (1985):

Diseases of joints. In "Anderson's pathology" edit. Kisane, J.M., Vol. II, 8th ed., P. 1828-1863, The C.V. Mosby Comp., St. Louis.

Siri, A.; Carnemolla, B. and Raffanti, S. (1984):

Fibronectin concentration in pleural effusion of patients with malignant and non-malignant diseases. Cancer Lett., 22: 1-8.

Sissons, H.A. (1979):

Rheumatoid arthritis-osteoarthritis. In: "Systemic pathology", 2nd edition, Vol. 5, Churchill Livingstone, Edinburgh, London, P. 2501.

- Snukler, N.M.; Edeiken, J. and Givillians, V.J. (1971):**
Ankylosis in ostoarthritis of the finger joint.
Radiology, 100: 515.
- Sokoloff, L. (1953):**
The heart in rheumatoid arthritis.
Am. Heart J., 45: 635.
- Sokoloff, L. (1979):**
Pathology of rheumatoid arthritis. In "Arthritis and Allied Diseases", Edited by McCarty, D.J. 9th Ed., P. 309-331. Philadelphia: Lea and Febiger.
- Solomon, L. (1972):**
Pathogenesis of osteoarthritis.
Lancet, 1: 1072.
- Stathakis, N.E.; Fountos, A. and Tsianos, E. (1981):**
Plasma fibronectin in normal subjects and in various diseases states.
J. Clin. Pathol. 30: 504-508.
- Steinbroker, O.; Traeger, C.H. and Batterman, R.C. (1949):**
Therapeutic criteria in rheumatoid arthritis.
JAMA, 140: 659-662.
- Stenman, S. and Vaheri, A. (1978):**
Distribution of a major connective tissue protein, fibronectin in normal human tissues.
J. Exp. Med., 147: 1054-1064.
- Stubbs, T.M.; Lazarchick, J. and Horger, E.O. (1984):**
Plasma fibronectin levels in pre-eclampsia.
Am. J. Obstetrics, Gynecology, 150: 885.
- Sukoff, M.H.; Ckadin, M.M. and Moran, T. (1972):**
Decompression for myelopathy caused by rheumatoid arthritis of the cervical spine.
Journal of Neurosurgery, 37: 493.

Teshima, R.; Treadwell, B.V.; Trachan, C.A. and Mankin, H.J. (1983):

Comparative rates of proteoglycans in normal and osteoarthritic chondrocytes.
Arth. Rheum., 26: 1225-1230.

Tyler, J.A. and Saklatvala, J. (1985):

Pig interleukin-1 induces resorption of cartilage proteoglycan and prevents synthesis of proteoglycan and collagen.
Br. J. Rheumatol., 24 (Suppl. 1): 150-155.

Vercellotti, G.M.; McCarthy, J.; Jacob, H.S. and Moldow, C.F. (1983):

Inflamed fibronectin: An altered fibronectin enhances neutrophil adhesion.
Blood, Vol. 62, P. 1063-69.

Villiger, B.; Broekelmann, T. and Kelly, D. (1981):

Bronchoalveolar fibronectin in smokers and non-smokers.
Am. Rev. Resp. Dis., 124: 652-654.

Vuoto, M.; Wrann, M. and Ruoslahti, E. (1977):

Similarity of fibronectin isolated from human plasma and fibroblast culture medium.
FEBS LETT, 82: 277.

Watson, R.G. (1965):

A more specific method for detecting and quantitating rheumatoid factors using a simple modification of the rheumatoid arthritis test.
Am. J. Clin. Path., 43:152.

Weiss, M.A.; Oai, B.S. and Oai, Y.M. (1979):

Immunofluorescent localization of fibronectin in the human kidney.
Lab. Invest., 41: 340-347.

Weiss, R.E. and Reddi, A.H. (1981):

Appearance of fibronectin during the differentiation of cartilage, bone and bone Marrow.
J. Cell. Biol., Vol. 88, P. 630-636.

Weissmann, G. (1972):

Lysosomal mechanisms of tissue injury in rheumatoid arthritis.
Eng. J. Med., 286: 141.

Weissmann, G.; Pearlstein, E.; Perez, H.D.; Falkow, S. and Hoffstein, S.T. (1983):

Neutrophils synthesize and deposit fibronectin on surfaces to which they attach.
Trans. Assoc. Am. Phys., Vol. 75, P. 4944-8.

Westergren, A. (1926):

Amer. Rev. Tuberc., 14: 94.

Williams, E.; Janney, P. and Peny, J. (1982):

Confirmational studies of fibronectin. Effects of PH, ionic strength and collagen binding.
J. Biol. Chem., 527: 1473.

Wilson, R.; Mammen, E.; Robson, M.; Hessers, J. and Depoli, (1986):

Antithrombin, prekallikrein and fibronectin in surgical patients.
Arch. Surg., 121 (6): 635.

Wisnieski, J.J. and Askari, A.D. (1981):

Rheumatoid nodulosis-a relatively benign rheumatoid variant.
Arch. Int. Med., 141: 615.

Wolf, L.; Timple, R.; Pecker, L. and Steffen, C. (1967):

A two-component system of human serum agglutination gelatin-coated erythrocyte.
Vox. Sang., 12: 443.

Wolley, D.E.; Crossley, M.J. and Evanson, J.M. (1977):

Collagenase at sites of cartilage erosion in the rheumatoid joint.
Arthr. Rheum., Vol. 20, P. 1231-9.

Woods, D.E.; Staus, D.C.; Johanson, W.G. and Bass, J.A. (1981):

Role of fibronectin in the prevention of adherence of pseudomonas aeruginosa to buccal cells.
J. Infect. Dis., 143: 779-784.

Wurster, N.B. and Lust, G. (1984):

Synthesis of fibronectin in normal and osteoarthritic articular cartilage.
Biochim. Biophys. Acta., Vol. 800, P. 52-8.

Wurster, N.B. and Lust, G. (1986):

Incorporation of purified plasma fibronectin into explants of articular cartilage from disease free and osteoarthritic canine joints.
J. Orthop. Res., Vol. (4), P. 409-19.

Yamada, K.M. and Olden, K. (1978):

Fibronectin-adhesive glycoproteins of cell surface and blood.
Nature, 275: 179.

Yamada, K.M. and Weston, J.A. (1974):

Isolation of major cell surface glycoprotein from fibroblasts.
Proc. Natl. Acad. Sci. USA., 71: 3492.

Yamada, K.M.; Yamada, S.S. and Pastan, I. (1976):

Cell surface protein partially restores morphology, adhesiveness and contact inhibition of movement to transformed fibroblasts.
Proc. Natl. Acad. Sci. USA., 73: 1217.

Ziff, M. and Baum, J. (1974):

Laboratory finding in rheumatoid arthritis. In "Arthritis and allied conditions" ed. by Hollander, J. MacCarty, D.J. 8th ed., Lea and Febiger, Philadelphia. P. 367.

Zvaifler, N. (1973):

Immunopathology of joint inflammation in arthritis.
Adv. Immunol., 16: 265.