

REFERENCES

VII. REFERENCES

- Abdel-Kawy, A.M. (1981): Physiochemical effects of entomophagous nematodes on their insect hosts. M.Sc. Thesis, Faculty of Agriculture, Cairo University.
- Abdel-Kawy, A.M. (1985): Biological control of some lepidopterous pests by the nematode *N. carpocapsae* and the physiological changes associated with it. Ph.D. Thesis, Fac. Agric., Cairo University.
- Abdel-Kawy, A.M.; M.H. El-Bishry and T.A.H. El-Kifl (1992): Controlling the leopard moth *Z. pyrina* by three entomopathogenic nematode species in the field. Bull.Fac. Agric., Cairo Univ., 43 (2): 769-780.
- Abdel-Kawy, A.M.; N.M.E. Saleh and A.M. El-Azzazi (1996): Using entomopathogenic nematodes in the control of two apple borers. Egyptian Journal of Biological Pest Control, 6 (2): 147-151.
- Abou El-Magd, A.A. (1995): Is *Nigilla sativa* (black seeds) capable of activating immune responses of *S. gregaria*. Ain Shams Science Bulletin, 33: 203-215.
- Abou El-Magd, A.A. and T.A.H. El-Kifl (1993): Cellular and humoral immune reactions of *S. littoralis* and *A. ipsilon* larvae against the nematode *H. bacteriophora*. J. Egypt, Ger. Soc. Zool., vol. 12 (A), Comparative Physiology, 475-488.
- Abou El-Magd, A.A.; M.F. Harb and M.S. El-Katatny (1991): Effect of fungal spores, fungal soluble proteins and cell wall B-1,3-glucans on the haemocytic aggregation (nodule formation) in *S. gregaria* Forsk.

- (Orthoptera: Acrididae). Proc. Zool. Soc. A.R. Egypt, vol. 22: 53-62.
- Abu Hatab, M; S. Selvan and R. Gaugler (1995):** Role of protease in penetration of insect gut by the entomopathogenic nematode *S. glaseri* (Nematoda: Steinernematidae). Journal of Invertebrate Pathology, 66 (2): 125-130.
- Ahmed, N.M. (1982):** Comparative studies on the effect of pesticides and the entomogenous nematodes in controlling some economic insects. Ph.D. Thesis, Fac. Agric., Cairo University, 157 pp.
- Akhurst, R.J. (1983):** Taxonomic study of *Xenorhabdus*, a genus of bacteria symbiotically associated with insect pathogenic nematodes. Int. J. System. Bact., 33 (1): 38-45.
- Akhurst, R.J. (1986):** Controlling insects in soil with entomopathogenic nematodes, pp. 265-267. In *R.A. Samson; J.M. Vlak, and D. Peters* [Eds.], Fundamental and Applied Aspects of Invertebrate Pathology.
- Andreadis, T.G. and D.W. Hall (1970):** *N. carpocapsae* encapsulation in *A. aegypti* and changes in host haemocytes and haemolymph proteins. Exp. Parasitol., 39: 252-261.
- Anonymous, S. (1987):** SAS user's guide: Statistics. pp. 1028. SAS Institute. Cary, N.C.0
- Baur, M.E.; H.K. Kaya and G.S. Thurston (1995):** Factors affecting entomopathogenic nematode infection of *P. xylostella* on a leaf surface. Entomologia-Experimentalis-et-Applicata, 77 (3): 239-250.

- Bedding, R.A. (1990): Logistic and Strategies for introducing entomopathogenic nematode technology into developing countries, in Entomopathogenic Nematodes in Biological Control (*Gaugler, R. and Kaya, H.K.*, Eds) CRC Press, Boca Raton Florida, USA, pp. 233-246.
- Blackshaw, R.P. and C.R. Newell (1987): Studies on temperature limitations to *H. heliothidis* activity. *Nematologica*, **33**: 180-185.
- Brehelin, M. and D. Zachary (1986): Insect haemocytes: A new classification to rule out the controversy. In: Immunity in Invertebrates, Brehelin, M. (ed.) Springer-Verlag 36-46.
- Brown, I.M. and R. Gaugler (1997): Temperature and humidity influence emergence and survival of entomopathogenic nematodes. *Nematologica*, **43** (5): 363-375.
- Bucher, G.E. (1960): Potential bacterial pathogens of insects and their characteristics. *J. Insect Pathol.*, **2**: 172-195.
- Cabanillas, H.E. and J.R. Raulston (1994): Pathogenicity of *St. riobravis* against corn earworm, *H. zea* (Boddie). *Fundam. Appl. Nematol.*, **17**: 219-223.
- Campbell, J.F. and R. Gaugler (1994): Entomopathogenic nematode vertical distribution in turfgrass soil. *J. Nematol.*, **26**, 540 pp.

- Chain, B.M. and R.S. Anderson (1982): Selective depletion of the plasmatocytes in *G. mellonella* following infection of bacteria. *J. Insect Physiol.*, 28: 377-384.
- Cheng, C. and R.F. Hou (1997): Effects of environmental factors on survival of the entomopathogenic nematode, *S. carpocapsae*. *Chinese Journal of Entomology*, 17 (2): 120-131.
- Choo, H.Y. and H.K. Kaya (1991): Influence of soil texture and the presence of roots on host finding by *H. bacteriophora*. *J. Invert. Pathol.*, 58: 279-280.
- Couterier, A. (1963): Recherches sur des Mermithidae nematodes parasites du hanneton commun *Me. melolontha*. *Annls Epiphyt.*, 14: 203-267.
- Creighton, C.S.; F.P. Cuthbert and J.R. Reid (1968): Susceptibility of certain Coleopterous larvae to the DD-136 nematode. *J. Invert. Pathol.*, 10: 368-373.
- Danilov, L.G. and F.V. Karpova (1990): Testing entomopathogenic nematodes against Acrididae. *Zashchita-Rastenii*, No. 7, 34-35.
- Dunn, P.E. and D.R. Drake (1983): Fate of bacteria injected into naive and immunized larvae of the tobacco hornworm, *M. sexta*. *J. Invert. Pathol.*, 41: 77-85.
- Dunphy, G.B. and J.M. Webster (1985): Influence of *S. feltiae* (Filipjev), DD-136 strain on the humoral and haemocytic responses of *G. mellonella* larvae to selected bacteria. *Parasitology*, 91 (2): 369-380.

- Dunphy, G.B. and J.M. Webster (1986a): Temperature effects on growth and virulence of *S. carpocapsae* strains and *H. heliothedis*. Journal of Nematology, 18 (2): 270-272.
- Dunphy, G.B. and J.M. Webster (1986b): Effect of the Mexican strain of *S. feltiae* and its associated bacterium *X. nematophilus* on *G. mellonella*. J. of Parasitology, 72 (1): 130-135.
- Dutky, S.R. (1959): Insect microbiology. Adv. Appl. Microbiol., 1: 175-200.
- Dutky, S.R. and S.W. Hough (1955): Note on a parasite nematode from codling moth larvae *Carpocapsae pomonella*. Proc. Helminth. Soc. Wash., 57: 244.
- Dutky, S.R.; J.V. Thompson and G.E. Cantwell (1964): A technique for the mass propagation of the DD-136 nematode. J. Insect Pathol., 6: 417-422.
- Elawad, S.A.; M.S. Abbas and N.G.M. Hague (1996): The establishment, reproduction and pathogenicity of a new species of *Steinernema* from the Sultanate of Oman in *G. mellonella*. Afro-Asian Journal of Nematology, 6 (1): 40-45.
- El-Bishry, M.H. (1989): Studies on the utilization of the entomogenous nematodes in controlling some pests in Egypt. Ph.D. Thesis, Fac. Agric. Cairo University.

- El-Kifl, T. M. (1980): Utilization of the nematode *N. carpocapsae* in the biological control of cotton leafworm *S. littoralis*. M.Sc. Thesis, Fac. Agric., Cairo Univ., 125 pp.
- El-Kifl, T. M. (1984): Factors affecting potentialities of entomogenous nematodes in the biological control of insect pests. Ph.D. Thesis, Fac. Agric., Cairo Univ., 124 pp.
- El-Maasarawy, S.A. (1980): Studies on the haemolymph of *P. ricini* Biosduval, under different conditions. M.Sc. Thesis, Fac. Agric., Cairo University.
- El-Maasarawy, S.A.S.; M.A.K. El-Sheikh and S.M. Ibrahim (1993): Haemocytes changes in Maize worm, *M. loreyi* (Dup.) parasitized by the braconid wasp, *Meteorus gyrator* THUN. Bull. Ent. Soc. Egypt Econ., Ser., 71: 19-31.
- FAO (1994): Formulation Mission Report on Emergency Prevention System (EMPRES) for Desert Locust. Part I, pp. 5-8, by Tsedeke Abate; Laurence Mc Culloch and Greg Spendjian.
- FAO (1996): Commission for controlling the desert locust in the central region. Report of the Work Shop on Evaluation of Spray Equipment Used in Desert Locust Control, 1 pp.
- Finney, D.J. (1952): Probit analysis. 2nd ed., Cambridge University Press, 218 pp.

- Finney, J.R. and G.F. Bennett (1984): *H. heliothidis*: A potential biocontrol agent of agricultural and forest pests in Newfoundland. J. Agric. Entomology, 1: 287-295.
- Fowler, H.G. and C.R. Garcia (1988): Nematodes (Rhabditida: *S. feltiae*) as natural control agents of mole crickets (Orthoptera: Gryllotalpidae): field and laboratory studies. Revista-Brasileira-de-biologia, 48 (4): 789-795.
- Gagen, S.J. and N.A. Ratcliffe (1976): Studies on the in vitro cellular reactions and fate of injected bacteria in *G. mellonella* and *P. brassicae* larvae. J. Invert. Pathol., 28: 17-24.
- Gaugler, R. (1981): Biological control potential of neoaplectanid nematodes. J. Nematol., 13: 241-249.
- Gaugler, R. and G.M. Boush (1979): Nonsusceptibility of rates of the entomogenous nematodes, *N. carpocapsae*. Environ. Entomol., 8, 658.
- Gaugler, R. and H.K. Kaya (1990): "Entomopathogenic Nematodes in Biological Control". CRC Press, Boca Raton Florida, USA.
- Gaugler, R.; T. McGuire and J. Campbell (1989): Genetic variability among strains of entomopathogenic nematode *S. feltiae*. J. Nematol., 21: 247-253.
- Geden, C.J. and R.C. Axtell (1988): Effect of temperature on nematode (*S. feltiae* [Nematoda: Steinernematidae]) Treatment of soil for control of lesser Mealworm (Coleoptera: Tenebrionidae) in Turkey houses. J. Econ. Entomol., 81 (3): 800-803.

- Geden, C.J.; R.C. Axtell and W.M. Brooks (1986): Susceptibility of the house fly, *M. domestica* (Diptera: Muscidae), to the entomogenous nematodes *S. feltiae*, *S. glaseri* (Steinernematidae), and *H. heliothidis* (Heterorhabditidae). J. Medical Entomology, 23 (3): 326-332.
- Geng, C. and P.E. Dunn (1989): Plasmacyte depletion in larvae of *M. sexta* following injection of bacteria. Developmental and Comparative Immunology, 13: 17-23.
- Georgis, R. and R. Gaugler (1991): Predictability in biological control using entomopathogenic nematodes. J. Econ. Entomol., 84: 713-720.
- Ghally, S.E. (1987): The comparative effect of two nematode species *S. feltiae* Filipjev and *H. bacteriophora* Poinar, on caterpillars of *E. kuehniella* Zeller (Lepidoptera: Pyralidae). J. Zool. Soci. Egypt.
- Ghally, S.E. (1988): Pathogenicity of the nematode *S. feltiae* Filipjev in relation to different insect hosts. Journal of the Egyptian Society of Parasitology, 18 (1): 297-304.
- Ghally, S.E.; O.S. El-Din and M.A. Amin (1988): Effects of the parasitic nematodes on total proteins and total lipids of *C. capitata* Wied, (Diptera: Trypetidae). Journal of the Egyptian Society of Parasitology, 18 (2): 619-627.
- Gingrich, R.E. (1964): Acquired humoral immune response of the large milkweed bug, *O. fasciatus* (Dallas), to injected materials. J. Insect Physiol., 10: 179-194.

- Glazer, I. (1991): Ecological considerations for entomopathogenic nematodes activity under sub-optimal conditions. Bulletin-SROP, 14 (7): 28-31.
- Glazer, I. (1992): Survival and Efficacy of *S. carpocapsae* in an exposed environment. Biocontrol Science and Technology, 2: 101-107.
- Glazer, I. and A. Navon (1990): Activity and persistence of entomopathogenic nematodes tested against *H. armigera* (Lepidoptera: Noctuidae). J. Econ. Entomol., 83: 1795-1800.
- Glazer, I. and M. Wysoki (1990): Steinernematid and heterorhabditid nematodes for biological control of the giant looper, *B. selenaria*. Phytoparasitica, 18 (1): 9-16.
- Glazer, I.; N. Liran; G.O.Jr. Poinar and P.H. Smits (1993): Identification and biological activity of newly isolated heterorhabditid populations from Israel. Fundamental and Applied Nematology, 16 (5): 467-472.
- Glinski, Z. and S. Klimont (1987): Invasion of *V. jacobsoni* Oud. and phagocytic activity of haemocytes of the worker honey bee, *A. mellifera* L. Preliminary observations on wintering bees. Annals-Universitatis-Mariae-Curie-Sklodowska-Sectio-DD.-Medicina-Veterinaria, 42: 1-7.
- Gordon, R.; W.J. Condon; W.J. Edgar and S.J. Bahie (1978): Effect of mermithid parasitism on the haemolymph composition of the larval black flies *P. mixtum* /*fuscum* and *S. venustum* (Diptera). Parasitol., 77: 367-374.

- Gorsuch, A.M. (1982): Regulations for the enforcement of the federal insecticide, fungicide and rodenticide Act (FIFRA) exemption from regulation of certain biological control agents. Fed. Regist., 47: 23928-23930.
- Govindarajan, R.; S. Jayaraj and K. Narayanan (1976): Studies on the effect of *B. thurigiensis* (Bacteria) on the castor semi-loopers, *A. Janata* L. (Lepidoptera: Noctuidae). Z. Angew Entomol., 80: 191-200.
- Gray, P.A. (1973): The Encyclopodia of microscopy and microtechnique. Library of Congress Catalogue Card No. 73-164.
- Gray, P.A. and D.T. Johnson (1983): Survival of the nematode *N. carpocapsae* in relation to soil temperature, moisture and time. J. Georgia Entomol. Soc., 18 (4): 452-460.
- Grewal, P.S.; S. Selvan and R. Gaugler (1994): Thermal adaptation of entomopathogenic nematodes: Niche breadth for infection, establishment and reproduction. J. Therm. Biol., 19 (4): 245-253.
- Griffin, C.T.; J. Moore and M.J. Doowns (1991): Occurrence of insect parasitic nematodes (*Steinernematidae* & *Heterorhabditidae*) in the Republic Ireland. Nematologica, 37: 92-100.
- Griffin, C.T.; W.R. Simons and P.H. Smits (1989): Activity and infectivity of four isolates of *Heterorhabditis* sp. J. Invert. Pathol., 53: 107-113.

- Gunnarsson, S.G.S. (1988): Effects in vivo of B,1-3, glucans from fungal cell walls on the desert locust, *S. gregaria*. J. Insect Physiol., 43 (1): 47-51.
- Gunnarsson, S.G.S. and A.M. Lackie (1985): Haemocytic aggregation in *S. gregaria* and *P. americana* as a response to injected substances of microbial origin. J. Invert. Pathol., 46: 312-319.
- Hara, A.H.; R. Gaugler; H.K. Kaya and I.M. Lebeck (1991): Natural populations of entomopathogenic nematodes (Rhabditida: Heterorhabditidae and Steinernematidae) from the Hawaiian Island. Environmental Entomology, 20: 211-216.
- Herron, G.A. and G.L. Baker (1991): The effect of host stage and temperature on the development of *Hexamermis* sp. (Nematoda: Mermithidae) in the Australian plague locust *C. terminifera* (Walker) (Orthoptera: Acrididae). Nematologica, 37 (2): 213-224.
- Hoffmann, D. and N.C.R. Brechelin (1971): Changes in the blood picture in male *L. migratoria* after injection of *B. thuringiensis*. Acad. Sci. Paris, Ser. D, 274: 422-425.
- Hoffmann, D. and M.A. Porte (1973): Sur la differenciation secretoire de celluli reticulaire de l'organe hemoatopoietique de *L. migratoria* immunisee contre *B. thuringiensis* Etude au microscope electrique, C.R. Acad. Sci. Paris, t. 276, D, 677.
- Hoffmann, D.; M. Brehelin and J.A. Hoffman (1974): Modifications of the haemogram and of the haemocytopoietic tissue of male adults of

- L. migratoria* after injection of *B. thuringiensis* (Bacteria). J. Invert. Pathol., 24: 238-247.
- Hoffman, J.A.; M.E. Stockel; A. Porte and P. Joly (1969): Ultrastructure des haemocytes de *L. migratoria* (Orthoptere). C.R. Acad. Sci. Paris, 266: 503-505.
- Hominick, W.M. and B.R. Briscoe (1990): Occurrence of entomopathogenic nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) in British soils. Parasitology, 100: 295-302.
- Horohov, D.W. and P.E. Dunn (1982): Changes in the circulating haemocyte population of *M. sexta* larvae following injection of bacteria. J. Invert. Pathol., 40: 327-339.
- Hudson, W.G. and K.B. Nguyen (1989): Effects of soil moisture, exposure time, nematode age and nematode density on laboratory infection of *S. vicinus* and *S. acletus* (Orthoptera: Gryllotalpidae) by *Neoaplectana* sp. (Rhabditida: Steinernematidae). Environmental Entomology, 18 (4): 719-722.
- Hung, S.Y. and D.G. Boucias (1992): Influence of *B. bassiana* on the cellular recognition response of the beet armyworm, *S. exigua*. J. Invert. Pathol., 60: 152-158.
- Hung, S.Y.; D.G. Boucias and A.J. Vey (1993): Effect of *B. bassiana* and *C. albicans* on the cellular defence response of *S. exigua*. J. Invert. Pathol., 61: 179-187.

- Hunter-Jones, P. (1961): Rearing and breeding locusts in the laboratory. Bull. Anti-locust Research Centre London, 12 pp.
- Ishibashi, N. and E. Kondo (1986): *S. feltiae* (DD-136) and *S. glaseri*: Persistence in soil and bark compost and their influence on native nematodes. J. Nematol., 18: 310-316.
- Jaques, R.P. (1967): Mortality of five apple insects induced by the nematode DD-136. J. Econ. Entomol., 60: 741-743.
- Jones, J.C. (1962): Current concepts concerning insect haemocytes. Amer. Zool., 2: 209-246.
- Jones, J.C. (1967): Changes in the haemocyte picture of *G. mellonella* (Linnaeus). Biol. Bull. (Woods Hole), 132: 211-221.
- Kamionek, M. (1977): Pathogenicity of the nematode *N. carpocapsae* Weiser in relation to different hosts. Bull. Acad. Pol. Sci. Cl., 25: 243-246.
- Kamionek, M. and H. Seryczynska (1976): Effect of nematodes and parasitic fungi on Colorado beetle. Bull. de l'Acad. Polonaise Scie., Biologiques, 24: 481-483.
- Kamionek, M.; I. Maslana and H. Sandner (1974): The survival of invasive larvae of *N. carpocapsae* Weiser in a waterless environment under various conditions of temperature and humidity. Zesz. Post. Nauk Roln., 154: 409-412.
- Kaya, H.K. (1977): Development of the DD-136 strain of *N. carpocapsae* at constant temperatures. J. Nematol., 9: 346-351.

- Kaya, H.K. (1978): Interaction between *S. carpocapsae* (Nematoda: Steinernematidae) and *A. militaris* (Hymenoptera: Brachonidae), a parasitoid of the army worm *Pseudaltia unipuncta*. J. Invert. Pathol., 31: 358-364.
- Kaya, H.K. (1985a): Entomogenous nematodes for insect control in IPM system, pp. 283-302. In M. Hoy & D.C. Herzog [eds.], Biological control in agricultural IPM systems. Academic, New York.
- Kaya, H.K. (1985b): Susceptibility of early larval stages of *P. unipuncta* and *S. exigua* to entomogenous nematode *S. feltiae*. J. Invert. Pathol., 46: 58-62.
- Kaya, H.K. (1986): *S. feltiae* use against foliage feeding insects and effect on non-target insects, pp. 268-270. In R.A. Samson; J.M. Vlak, and D. Peters [Eds.], Fundamental and Applied Aspects of Invertebrate Pathology.
- Kaya, H.K. (1993): Entomogenous and entomopathogenic nematodes in biological control. In plant parasitic nematode in temperate agriculture. Edited by K. Evans; D.L. Trudgill and J.M. Webster. CAB International, Wallingford, Oxon OX10 8DE UK. pp. 565-591.
- Kaya, H.K and R. Gaugler (1993): Entomopathogenic nematodes. Ann. Rev. Entomol., 38: 181-206.
- Klein, M.G. (1990): Nematode efficacy against soil-inhabiting insect pests. PP. 195-214 in R. Gaugler and H.K. Kaya (eds): Entomopathogenic nematodes in biological control. Boca Raton, Fl: CRC Press.

- Kondo, E. (1987): Size related susceptibility of *S. litura* (Lepidoptera: Noctuidae) larvae to entomogenous nematode *S. feltiae* (DD-136). Appl. Entomol. Zool., 22 (4): 506-509.
- Kondo, E. (1989): Studies on the infectivity and propagation of entomogenous nematodes *Steinernema* sp. (Rhabditida: Steinernematidae) in the common cutworm, *S. litura* (Lepidoptera: Noctuidae). Bull. Fac. Agric. Saga Univ., 67: 1-88.
- Kondo, E. and N. Ishibashi (1987): Ultrastructure differences between the propagative and dispersal form in pine wood nematode, *Bursaphelenchus lignicolus*, with reference to survival. Appl. Entomol. Zool., 13: 1-11.
- Koppenhofer, A.M. and H.K. Kaya (1995): Density-dependent effects on *S. glaseri* (Nematoda: Steinernematidae) within an insect host. Journal of Parasitology, 81 (5): 797-799.
- Kung, S.P. (1990): Abiotic factors affecting the persistence of two entomopathogenic nematodes, *S. carpocapsae* and *S. glaseri* (Nematoda: Steinernematidae) in the soil. Dissertation Abstracts International. B, Sciences and Engineering 51: 4, 1620; Abstract of Thesis, Rutgers State University of New Jersey, USA, 134 pp.
- Kung, S.P.; R. Gaugler and H.K. Kaya (1991): Effects of soil temperature, moisture and relative humidity on Entomopathogenic nematode persistence. J. Invert. Pathol., 57: 242-249.

- Lacey, L.A.; H.K. Kaya and R. B. Court (1995): Dispersal of *S. glaseri* (Nematoda: Steinernematidae) in adult Japanese beetles, *P. japonica* (Coleoptera: Scarabaeidae). *Biocontrol Science and Technology*, 5 (1): 121-130.
- Lindegren, J.E. and P.V. Vail (1986): Susceptibility of Mediterranean fruit fly, melon fly, and oriental fruit fly (Diptera: Tephritidae) to the entomogenous nematode *S. feltiae* in laboratory tests. *Environment. Entomol.*, 15: 465-468.
- Liu, X.L. and J.X. Wang (1989): Preliminary study on pathogenicity of different strains of the nematode *S. glaseri* steiner to the sugarcane beetle *Alissonotum impressicolle* Arrow. *Natural Enemies of Insects*, 3 (11): 101-103.
- Lopez-Lastra, C.C. and D.G. Boucias (1994): Studies on the cellular reactions of *S. exigua* larvae infected with the fungus *N. rileyi*. *J. Invert. Pathol.*, 63: 101-102.
- Lowery, O.H.; N.J. Rosebrough; A.L. Farr and R.I. Randall (1951): Protein measurement with the folin phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- Lysenko, O. (1966): Bacterial toxins. In "Insect Physiology and Microbial Control" (P.A. Van der Lann, Ed.), North Amesterdam: 214-237.
- Mac-Vean, C.M.; J.W. Brewer and J.L. Capinera (1982): Field tests of antidesiccants to extend the infection period of an entomogenous nematode, *N. carpocapsae* against the Colorado potato beetle. *J. Econ. Entomol.*, 75: 97-101.

- Milstead, J.E. (1981): Influence of temperature and dosage on mortality of seventh instar larvae of *G. mellonella* (Insecta: Lepidoptera) caused by *H. bacteriophora* (Nematoda: Rhabditida) and its bacterial associate *X. luminescens*. *Nematol.*, 27: 167-175.
- Miselyurenen, I.S. (1976): Pathogenic changes in the morphology and number of haemolymph haemocytes of cabbage butterfly larvae infected with *intobacterin*³. *Thistologia*, 18 (10): 1220-1225.
- Mitro, S. (1994): Cellular defence reaction of the haemolymph, and description of specific cells in the seminal plasma of *A. mellifera* L. *Apidologie*, 25 (4): 361-366.
- Molyneux, A.S. (1985): Survival of infective juveniles of *Heterorhabditis* sp. and *Steinernema* sp. (Nematoda: Rhabditida) at various temperatures and their subsequent infectivity for insects. *Rev. Nematol.*, 8: 165-170.
- Molyneux, A.S. (1986): *Heterorhabditidis* sp. and *Steinernema* (= *Neoaplectana*) sp.: Temperature and aspects of behavior and infectivity. *Exp. Parasitol.*, 62: 169-180.
- Molyneux, A.S.; R.A. Bedding and R.J. Akhurst (1983): Susceptibility of larvae of the sheep blowfly, *L. cuprina* to various *Heterorhabditis* sp., *Neoaplectana* sp. and undescribed steinernematid (Nematoda). *J. Invert. Pathol.*, 42: 1-8.
- Nicolas, D.; N.D. Epsky and J.L. Capinera (1995): Susceptibility of *M. sanguinipes* (Orthoptera: Acrididae) nymphs to *S. carpocapsae*

- and *S. scapterisci* (Nematoda: Steinernematidae). *Environmental Entomology*, 24 (3): 762-769.
- Ogel, S. (1955): A contribution to the study of blood cells in Orthoptera. *Commun. Fac. Sci. Ankara, C*, 4: 15-41.
- Parkman, J.P.; J.H. Frank; K.B. Nguyen and G.C.Jr. Smart (1993a): Dispersal of *S. scapterisci* (Rhabditida: Steinernematidae) after inoculative applications for mole cricket (Orthoptera: Gryllotalpidae), control in pastures. *Biological control*, 3 (3): 226-232.
- Parkman, J.P.; W.G. Hudson; J.H. Frank; K.B. Nguyen and G.C.Jr. Smart (1993b): Establishment and Persistence of *S. scapterisci* (Rhabditida: Steinernematidae) in field populations of *Scapteriscus* sp. mole crickets (Orthoptera: Gryllotalpidae). *Journal of Entomological Science*, 28 (2): 182-190.
- Parkman, J.P.; J.H. Frank; K.B. Nguyen and G.C.Jr. Smart (1994): Inoculative release of *S. scapterisci* (Rhabditida: Steinernematidae) to suppress pest mole crickets (Orthoptera: Gryllotalpidae) on golf courses. *Environmental Entomology*, 23 (5): 1331-1337.
- Patel, M.C. and R.V. Vyas (1995): Efficacy of *S. glaseri* against *H. armigera* on chickpea in pots. *International Chickpea and Pigeonpea-Newsletter*. No. 2, 39-40.
- Patel, M.N.; R.N. Perry and D.J. Wright (1997): Desiccation survival and water contents of entomopathogenic nematodes, *Steinernema* sp.

- (Rhabditida: Steinernematidae). International Journal for Parasitology, 27 (1): 61-70.
- Poinar, G.O.Jr. (1966): The presence of *A. nematophilus* in the infective stage of *Neoaplectana* sp. (Nematoda). Nematologica, 12: 105-108.
- Poinar, G.O.Jr. (1979): Nematodes for biological control of insects. CRC Press, Boca Raton, Fl., 277 & 249 pp.
- Poinar, G.O.Jr. (1986): Entomogenous nematodes, pp. 95-121. In B.D. Franz [Eds], Biological Plant and Health Protection. Fischer, Stuttgart.
- Poinar, G.O.Jr. (1990): Taxonomy and biology of Steinernematidae and Heterorhabditidae. In Entomopathogenic Nematodes in Biological Control (Gaugler, R. and Kaya, H.K., Eds) CRC Press, Boca Raton, Florida, USA, 23-61.
- Poinar, G.O.Jr. and G.M. Thomas (1965): A new bacterium, *A. nematophilus* (Achromobacteraceae: Eubacteriales) associated with a nematode. Int. Bull. Bacteriol. Nematol. Taxon., 15: 249-252.
- Poinar, G.O.Jr. and G.M. Thomas (1966): Significance of *A. nematophilus* Poinar and Thomas in the development of the nematode, DD-136. Parasitol., 56: 385-390.
- Poinar, G.O.Jr. and G.M. Thomas (1967): The nature of *A. nematophilus* as an insect pathogen. J. Invert. Pathol., 9: 510-514.

- Pryce, M.J.; W.P. Aston and J.S. Chadwick (1990): Cane sugar factor as an inducing agent of immunity in *G. mellonella*. *Developmental and Comparative Immunology*, 14: 369-378.
- Ramadan, R.H. (1994): Susceptibility of different pests to infection of entomogenous nematodes. Ph.D. Thesis, Faculty of Science, Zagazig University.
- Ratcliffe, N.A. (1975): Cellular defence reactions of insect haemocytes in vitro phagocytosis in a new suspension culture system. *J. Invert. Pathol.*, 26: 225-233.
- Ratcliffe, N.A. and S.J. Gagen (1976): Cellular defence reactions of insect haemocytes in vitro: Nodule formation and development in *G. mellonella* and *P. brassicae* larvae. *J. Invert. Pathol.*, 28 (3): 373-382.
- Ratcliffe, N.A. and J.B. Walters (1983): Studies on the in vivo cellular reactions of insects: Clearance of pathogenic and non-pathogenic bacteria in *G. mellonella* larvae. *J. Insect Physiol.*, 29: 407-415.
- Robert, G.D. Stell and J.H. Torrie (1980): Principles and procedures of statistics, a biometrical approach (2nd Edition). Mc. Jraw-Hill Book Campany New York st.
- Rosenberger, C.R. and J.C. Jones (1960): Studies on total blood counts of the Southern armyworm larva, *Prodenia eridania* (Lepidoptera). *Ann. Entomol. Soc. Ameri.*, 53: 351-355.

- Ryan, N.A. and R.D. Karp (1993): Stimulation of haemocyte proliferation in the american Cockroach (*P. americana*) by the infection of *Enterobacter cloacae*. J. Insect Physiol., 39 (7): 601-608.
- Saleh, M.M.E. and Tayseer, A.H. El-Kifl (1994): Virulence of certain entomopathogenic nematodes to the diapausing larvae of the European corn borer, *O. nubilalis* (Hbn.). Egypt. J. Biol. Pest Control, 4 (2): 125-131.
- Saleh, M.M.E.; A.M. Semeada and E.A. El-Shazly (1994): Virulence of native and foreign entomopathogenic nematodes to the apple borer, *Z. pyrina*. Egypt. J. Biol. Pest Control, 4 (2): 107-112.
- Salt, G. (1970): "The cellular defence reactions of insects". Monographs in Experimental Biology. Vol.16 . Cambridge Univ. Press, London.
- Schmiedege, D.C. (1963): The feasibility of using of a neoaplectanid nematode for control of some forest insect pests. J. Econ. Entomol., 56: 427-431.
- Seryczynska, H. and M. Kamionek (1977): Stimulation of phagocytic cells of *G. mellonella* by microorganisms. Bull. Acad. Pol. Sci. Ser. Sci. Biol., 25: 173-176.
- Shamseldean, M.M. (1994): Effects of temperature on survival and infectivity of Egyptian heterorhabditid nematode isolates. Egypt. J. Appl. Sci., 9 (9): 53-59.

- Shamseldean, M.M. and M.M. Abd-Elgawad (1994): Laboratory evaluation of six Egyptian isolates of heterorhabditid nematodes for control of the red palm weevil. *Egypt. J. Appl. Sci.*, 9 (3): 670-679.
- Shamseldean, M.M. and M.M. Abd-Elgawad (1995): Survival and infectivity of entomopathogenic nematodes under environmental stress. *Anz. Schadlingskde., Pflanzenschutz, Umweltschutz*, 68: 31-33.
- Shamseldean, M.M. and M.M. Abd-Elgawad (1996): Survival and recycling of an Egyptian isolate of *H. indicus* compared to *S. glaseri*. *Afro-Asian Journal of Nematology*, 6 (1): 94-97.
- Shamseldean, M.M. and A.A. Ismail (1997): Effect of the nematode *H. bacteriophora* and the bacterium *B. thuringiensis* as integrated biocontrol agents of the black cutworm. *Anz. Schadlingskde., Pflanzenschutz, Umweltschutz*, 70: 77-79.
- Shamseldean, M.M.; M.M. Abd-Elgawad and A.A. Atwa (1996a): Evaluation of four entomopathogenic nematodes against *S. littoralis* (Lepidoptera: Noctuidae) larvae under different temperatures. *Anz. Schadlingskde., Pflanzenschutz, Umweltschutz*, 69: 111-113.
- Shamseldean, M.M.; M.M. Abd-Elgawad and M.H. El-Bishry (1994): Comparative infectivity of native heterorhabditid nematodes and the potential use of one isolate against some economically important pests in Egypt. *Egypt. J. Biol. Pest Control*, 4 (2): 133-139.
- Shamseldean, M.M.; A.B. Abou El-Sooud; M.M. Abd-Elgawad and M.M. Saleh (1996b): Identification of a new heterorhabditid species from

- Egypt, *H. tysearae*, N. sp. (Rhabditida: Heterorhabditidae). Egypt. J. Biol. Pest Control, 6 (2): 129-138.
- Shamseldean, M.M.; A.B. Abou El-Sooud and M.M. Saleh (1996c): Isolation of *S. carpocapsae* (Weiser, 1955) Poinar, 1990 (Steinernematidae), *H. bacteriophora* Poinar, 1976 and *H. indicus* Poinar, *et al.* 1992 (Heterorhabditidae) as a first record from Egypt. Egypt. J. Biol. Pest Control, 6 (2): 185-199.
- Shanti, A.N. and C.V. Sivakumar (1993): Comparative virulence of *S. glaseri* (Steiner) and *S. carpocapsae* (Weiser) to the chafer *H. consanguinea* (Coleoptera: Scarabaeidae). Indian Journal of Nematology, 21 (2): 149-152.
- Shapiro, M. (1966): Pathogenic changes in the blood of the greater wax moth, *G. mellonella* (Linnaeus), during the course of starvation and nucleopolyhedrosis. Ph.D. Thesis, University of California, Berkeley, California.
- Shapiro, M. (1967): Pathogenic changes in the blood of the greater wax moth, *G. mellonella* (Linnaeus), during the course of nucleopolyhedrosis and starvation. I. Total haemocyte count. J. Invert. Pathol., 9: 111-113.
- Sikora, R.A.; I.E.M. Salem and F. Klingauf (1978): Susceptibility of *S. littoralis* to *N. carpocapsae* and observations on the effect of environmental factors on insect mortality. Proc. 4th Conf. Pest Control, NRC, 11: 490. Cairo.

- Sikora, R.A.; I.E.M. Salem and F. Klingauf (1979): Susceptibility of *S. littoralis* to *N. carpocapsae* and the importance of environmental factors on an insect control program. Med. Fac. Landbouww. Rijks Univ., Gent., 44: 309-321.
- Simons, W.R. and G.O.Jr. Poinar (1973): The ability of *N. carpocapsae* (Nematoda: Steinernematidae) to survive extended periods of desiccation. J. Invert. Pathol., 22: 228-230.
- Surrey, M.R. and D.A. Wharton (1995): Desiccation survival of the infective larvae of the insect parasitic nematode, *H. zealandica* Poinar. International Journal for Parasitology, 25 (6): 749-752.
- Swain, S.B.; S. Litting; F.J. Barlett and M.F. Gordon (1944): Studies on nematodes of the genus *Neoaplectana*, factors in the biological control of the white-fringed beetles. Special USDA Report 7-1, Gulfport, Miss. 33 pp.
- Triggiani, O. and G.O.Jr. Poinar (1976): Infection of adult Lepidoptera by *N. carpocapsae* (Nematoda). J. Invert. Pathol., 27: 413-414.
- Uvarov, B. (1966): Grasshoppers and locusts. A handbook of general acridology, vol. 1, Cambridge University Press.
- Vyas, R.V. and D.N. Yadav (1993): *Steinernema glaseri* (Steiner) Travassos, a biological control agent of root grub *H. consanguinea* Blanchard. Pakistan Journal of Nematology, 11 (1): 41-44.
- Wang, Y.L.; J.F. Campbell; R. Gaugler and Y. Wang (1995): Infection of entomopathogenic nematodes, *S. glaseri* and *H. bacteriophora*

- against *P. japonica* (Coleoptera: Scarabaeidae) larvae. J. Invert. Pathol., 66 (2): 178-184.
- White, G.F. (1927): A method for obtaining infective nematode larvae from cultures. Science, 66: 302-303.
- Wienands, A. and G. Madel (1988): Haemocytes of honeybee, *A. mellifera*, and their changes during *Varroa* disease. Entomologia Generalis, 14 (2): 81-92.
- Wienands, A.; H. Strich and G. Madel (1987): Haemocytic and bacteriological aspects of the haemolymph of *A. mellifera* L. (Hymenoptera: Apidae). Zentralblatt-fur-Bakteriologie, Mikrobiologie-und-Hygiene, A. 265, 489; Ba (Apic. Abst. 1264/89).
- Wijbenga, J. (1991): Possibility of obtaining a monoxenic culture of the north west European insect parasitic nematode *Heterorhabditis* sp. Mededelingen-Van-de-Faculteit-Landbouwwetenschappen, Rijksuniversiteit Gent, 56 (3b): 1297-1304.
- Williams, E.C. and K.F.A. Walters (1996): Strategic incorporation of biocontrol agents for managing alien insect pests. Brighton Crop Protection Conference: Pests and Diseases, vol. 3, Proceedings of an International Conference, Brighton, UK, 1071-1076.
- Witting, G. (1965): Phagocytosis by blood cells in healthy and diseased caterpillars. I. phagocytosis of *B. thuringiensis* Berliner in *P. unipuncta* (Haworth). J. Invert. Pathol., 7: 474-488.

- Witting, G. (1966): Phagocytosis by blood cells in healthy and diseased caterpillars. II. A consideration of the method of making haemocyte counts. *J. Invert. Pathol.*, 8: 461-477.
- Yamanaka, S.; H. Tanabe and K. Takeuchi (1995): Vertical dispersal and infectivity of *S. glaseri*, *S. anomali* and *S. Kushidai* (Nematoda: Steinernematidae). *Japanese Journal of Nematology*, 25 (1): 24-32.
- Zervos, S.; S.C. Johnson and J.M. Webster (1991): Effect of temperature and inoculum size on reproduction and development of *H. heliothidis* and *S. glaseri* (Nematoda: Rhabditoidea) in *G. mellonella*. *Canadian Journal of Zoology*, 69 (5): 1261-1269.
- Zidan, Z.H.; G.M. Moawad; A.I. Gadallah and F.E. El-Sweeki (1996): Biochemical aspects of the cotton leafworm larvae *S. littoralis* (Boisd) as affected by soft nontoxic insecticides. Proceedings: Sixth conference of agricultural development research, Cairo. *Annals of Agricultural Science, Cairo*, No. special issue, 233-244.