

# References

**Abramovici D, Friedman S, Mercer BM, Audibert F, Koal and Sibai B:** National outcome in severe preeclampsia at 24-36 weeks gestation = dose the HELLP syndrome matte? *Am J. Obstet Gynecol 1999; 180: 221-225*

**Abuja PM:** Ascorbate prevents pro-oxidant effects of urate in oxidation of human low density lipoprotein. *FEBS Lett. 1999; 446: 305-8.*

**Agraharkar M and Kelly B:** Renal Disease and Pregnancy. 2003; *emidicine.com, Inc.*

**Ahmed Y, van Iddekinge B, Paul C, Sullivan MHF and Elder MG:** Retrospective analysis of platelet members and volumes in normal pregnancy and in preeclampsia. *Br J. Obstet Gynecol 1993; 52: 100-104.*

**Alexander BT, Kassab SE, Miller MT, Abram SR, Reckelhoff JF, Bennett WA, and Granger JP:** Reduced uterine perfusion pressure during pregnancy in the rat is associated with increases in arterial pressure and changes in renal nitric oxide. *Hypertension; 2001; 37: 1191-5.*

**Alexander BT, Reckelhoff JF, Kassab S, and Granger JP:** Differential expression of renal nitric oxide synthase isoforms during pregnancy in rats. *Hypertension 1999; 33: 435-439, 1999.*

**American Collage of Obstetric and Gynecology (ACOG) Committee:** Diagnosis and management of preeclampsia and eclampsia. *Obstet Gynecol; 2001; 98: 159-67.*

**Anderson, S: Proteinuria.** In: Greenburg, A. editor. *Primer on Kidney Diseases* (2nd edition). Boston: National Kidney Foundation; Academic Press: 1998; *Pp. 42-6.*

**Aquiline J, Barnett a, Thompson O and Harrington K:** Second trimester maternal seum inhibin A concentrations as an early marker for preclampsia. *Am J Obstet Gynecol 1999; 181: 131-137.*

- Arbogast BW, Leeper SC, Merrick RD, Live KE and Taylor RN:** Hypothesis which plasma factors ring about disturbance of endothelial function in preeclampsia? *Lancet AmJ* 1994; 343: 340-341.
- Arngrimsson R, Bjornsson S and Geirsson RT:** Genetic and familial predisposition to eclampsia and pre-eclampsia in a defined population. *Br. J Obstet Gynecol* 1990; 97: 762-769.
- Arngrimsson R, Hayward C and Nadaud S:** Evidence for a familial pregnancy induced hypertension locus in the e NOS-gene region. *Am J Hum Genet* 1997; 61: 354-362.
- Arngrimsson R, Purandare S and Connor M:** Angiotensinogen-a candidate gene involved in preeclampsia. *Nat Genet* 1993; 4: 115-124.
- Ashour AMN, Lieberman ES, Wilkins Haug LE and Repke JT:** The value of elevated second trimester  $\beta$ -human chorionic gonadotropin in predicting development of preeclampsia. *Am J Obstet Gynecol* 1997; 176: 438-440.
- August P and Lindheimer MD:** Pathophysiology of preeclampsia. *Hypertension*; 1995; 142: 2407-26.
- Baker PN, Krasnow J, Roberts JM and Yeo K:** Elevated serum levels of vascular endothelial growth factor in patients with preeclampsia. *Obstet Gynecol* 1995; 86: 815-817.
- Bardeguet AD, McNeerney R, Frieri M, Verma UL and Tejani N:** Cellular immunity in preeclampsia, alterations in T-lymphocyte subpopulations during early pregnancy. *Obstet Gynecol* 1991; 77: 859-866.
- Barron WM, Lindheimer MD and Davison JM:** Hypertension in medical disorders during pregnancy, United States of America; Mosby. 2000; 58:45-48.
- Barton JR, O'Brien JM, Bergauer NK, Jacques DL and Sibai BM:** Mild gestational hypertension remote from term: progression and outcome. *Am J Obstet Gynecol* 2001; 184: 979-83.
- Baylis C, Beinder E and Suto T:** Recent insights into the roles of nitric oxide and renin angiotensin in the pathophysiology of preeclamptic pregnancy. *Semin Nephrol.* 1998; 18:208-230.

**Beaufils M:** Hypertensive disorders of pregnancy. *La Revue de Médecine Interne*; 2002; 23(11) : 927-38.

**Benyo DF, Adatisa P and Conrad KP:** Levels of inflammatory cytokines in normal term and preeclamptic placentas and their regulation by oxygen. *Gynecol Investig* 2000; (*Abstr* 894), 7 (1) (*Suppl*).

**Bewley S, Cooper D and Campbell S:** Doppler investigation of uteroplacental blood flow resistance in the second trimester: screening study for preeclampsia and intrauterine growth retardation. *Br J Obstet Gynecol* 1991; 98: 871-874.

**Biagi G, De Rosa V, Pelusi G, Scagliarini G, Sani G and Coccheri S:** Increased placental production of leukotriene B in gestational hypertension. *Thromb Res*. 1990; 60: 377-381.

**Bosolo P, O Herlihy C and Conroy R:** Maternal Central hemodynamics in hypertensive disorders of pregnancy. *AM J Obstet Gynecol* 2001; 15:178-861.

**Bower S, Bewley S and Campbell S:** Improved prediction of preeclampsia by two-stage screening of uterine arteries using the early diastolic notch and the color Doppler imaging. *Obstet Gynecol* 1993; 82: 78-82.

**Bower Sarah; Katharin Schlter and Stuart Campbell:** Doppler ultrasound screening as part of routine antenatal scanning; prediction of preeclampsia and intrauterine growth retardation. *Br. J. Gynecol*. 1993; 100: 625-629.

**Brookes MB, Sayah AJ, Talavera F, Zwanger M and Halamka J:** Pregnancy and preeclampsia. 2004; *eMedicine.com, Inc*.

**Brown MA, Gallery EDM, Ross Smr and Esber RP:** Sodium excretion in normal and hypertensive pregnancy. A prospective study. *AM J Obstet Gynecol* 1988; 159:297-307.

**Brown MA, Zammit VC and Mitor DM:** Extracellular fluid volumes in pregnancy – induced hypertension. *J Hypertens* 1999; 10 : 821-829.

- Brown MS and Gallery EDM:** Volume homeostasis in normal pregnancy and preeclampsia: physiology and clinical implications. *Clin Obstet Gynecol.* (Baillier's) 1994; *8*: 297-320.
- Brubaker DB, Ross MG and Marinoff D:** The function of elevated plasma fibronectin in preeclampsia. *Am J Obstet Gynecol.* 1992; *166*: 526-531.
- Brunner HIR and Gavras H:** Vascular damage in hypertension. *Hosp Pract.* 1975; *10*: 97-99.
- Buchbinder A, Sibai BM, Caritis S, Macpherson C, Hauth J and Lindheimer MD:** Adverse perinatal outcomes are significantly higher in severe gestational hypertension than in mild preeclampsia. *Am J Obstet Gynecol;* 2002; *186*: 66-71.
- Butler R, Morris AD, Belch JJ, Hill A and Struthers AD:** Allopurinol normalizes endothelial dysfunction in type 2 diabetics with mild hypertension. *Hypertension;* 2000; *35*: 746-51.
- Byrd DE and Riely CA:** liver disease in preeclampsia. *Gastroenterologist,* 1996; *4*: 65-69.
- Cant NF, Worley FJ and Everett RB:** Control of vascular responsiveness during human pregnancy. *Kidney Int.* 1980; *253*-258.
- Caron C, Goudemaud J, Marey A, Beague D, Ducroux G and Drouvin F:** Are haemostatic and fibrinolytic parameters predictors of preeclampsia in pregnancy- associated hypertension? *Thromb Haemost* 2000; *66*: 410
- Carroll G, Duley L, Bellizain JM and Villar J:** Calcium supplementation during pregnancy. A systematic review of randomized controlled trials. *Br J Obstet Gynecol* 1994; *101*: 753-8.
- Catella F, Lawson JA, Fitzgerald DJ and Fitzgerald GA:** Endogenous biosynthesis of arachidonic acid peroxides in humans increased formation in pregnancy induced hypertension. *Proc Na Acad Sci USA,* 1990; *87*: 5893-5899.
- Chappell L and Bewley S:** Preeclamptic toxemia: The role of uterine artery Doppler. *Br J Obstet Gynecol* 1998; *105*: 379-383.
- Chappell LC, Paul T, Annette B Frank RM and Kelly J :**A longitudinal study of biochemical variables in women at risk of preeclampsia *Am J Obstet Gynecol;* 2002; *187*: 127-36.

- Chen G, Wilson R, McKillop J and Walker J :** The role of cytokines in the production of prostacyclin and thromboxane in human mononuclear cells. *Immunol invest* 1994; *23*: 269-279.
- Chen G, Wilson R, Wang S H, Zheng H Z, Walker J and McKillop J H:** Tumor necrosis factor alpha (TNF alpha) gene polymorphism and expression in preeclampsia. *Clin Exp Immunol* 1996; *104*: 154-159.
- Chesley LC:** Diagnosis of preeclampsia. *Obstet Gynecol* 1985; *65*: 423-327.
- Chumbely G, King A, Robertson K. Holmes N and Loke YW:** Resistance of HLA-G and HLA-A2 transfectants to lysis by decidual NK cells. *Immunol* 1994; *155*: 312-322.
- Chung JK, Roman C and Heymann MA:** Effect of endothelium – derived relaxing Am J Obstet Gynecol. 1992; *166*: 727-729.
- Chung YK, Copel JA, Lee IS, Chou K and Hsu CD:** Serum uric acid levels in preeclamptic multiple gestations. *J Soc Gynecol Invest.*; 1996; *3*(2): *Suppl. 2*.
- Cincotta RB and Brennecke SP:** Family history of preeclampsia as a predictor for preeclampsia in primigravida. *Int. J. Gynecol. Obstet.* 1998; *60* (1): 23-30.
- Clark BA, Halvorson L, Sachs B and Epstein FH:** Plasma endothelin levels in preeclampsia: elevation and correlation with uric acid levels and renal impairment. *Am J Obstet. Gynecol.*; 1992; *166*: 962-8.
- Clark BA, Halvorson L, Sachs B and Epstein FH:** Plasma endothelin levels in preeclampsia; Elevation and correlation with uric acids levels and renal impairment. *Am J Obstet Gynecol.* 1992; *166*: 962-966.
- Clark SL, Greenspoons JS and Aldahi D:** Severe preeclampsia with persistent oliguria; Management of hemodynamic subsets. *Am J Obstet Gynecol.* 1986; *154*: 490-494.
- Cnattingius S, Mills JL, Yuen J, Eriksson O and Salonen H:** The paradoxical effect of smoking in preeclampsia pregnancies smoking reduces the incidence but increases the rates of

perinatal mortality, abruptio placentae, and intrauterine growth restriction. *Am J Obstet Gynecol* 2001;177:156-161.

**Cockell A P, Learmont J G, Smarason AK, Redman C WG, Sargent I L and Poston L:** Human placental syncytiotrophoblast microvillous membranes impair maternal vascular endothelial function. *Br J Obstet Gynecol* 1997;104: 235-240.

**Collr Bell A, Adair LS and Popkin BM:** Ethnic differences in the association between body mass index and hypertension. *Am J epidemiology* 2002;155: 346-353.

**Conde- Agudelo A and Belizan JM:** Risk faotors for pre-eclampsia in a large cohort of Latin American and Caribbean Women. *Br. J. Obstet Gynecol* 2000;107 (1), PP- 75-83.

**Conde- Agudelo A, Lede R and Belizan J:** Evaluation of methods used in the prediction of hypertensive disorders of pregnancy. *Obstet Gynecol Sury* 1994; 49: 210-222.

**Conde-Agudelo A, Lede R and Belizan J:** Evaluation of methods used in the prediction of hypertensive disorders of pregnancy. *Obstet Gynecol*; 1994; 49: 210-21.

**Conrad KP and Lindheimer MD:** Renal and cardiovascular changes in normal pregnancy and preeclampsia. In Lindheimer MD, Roberts JM Cunningham FG, Editors. Chesley's hypertensive disorders in pregnancy. Standford, CT Appleto & Lange, 1999; pp 395-429.

**Cotter A, Molloy A and Scoot Daly SF:** Elevated plasma homocysteine in early pregnancy: A risk factor for the development of severe preeclampsia. Presented at the 21st Annual Meeting of the Society of Maternal – Fetal Medicine. 2001 February; held in Reno, Nevada: 5-10

**Cuckle H, Sehmi I and Jones R:** Maternal serum inhibin A can predict preeclampsia. *Br J Obstet Gynecol* 1998;105: 1101-1106.

**Cunningham FG, MacDonald PC and Gant NF:** Hypertensive Disorders in Pregnancy. In Williams Obstetrics 20<sup>th</sup> Edition, United States of America: Appleton & Lange, 1997; PP.693-716.

- Cunningham FG, Norman F Gant and Renneth Dleveno:** Hypertensive Disorders in Pregnancy. In Williams Obstetrics 21<sup>th</sup> Edition, United states of America: Appleton & lange, 2001; PP.567-618.
- Cuo G, Lade JA and Wilton AN:** Genetic susceptibility to preeclampsia and chromosome 7 q 36. Hum-Reprod. 1999;105: 641-647.
- Davidge ST, Hubel CA. Brayden RD, Capeless EC and McLaughlin MK:** Sera antioxidant activity in uncomplicated and preeclamptic pregnancies. Obstet Gynecol. 1992; 79: 897-913.
- Davison J and Baylis C:** Renal disease In Medical Disorders in Obstetric Practice, 3<sup>rd</sup> Black well Scientific Publications' 1996; 226-305.
- Davison J and Baylis C:** Pregnancy in patients with underlying renal disease. In: Davison AM, Camerson JS, Grunfeld J-P, Kerr DNS, Ritz E & Wiewals CG (eds) Oxford University Press 1998; PP. 2327-2335.
- De Man JG, De Winter BY, Moreels TG, Herman AG, Pelckmans PA:** S-nitrosothiols and the nitrenergic neurotransmitter in the rat gastric fundus: effect of antioxidants and metal chelation. Br J Pharmacol; 1998;123: 1039-46.
- Dekker GA and Sibai BM:** Early detection of preeclampsia. Am J Obstet Gynecol; 1991; 165: 160-72.
- Dekker GA and Sibai BM:** Etiology and pathogenesis of preeclampsia Current concepts. Am J Obstet Gynecol 1998;179: 1359-1365.
- Dekker GA, Kraayenbrink AA, Zeeman G and Kamp J:** Increased Plasma level of the novel vasoconstrictor peptide endothelin in sever preeclampsia. Eur J Obstet Gynecol Reprod Biol. 1991; 40: 215-220.
- Demianlzik NN, Xiong, Sannders LD, Wang FL and Freaser WD:** Impact of preeclampsion and gestationd hypertension on birthweight by gestational age. Am J Epidemiology 2002;155: 203-209.

- Dizontownson DS, Nelson LM, Easton K and Ward K:** The factor -V Leiden mutation may predispose women to severe preeclampsia. *Am J Obstet Gynecol* 1996;175: 902-905.
- Dudley DJ, Hunter C, Mitchell MD, Varner MW and Gately M:** Elevations of serum interleukin-1 concentrations in women with severe preclampsia and HELLP syndrome. *J Reprod Immunol* 1996; 31: 97-101.
- E.N Gofton. V. Capewell, R. Natale and R.J.Gratton:** Outcomes of women with gestational hypertension. *Am. J obsret. Gynecol* 2001;185-798-803.
- Easterling TR, Benedetti TJ, Schumucker BC and Millard SP:** Maternal hemodynamics in normal and preeclamptic pregnancies. A longitudinal study. *Obstet Gynecol.* 1990; 76: 1061-1064.
- Elias AN, Vaziri ND, Pandian MR, Powers DR and Domurat E:** Atrial natriuretic peptide and arginine vasopressin in pregnancy and pregnancy-induced hypertension. *Nephron Am.* 1988; 49:140-142.
- Endresen MG, Lorentzen B and Henriksen T:** Increased lipolytic activity and high ratio of free fatty acids to albumin in sera from women with preeclampsia leads to triglyceride accumulation in cultured endothelial cells. *Am J Obstet Gynecol* 1992;167: 440-447.
- Endresen MJ, Tosti E, Heimli H, Lorentzen B and Henriksen T:** Effects of free fatty acids found increased in women who develop preeclampsia on the ability of endothelial cells to produce prostacyclin cGMP and inhibit platelet aggregation. *Scand J Clin Lab invest Hen* 1994; 54:549-557.
- Endresen MJ, Tosti E, Lorentzen and Henniksen T:** sera of preeclamptic women are not cytotoxic to endothelial cells in culture. *Am J Obstet Gynecol.* 1995;172: 976-979.
- Eneroth E, Remberger M and Vahlne A:** Increased serum concentrations of interleukin -2- receptor in the first trimestr in women who later developed severe preeclampsia. *Acta Obstet Gynecol Scand.* 1998; 77: 591-593.

**Espana F, Gilibert J, Aznar, J, Estelles A, Kobayashi T and Friffin JH:** Complexes of activated protein C with  $\alpha_1$ -antitrypsin in normal pregnancy and in severe preeclampsia. *Am J Obstet Gynecol.* 1991; 164: 1310-1313.

**Faas MM, Schuiling GA, Linton EA, Sargent IL and Redman CWG:** Activation of Peripheral leukocytes in late pregnancy and experimental preeclampsia. *Am J Obstet Gynecol* 2000; 182: 351-355.

**Fahlen M and Agraharkar M:** Nephropathy, Uric acid. 2003; *eMedicine.com, Inc.*

**Farquharson CA, Butler R, Hill A and Belch JJ:** Allopurinol improves endothelial cell dysfunction in chronic heart failure. *Circulation*; 2002; 106: 221-6.

**Ferrazzani S, De Carolis S, Pomini F, Testa AC, Mastromarino C and Caruso A:** The duration of hypertension in the puerperium of preeclamptic women: relationship with renal impairment and week of delivery. *Am. J. Obstet. Gynecol.*; 1994; 171: 506-12.

**Ferris TF:** Prostanoids in normal and hypertensive pregnancy. *Hand book of hypertension in pregnancy.* Amsterdam EL Sevier, 1988; 10: 102-117.

**Fievet P, Peskov I and Desai I:** Plasma renin activity, blood uric and plasma volume in P.I.H. *Nephron* 40 quoted from high risk pregnancy by A. Calder 1995; 53:998-1102.

**Fisher SJ and Roberts JM:** Defects in placentation and placental perfusion. In Lindheimer MD, Roberts JM, Cunningham FG, editors: *Chesley's hypertensive disorders in pregnancy* standard CT, 1999; *Appleton & Lange*, pp 387-394.

**Fitzgerald DJ, Rocki W, Murray R, Mayo G and Fitzgerald GA:** Thromboxane A<sub>2</sub> synthesis in pregnancy-induced hypertension. *Lancet*, 1990; 335: 751-758.

**Fox H:** The placenta in pregnancy hypertension in Rubin PC *Handbook of hypertension.* Vol 10 hypertension in pregnancy. Elsevier Science. Amsterdam. 1988; 16:37-42.

**Franx A, Steegers EA and Deboot:** Sodium- blood pressure interrelationship in pregnancy. *J Hum Hypertension* 1999; 13: 159-166.

- Friedl HP, Till GO, Ryaa US and Ward PA:** Mediator induced activation of xanthine oxidase in endothelial cells *FASEB J* 1989; 3: 2512-2518.
- Friedman SA and Lindhemier MD:** Prediction and Differential Diagnosis Hypertensive Disorders in Pregnacy, 2<sup>nd</sup> ed. Standford, CT, Appleton & Lange, 1999; 20:45-52.
- Gaber LW and Lindheimer MD:** Pathology Kidney, Liver and brain. In Lindhemier MD, Roberts JM. Cunningham FG, editors; Chesley's hypertensive disorders in pregnancy. Standford, CT 1999; *Appleton & Lange*, pp 231-292.
- Gant NF, Chand S, Worley RJ, Whalley PJ, Crosby UD and MacDonald PC:** A clinical test useful of predicting the development of acute hypertension in pregnancy. *Am J Obstet Gynecol* 1976; 120:12-19.
- Genbacev O, Joslin R and Damsky CH:** Hypoxia alters early gestation human cytotrophoblast differentiation/invasion in vitro and models the placental defects that occur in preeclampsia. *J Clin Invest.* 1996; 97:540-550.
- Gibson P and Carson M:** Hypertension and pregnancy. 2004; *eMedicine.com Inc.*
- Goldman RK, Vlessis AA and Trunkey D:** Nitrosothiol quantification in human plasma. *Anal Biochem.* 1998; 259: 98-103.
- Graeff H, von Hugo R and Schrock R:** Recent aspects of hemostasis, hematology and hemorheology in preeclampsia-eclampsia. *Eur. J. Obstet. Gynecol. Reprod. Biol.*; 1984; 17: 91-102.
- Greer IA, dawes J, Johnston T A and Calder A:** Neutrophil activation is confined to the maternal circulation in pregnancy induced hypertension. *Obstet Gynecol* 1991; 78: 29-32.
- Greer IA, Lyall F, Perera T, Boswell F and Macaa LM:** Increased concentrations of cytokines interleukin-6 and interleukin-1 receptor antagonist in plasma of women with preeclampsia. A mechanism for endothelial dysfunction? *Obstet Gynecol* 1994; 84:937-940.

- Gurski M, Krueger S, Jurcovicova J, Brown E, Gonzalez E, Lewis D, Brooks G and Edwards M:** Detection of platelet derived growth factor and its down regulation by magnesium in the preeclamptic placenta. *Am J Obstet Gynecol.* 1996; 174:448-451.
- Halligan A, Bonnar J, Sheppard B, Darling M and Walshe J:** Haemostatic, fibrinolytic and endothelial variables in normal pregnancies and preeclampsia. *Br. J Obstet Gynecol* 1994; 101: 488-492.
- Hamai Y, Fujii T and Yamashita T:** Pathogenetic implication of interleukin -2 expressed in preeclamptic decidual tissues, a possible mechanism of deranged vasculature of the placenta associated with preeclampsia. *Am J Reprod Immunol.* 1997; 38:83-88.
- Hansen JP:** Older maternal age and pregnancy outcome; a review of the literature. *Obstet Gynecol* 1996; 41: 726-728
- Hara N, Fujii T and Okai T:** Histochemical demonstration of interleukin-2 in decidua cells of patients with preeclampsia. *Am J Reprod Immunol.* 1995; 34:44-51.
- Harris MD, Siegel LB and Alloway JA:** Gout and hyperuricemia. *Am Fam Physician;* 1999; 59(4): 925-34.
- Hauth J, Owen J, Thorn E, Goldenberg R, Sibai B and McNeils D :** Prevention of preeclampsia with low-dose aspirin in nulliparas. *J Soc Gynecol Invest.;* 1996; 3(2): 86A.
- Hauth JC, Ewell MG, Levine RL, Esterlitz JR, Sibai BM and Curet LB:** Pregnancy outcomes in healthy nulliparas women who subsequently developed hypertension. *Obstet Gynecol;* 2000; 95:24-8.
- Hayman R:** Hypertension in pregnancy. *Current Obstet Gynecol;* 2004; 14(1): 1-10
- Hnat MD, Sibai BM, Caritis S, Hauth J, Lindheimer MD and MacPherson C :** Perinatal outcome in women with recurrent preeclampsia compared with women who develop preeclampsia as nulliparas. *Am J Obstet Gynecol;* 2002; 186:422-6.
- Hogg N:** Biological chemistry and clinical potential of S-nitrosothiols. *Free Radic Biol Med.;* 2000; 28: 1478-86.

- Horstkamp BS, Kiess H and Kramer J:** Activated protein C resistance shows and association with pregnancy-induced hypertension. *Hum-Reprod.* 1993; 49: 494-505.
- Hsu C-D, Iriye B and Jonson TRB:** Elevated circulating thrombomodulin in severe preeclampsia. *AM J Obstet Gynecol.* 1999; 14: 31-37.
- Hubel CA:** Oxidative stress in the pathogenesis of preeclampsia. *Proc. Soc. Exp. Biol. Med.;* 1999; 222: 35.
- Hubel CA, Kagan VE, Kisin ER, McLaughlin MK, Roberts JM:** Increased ascorbyl radical production and ascorbate depletion in plasma from women with preeclampsia: implications for oxidative stress. *Free Radic Biol Med.;* 1997; 23: 596-609.
- Hubel CA, McLaughlin MK, Evans RW, Hawth JA, Sims CJ and Roberts JM:** Fasting serum triglycerides, free fatty acids, positively correlated and decrease within 48 hours postpartum. *Am J Obstet Gynecol* 1993; 169: 148-149.
- Hubel CA, Roberts JM and Taylor RN:** Lipid peroxidation in pregnancy new perspectives on preeclampsia. *Am J Obstet Gynecol.* 1996; 174: 975-979.
- Husu CD, Johnson TRB, Hong SF and Chan DW:** Altered circulating thrombomodulin and fibronectin levels as evidence for different degrees of vascular endothelial damage between severe and mild preeclampsia. *J Mat FET Invest.* 1989; 161: 1025-1034.
- Irion O, Masse J, Forest JC and Moutquin JM:** Prediction of preeclampsia, low birthweight of gestation and prematurity by uterine artery blood flow velocity waveform analysis in low risk nulliparous women. *B J obstet Gynecol* 1995; 5: 65-69.
- Iwasaki R, Ohkuchi A, Furuta I, Ojima T, Matsubara S, Sato I:** Relationship between blood pressure level in early pregnancy and subsequent changes in blood pressure during pregnancy. *Acta Obstet Gynecol Scand;* 2002; 81: 918-25.
- Jacobson SL, Imhof r. Manning N. Mannion V. Little D. Rey E and edman C:** The value of Doppler assessment of the uteroplacental circulation in predicting preeclampsia or intrauterine growth retardation. *Am J Obstet Gynecol* 1998; 105: 422-427.

**John R, Barton, John M, O'Brien, Nikik, Bergauer, Baha M and Sibai:** Mild gestational hypertension remote from term; Progress and outcome. *Am. J obsret Gynecol.* 2001; 162: 110-116.

**Johnson RJ, Kang D AND Feig D:** Is there a pathogenic role for uric acid in hypertension and cardiovascular and renal disease. *Hypertension*; 2003; 41: 1183-90.

**Johnson RJ, Kivlighn SD and Kim YG:** Reappraisal of the pathogenesis and consequences of hyperuricemia in hypertension, cardiovascular disease, and renal disease. *Am J Kidney Dis*; 1999; 33(2): 225-34.

**Jones DC and Hayslett JP:** Outcome of pregnancy in women with moderate or severe renal insufficiency. *N Engl J Med* 2001; 185: 798-803.

**Juriscova A, Casper RF, MacLusky NJ, Mills GB and Librach CL:** HLA-G expression during preimplantation human embryo development. *Proc Natl Acad Sci USA* 1995; 346: 1122-1126.

**Kagan VE, Tyurin VA, Borisenko GG, Fabisiak JP, Hubel CA, Ness RB, Gandley R, McLaughlin MK and Roberts JM:** Mishandling of copper by albumin: role in redox-cycling and oxidative stress in preeclampsia plasma. *Hypertens Pregnancy*. 2001; 20(3): 221-41.

**Kang D, Nakawaga T, Feng L, Truong L, Harris RC and Johnson RL :**A role for uric acid in the progression of renal disease. *J. Am. Soc. Nephrol.*; 2002; 13: 2888-97.

**Katz AI, Davison JM, Hayslett JP, Singson E and Lindheimer MD:** Pregnancy in women with kidney disease. *Kidney Int* 1996; 93: 161-166.

**Kelton JG, Hunter DJS and Naeme PB:** A platelet function defect in preeclampsia. *Am J Obstet Gynecol.* 1980; 18, 192-206.

**Kingdom J C, Macara I M and Whittle M J:** Fetoplacental circulation in health and disease. *Arch Dis Child* 1985; 65: 107-110.

**Kirshon B, Lee W, Mauer MD and Cotton DB:** Effects of low-dose dopamine therapy in the oliguric patient with preeclampsia. *Am J Obstet Gynecol.* 1994; 70: 161-163.

**Klein J and Ohuigin C:** The conundrum of nonclassical major histocompatibility complex genes. *Proc Natl Acad Sci USA* 1988; 159: 604-606.

**Knapp RG and Miller MC:** Clinical epidemiology and biostatistics. National Medical Series from Williams and Wilkins. Baltimore, Hong Kong, London, Sydney. Egyptian Edition 1994; 91: 625-627.

**Knuist M, Bonsel GJ, Zondervan HA and Treffers PE:** Low sodium diet and pregnancy-induced hypertension; A multicentre randomized controlled trial. *Br J Obstet Gynecol* 1998; 15:209-212.

**Knuist M, Bonsel GJ, Zondervan HA, Treffers PE:** Intensification of fetal and maternal surveillance in pregnant women with hypertensive disorders. *Int J Gynecol Obstet*; 1998; 61:127.

**Kupferminc MJ, Peacan AM, Wigton TR, Rehnberg KA and Socol ML:** Tumor necrosis factor- $\alpha$  is elevated in plasma and amniotic fluid of patients with severe preeclampsia. *Am J Obstet Gynecol* 1998; 105: 430-438.

**Kurdas C:** New guidelines for detecting and managing hypertension in pregnancy. *Contemporary Ob/Gyn*; 2001; 2:15-25.

**Kurki T, Hillesmaa V and Raitasalo R:** Depression and Anxiety in Early Pregnancy and Risk for Preeclampsia. *Obstet Gynecol* 1994; 170: 1752-1759.

**Kyle PM, Clark D, Buckley J, Kissane S, Coats M. de Swiet and Redman:** Second trimester ambulatory blood pressure in nulliparous pregnancy; a useful screening test for preeclampsia. *Br J Obstet Gynecol* 2000; 95: 487-490.

**Lapidus AM and Voto LS:** Effects of preeclampsia on the mother, fetus and child. 1999; *OBGYN.net Publications*.

**Leduc L, Wheeler JM, Kirshon B, Mitchell P and Cotton DB:** Coagulation profile in severe preeclampsia. *Obstet Gynecol*. 1996; 100: 914-919.

- Levine RJ:** Should the definition of preeclampsia include a rise in diastolic blood pressure  $\geq$  15mm Hg? *Am J Obstet Gynecol* 2000; *79*: 14-19.
- Lim KH, Friedman SA and Ecker JL:** The clinical utility of serum uric acid measurements in hypertensive diseases of pregnancy. *AM J Obstet Gynecol.* 2000; *182*: 225-227.
- Lindheimer MD and Katz AI:** Renal physiology and disease in pregnancy. In Seldin DW, Giebisch G, editors. *The kidney physiology and pathophysiology*, ed 2, New York 1998; *178*: 1067-1071.
- Lindhemimer M and Woodruff TK:** Activin A, inhibin A, and preeclampsia. *Lancet* 1992; *765*: 3391-3401.
- Lockwood CJ and Peters JH:** Increased plasma levels of EDI + cellular fibronectin precede the clinical signs of preeclampsia. *AM J Obstet Gynecol.* 1997; *349*: 1266-1273
- Loke YW and King A:** Immunology of human placental implantation; Clinical implications of our current understanding. *Mol Med Today* 1990; *162*: 358-362.
- Lorentzen B, Dreven CA, Endresen MJ and Henriksen T:** Fatty acid path of esterified and free fatty acids in sera of women with normal and preeclamptic pregnancy. *Br J Obstet* 1997; *65*: 153-159.
- Lyall F, Greer I A, Boswell F, Macara L M, Walker J J and Kingdom J C P:** The cell adhesion molecule, VCAM 1, is selectively elevated in serum; in preeclampsia does this indicate the mechanism of leukocyte activation? *Br. J Obstet Gynecol* 1995; *102*: 530-537.
- Lyndon MH:** Metabolism of uric acid in normal and toxemic pregnancy. *Mayo Clin Proc*; 1998; *53*: 743-51.
- Mabie WC, Ratts TW and Sibai BM:** The central hemodynamics of severe preeclampsia. *AMJ Obstet Gynecol.* 1994; *101*: 485-487.
- MacGillivray:** Preeclampsia. Hypertensive disease of pregnancy. *Lancet I* 1988; *161*: 1443-1448.

- Many A, Friedman SA, Hubel CA and Roberts JM:** Xanthine peroxidase activity in preeclamptic women is higher in invasive but not in villous trophoblast *J Soc Gynecol Invest.*; 1996; 3(2): 86A.
- Many A, Hubel CA and Roberts JM:** Hyperuricemia and xanthine peroxidase in preeclampsia, revisited. *Am J Obstet Gynecol*; 1996; 174(1): 288-91.
- Many A, Zhou Y, Fischer SJ and Roberts JM:** Xanthine oxidase/dehydrogenase immunofluorescent staining in invasive cytotrophoblast in preeclampsia. *J Soc Gynecol Invest.* 1995; 2(2): 199.
- Mastrogiannis DS, O'Brien WF, Krammer J and Benoit R:** Potential role of endothelin -1 in normal and hypertensive pregnancies. *Am J obstet Gynecol.* 1983; 345:78-86.
- Mazzali M, Hughes J, Kim YG:** Elevated uric acid increases blood pressure in the rat by a novel crystal - independent mechanism. *Hypertension*; 2001; 38: 1101-6.
- McMahon MJ, Ananth CV and Liston RM:** Gestational diabetes mellitus. Risk factors, Obstetric complications and infant outcomes. *J Reprod Med* 1991; 165: 1711-1721.
- Merviel P, Ba R, Beaufils M, Breart G, Salat-Baroux J and Uzan S :** Lone hyperuricemia during pregnancy: Maternal and fetal outcomes. *Eur. J. Obstet. Gynecol. Reprod. Biology*; 1998; 77(2): 145-50.
- Meye NL, Mercer BM, Friedman SA and Sibai BM:** Urinary dipstick protein; A poor predictor of absent or severe proteinuria. *Am J Obstet Gynecol.* 1998; 43: 392-406.
- Mikhail MS, Anyaegbunam A, Garfinkel D, Palan PR and Basu J:** Reduced ascorbic acid, alpha tocopherol and beta carotene in women with preeclampsia. *Am J Obstt Gynecol* 1994; 170: 137-141.
- Millar JGB, Campbell SK, Albano JDM, Higgins BR and Clark AD:** Early prediction of preeclampsia by measurement of kallikrein and creatinine on a random urine sample. *Br J Obsret Gynecol* 1994; 171: 150-157.

- Minamiyama T, Takemura S and Inoue M:** Effect of thiol status on nitric oxide metabolism and circulation. *Arch Biochem Biophys.*; 1997; 341:186-92.
- Mitchell MD and Koenig JM:** Increased production of 15-hydroxyeicosatetraenoic acid by placentae from pregnancies complicated by pregnancy-induced hypertension. *AmJ obstet Gynecol.* 1996; 103: 421-430.
- Mittendorf R, Iain KY Wilkams MA and Walker CK:** Preeclampsia case control study of risk factors and the interactions. *J Reprod Med* 1991; 43:61-68.
- Morris MC, Twickler DM, Hatab MR, Clarker GD, Peshok RM and Cunningham FG:** Cerebral blood flow and cranial magnetic resonance imaging in eclampsia and severe preeclampsia. *Obstet Gynecol* 1996; 41: 491-496.
- Morriss NH, Eaton BM and Dekker G:** Nitric Oxide the endothelium pregnancy and preeclampsia. *BrJ Obstet Gynecol.* 1997; 76:663-669.
- Murai JT, Muzykanskiy E and Tay Lor RN:** Maternal and fetal modulators of lipid metabolism correlate with the development of preeclampsia. *Metabolism* 1996; 103: 4-12.
- Mustard JF, Murphy EA, Ogrylo MA and Smythe, HA:** Blood coagulation and platelet economy in subjects with primary gout. *J. Can. Med. Assoc.*; 1963; 89: 1207-11.
- Muttukrishna S, Knight PG, Groome NP, Redman CW and Ledger WL:** Activin-A and inhibin-A as possible endocrine markers for preeclampsia. *Lancel* 1997; 46: 963- 967.
- Myatt L, Brewer A and Brockman DE:** The action of nitric oxide in the perfused human fetal placental circulation. *Am J Obstet Gynecol.* 1997; 349: 1285-1289.
- Myatt L, Brewer AS, Langdon G and Brockman DE:** Attenuation of the vasoconstrictor effects of thromboxane and endothelin by nitric oxide in the human fetal-placental circulation. *Am J Obstet gynecol* 1991; 164: 687-692.
- Myatt L:** Vaso-active factors in pregnancy. *Fetal and Matrnal Medicine Review* 1992; 166:224

- Naidu K, Moodley J and Corr P:** Single photon emission and cerebral computerized tomographic scan and transcranial Doppler sonographic findings in eclampsia. *Br J. Obstet Gynecol.* 1992; 2: 15-36.
- National High Blood Pressure Education Program, (NHBPEP):** Working group report on high blood pressure in pregnancy. *Am J Obstet Gynecol;* 2000; 183: 51-22.
- Neerhof MG:** Pregnancy in the chronically hypertensive patient. *Clinic perinatol* 1997; 104: 1165-1172.
- Nisell H, Carlstrom J, Cizinsky, S, Grunewald C, Nylund L and randma I:** Atrial natriuretic peptide concentrations and hemodynamic effects of acute plasma volume expansion in normal pregnancy and preeclampsia. *Obstet Gynecol*, 1997; 24: 391-406.
- Nolan TE, Smith RP and devoe LD:** Maternal plasma D- dimer levels in normal and complicated pregnancies. *Obstet Gynecol.* 1992; 79: 902-915.
- North RA, Taylor RS and Schellengberg JC:** Evaluation of a definition of preeclampsia. *Br J Obstet Gynecol* 1999; 81: 235-240.
- Nova A. Sibai BM, Barton JR, Mercer BM and Mitchell MD:** Maternal plasma level of endothelin is increased in preeclampsia. *Am J Obstet Gynecol* 1999; 106: 767-776.
- Obrien WF:** predicting preeclampsia. *Obstet Gynecol* 1994; 165: 724-736.
- Oian P, Maltau JM and Noddeland H:** Transcapillary fluid balance in preeclampsia. *Br J. Obstet Gynecol.* 2001; 75: 445-451.
- Oosterhof H; Voorhoeve PG and Aamoudse JG:** Enhancement of hepatic artery resistance to blood flow in preeclampsia in presence or absence of HELLP syndrome (hemolysis, elevated liver enzymes, and low platelets). *Am J Obstet Gynecol.* 1986; 93-235-239.
- Paarlberg KM, DeJong CLD, Van Geijn HP, van Kamp GJ, Heinen AG and Dekker GA:** Total plasma fibronectin as a marker of pregnancy-induced hypertensive disorders: Alongitudinal study. *Obstet Gynecol* 1994; 171: 526-534.

**Palmer FMJ, Ashton DS and Moncada S:** Vascular endothelial cells synthesize nitric oxide from L-arginin, *Nature, AmJ obstet Gynecol 1999; 16:76-79.*

**Pelle G. Lindovist and Karel Marsal:** Moderate smoking during pregnancy is associated with a reduced risk of preeclampsia. *Acta Obstet Gynecol Scand 1988; 333: 664-672.*

**Petraglia F, Florio P, Benedetto C, Gallo C, Woods RJ, Genazzani AR and Lowry PJ:** High levels of corticotropin-releasing factor (CRF) are inversely correlated with low levels of maternal CRF-binding protein in pregnant women with pregnancy-induced hypertension. *J clin Endocrinol metab 1999; 78: 693-697.*

**Pickele C.J., Symonds E.M and Broughton Pipkin:** Fetal outcome in a randomized controlled trial of labetalol versus placebo in P.I.H.Br.J.Obstet Gynecol 1996; *81: 852-857.*

**Poul W. Whitecar, Kim A Boggess, John Mthrop and Douglas D.Taylor:** Altered expression of TCR- CD3 $\xi$  induced by sera from women with preeclampsia. *AmJ. Obst. Gynecol 1996; 15: 7-23.*

**Powers RW. Evans RW. Ness RB, Crombleholme WR and Roberts JM:** Homocysteine is increased in preeclampsia but not in gestational hypertension. *J Sur Gyecol Investig 2001; 185: 812-818.*

**Pritchard JA and Cunningham FG and Mason RA:** Coagulation changes in eclampsia; their frequency and pathogenesis. *Am J Obstet Gynecol. 2000; (Abstr 375). 7 (1) (Suppl).*

**Qureshi AL, Frankel MR and Ottenlips JR:** Cerebral hemodynamics in preeclampsia and eclampsia. *Arch Neurol. 1976; 124: 855-861.*

**R.A. Fay:** Uric acid in pregnancy and preeclampsia: An alternative hypothesis. *Aust. NZ J. Obstet. Gynecol.; 1990; 30: 141-2.*

**Railton A and Allen DG:** Management and outcome of pregnancy complicated by severe preeclampsia of early onset. *Med.J 1996; 53: 1226-1231.*

**Raps EC, Galetta SL and Broderick M:** Delayed peripartum vasculopathy: Cerebral eclampsia revisited. *Ann Neurol. 1997; 72:608-610.*

**Rayman MP, Bode P and Redman CW:** Low selenium status is associated with the occurrence of the pregnancy disease preeclampsia in women from the United Kingdom. *Am. J. Obstet. Gynecol.*; 2003; 189: 1343-9.

**Redman CW, Williams GF, Jones DD and Wilkinson RH:** Plasma urate and serum deoxycytidylate deaminase measurements for the early diagnosis of preeclampsia. *Br J Obstet Gynaecol*; 1977; 84: 904-8.

**Redman CWG, Sacks GP and Sargent IL:** Preeclampsia an excessive maternal inflammatory response to pregnancy. *AM J Obstet Gynecol*. 1993; 33: 222-225.

**Regan CL and Fitzgerald DJ:** Altered isoprostane to prostocyclin ratio in preeclampsia. *Am J Obstet Gynecol* 1999; 108: 499-503.

**Riedel H, Bahlmann J and Eisenbach GM:** Results of a prospective study of toxemia in pregnancy. *Contrib. Nephrol.* 1981; 25: 137-44.

**Riely CA, Latham PS and Romero R:** Acute fatty liver of pregnancy. *Ann Intern Med*. 1997; 176:510-522.

**Roberts J M:** Preeclampsia not simply pregnancy induced hypertension. *Hosp Pract* 2001; 341: 1447-1451.

**Roberts JM:** Preeclampsia: what we know and what we do not know. *Semin Perinatol.*; 2000; 24:24-8.

**Roberts JM, Taylor RN, Musci TJ, Rodgers GM, Hubel CA and Mc Laughlin MK:** Preeclampsia: an endothelial cell disorder. *Am. J. Obstet. Gynecol.*; 1989; 161: 1200-4.

**Roberts JM:** Endothelial dysfunction in preeclampsia. *Semin Repord Endocrinol*. 1995; 30: 25-33.

**Robertson WB and Khong TY:** Pathophysiology of uteroplacental bed In Sharp F, Symonds EM, edios. *Hypertension in pregnancy*. Ithaca (NY); Perinatology Press 2001; 16:5-15.

- Robson, Davison J and Baylis C:** Hypertension and renal disease .cli obstet gynecol; 1999; 8,353-373.
- Rodgers GM, Taylor RN and Roberts JM:** Preeclampsia is associated with a serum factor cytotoxic to human endothelial cells. Am. J. Obstet. Gynecol.; 1988; 59: 908-14.
- Saloner ROS H, Cnattingius S and Lipworth L:** Comparison of risk factors cohort study. AmJ Epidemiol 1987; 66: 101-18.
- Samuels P, Main EK, Tomaski A, Mennuti MT, Gabbe SG and Cines DB:** Abnormalities in platelet antiglobulin tests in preeclamptic mothers and their neonates. Am J Obstet Gyncol 1998; 147-1062-1070.
- Samuels P, Main EK, Tomaski A, Mennuti MT, Gabbe SG and Cines DB:** Abnormalities in platelet antiglobulin tests in preeclamptic mothers and their neonates. Am J Obstet Gynecol 1987; 157: 109-115.
- Sanchez Ramos L, Briones DK and Kaunitz AM:** Prevention of P.I.H. by calcium supplementation in angiotension II secretive patient. Obstet Gynecol 1987; 157: 109-113.
- Sanchez Ramos L, Jones DC and Cullen MT:** Urinary calcium as an early marker for preeclampsia. Obstet Gynecol 1994; 84: 349-53.
- Sanchez-Lozada LG, Tapia E, Avila-Casado C:** Mild hyperuricemia induces glomerular hypertension in normal rats. Am. J. Physiol. Renal Physiol.; 2002; 283: 51105-10.
- Sanchez-Ramos L, Jones DC and Cullen MT:** Urinary calcium as an early marker for preeclampsia. Obstet Gynecol 1991; 77: 685-693.
- Santos CX, Anjos EI and Augusto O:** Uric acid oxidation by peroxynitrite: multiple reactions, free radical formation, and amplification of lipid oxidation. Arch. Biochem. Biophys. 1999; 372: 285-94.
- Sargent I L:** Maternal and fetal immune responses during pregnancy. Exp Clin Immunoglobulins 1991; 77: 685-692.

**Sastry BV, Hemontolor ME and Olenick M:** Prostaglandin E2 in human placenta; Its vascular effects and activation of prostaglandin E2 formation by nicotine and cotinine. *Pharmacology* 1999; *10*: 95-102.

**Sattar N, GAW A, Packard Gj and Greer IA:** Potential pathogenic roles of aberrant liporprotein and fatty acid metabolism in preeclampsia. *Br J Obstet Gynecol* 1999; *58*: 70-86.

**Saudan P, Brown MA, Buddle ML and Jones M:** Does gestational hypertension become preeclampsia? *Br J Obstet Gynecol*; 1998; *105*: 1177-84.

**Saudan P, Brown MA, Buddle ML and Jones M:** Does gestational hypertension become preeclampsia? *Br J Obstet Gynecol* 1996; *103*: 614-620.

**Schackis RC:** Hyperuricaemia and preeclampsia: is there a pathogenic link? *Medical Hypotheses*. 2004; *Sincedirect.com*

**Scholtes MCW, Geretsen G and Hoak HI:** Factor VIII ration in normal and pathological pregnancy. *Europ J Obstet Gynecol* 1998; *105*: 1177-1184.

**Sen C, Madazh R, Kavuzlu C, Ocak V and Tolun N:** The value of antithrombin-III and fibronectin in hypertensive disorders of pregnancy. *J Perinat Med*. 1983; *723*:54-60.

**Seshadri I and Venkataraman I:** Hypertension in pregnancy; outcomes proteinuric and nonproteinuric. *J Reprod Med* 1994; *765*:22-29.

**Sheehan HL and Lynch JB:** Cerebra lesions In; *Pathology of Toxemia of Pregnancy*. Baltimore, Villiams & Wilkins, 1997; *42*:88-90.

**Sibai BM:** Diagnosis and management of gestational hypertension and preeclampsia. *Obstet Gynecol*; 2003; *102*(1): 181-192.

**Sibai BM, Caritis S, Hawth J, Lindheimer MD, MacPherson C and Klebanoff M:** Hypertensive disorders in twin versus singleton gestations. National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. *Am J Obstet Gynecol*; 2000; *182*: 938-42.

- Sibai BM, Mercer BM, Schiff E and Friedman SA:** Aggressive versus expectant management of severe preeclampsia at 28 to 32 weeks gestation a randomized controlled trial. *Am J Obstet Gynecol* 1973; *179*:877-883.
- Sibai BM, Taslimi MM and El- Nazer A:** Maternal perinatal outcome associated with the syndrome of hemolysis. Elevated liver enzymes, and low platelets in severe preeclampsia-eclampsia. *AM J Obstet Gynecol*. 1995; *171*: 818 -822.
- Sibai BM:** Thrombophilias and adverse outcomes of pregnancy-what should a clinician do? (Editorial) *N Engl J Med* 1986; *155*: 501-509.
- Silver HM, Seebeck M and Carlson R:** Comparison of total blood volume in normal preeclamptic and nonproteinuric gestational hypertensive pregnancy by simultaneous measurement of red blood cell and plasma volumes. *AM J obstet Gynecol*. 1999; *340*: 50-55.
- Spellacy WN, Hander A and Ferre CD:** A case control study of 1253 twin pregnancies. *Obstet Gynecol* 1999; *179*: 87-93.
- Spencer JAD, Smith MJ and Cederholm- Williams:** Influence of preeclampsia on concentration of haemostatic factors in mothers and infants. 1999; *98*:122-134.
- Srivastava ED, Hallet MB and Rhodes J:** Effect of nicotine and cotinine on the production of oxygen free radicals by neutrophils in smokers and nonsmokers. *Hum Toxicol Am J Obstet Gynecol* 1983; *34*:876-880.
- Staff AC, Ranheim T, Koury J and Henriksen T:** Increased Contents of phospholipids, cholesterol, and lipid peroxides in deciduas basalis in women with preeclampsia. *Am J Obstet Gynecol* 1989; *8*: 461-463.
- Stamilio DM, Sehder HM, Morgan MA, Probert K and Macones GA:** Can antenatal clinical and biochemical markers predict the development of severe preeclampsia? *Am J Obstet Gynecol* 1999; *180*:587-598.
- Starzyk KA, Salafia CM and Pezzullo JC:** Quantitative differences in arterial morphometry define the placental bed in preeclampsia. *Human Pathology*. 2000; *182*:589-597.

**Taufield PA, Alas KL and Rensick LM:** Hypocalciuria in preeclampsia. *Am J Obstet Gynecol* 1997; 28: 353-358.

**Taylor RN and Roberts JM:** Pathogenesis of preeclampsia, endothelial cell dysfunction. In Lindheimer MD, Roberts JM, Cunningham FG, editors; Chesley's hypertensive disorders in pregnancy. Stanford, CT, Appleton & Lange, 1998; 16: 17-31.

**Taylor RN, Groot CJ and Cho UK:** Circulating factors as markers and mediators of endothelial cell dysfunction in preeclampsia. *Semin Reprod Endocrinol.* 1987; 316: 715-718.

**Torres PJ, Escobar G, Palacio M, Gratacos E, Alonso PL and Ordinas A:** Platelet sensitivity to prostaglandin E, inhibition is reduced in preeclampsia but not in nonproteinuric gestational hypertension. *Br J Obstet Gynecol* 1999; 19: 344-354.

**Tranquilli AL, Giannubilo SL, Dell'Uomo B and Corradetti A:** Prediction of gestational hypertension or intrauterine fetal growth restriction by mid-trimester 24-h ambulatory blood pressure monitoring. *Inter. J. Gynecol. Obstet.*; 2004; 85: 126-31

**Tyurin VA, Liu SX, Tyurina YY, Sussman NB, Hubel CA, Roberts JM, Taylor RN, Kagan VE:** Elevated Levels of S-Nitrosoalbumin in Preeclampsia Plasma. *Circulation Research*; 2001; 88: 1210-5

**Walker JJ, Cameron AD, Bjornsson S, Singer CR and Fraser C:** Can platelet volume predict progressive hypertensive disease in pregnancy? *Am J Obstet Gynecol* 1987; 103: 19-24.

**Walsh SW:** Maternal placental interactions of oxidative stress and antioxidants in preeclampsia. *Semin Reprod Endocrinol.* 1989; 332: 177-188.

**Walsh SW:** Low dose aspirin, treatment for imbalance of increased thromboxane and decreased prostacyclin in preeclampsia. *Am J Perinat* 1998; 16: 93-104.

**Walsh SW:** Preeclampsia: An imbalance in placental prostacyclin and thromboxane production. *Am J Obstet Gynecol*, 1989; 196: 11-17.

**Walsh SW:** Lipid peroxidation in pregnancy. *Hypertens Pregn* 1985; 152: 335.

- Wang Y, Walsh SW, Guo J and Zhang J:** The imbalance between thromoxane and prostacyclin in preeclampsia is associated with an imbalance between lipid peroxides and vitamin E in maternal blood. *Am J Obstet Gynecol* 1985;152: 335-340.
- Ward PA:** Mechanisms of endothelial cell killing by H<sub>2</sub>O<sub>2</sub> or products of activated neutrophils. *AmJ Med.* 1991; *a*; 165: 1695-1703.
- Weinstein L:** Syndrome of hemolysis elevated liver enzymes, and low platelet count a severe consequence of hypertension in pregnancy. *Am. J. Obstet. Gynecol.* 1991; 91: 895-898.
- Weir RJ, Fraser R, Lever AF, Morton JJ, Brown JJ, Krazewski A, McIllevine GM, Robertson JIS and Tree M:** Plasma renin, substrate, angiotensin II, and aldosterone in hypertensive disease of pregnancy. *Lancet*, 1982;142: 159-167.
- Wells M and Bulmer J N:** The human placental bed; histology, immunocytochemistry and pathology *Histopathology*. *AmJ Obstet and Gynecol* 1973; 1: 291-296.
- Wergeland E and Strand K:** Work pace control and pregnancy health in a population – based sample of employed women. *Br J Obstet Gynecol* 1988; 13: 483-498.
- Winocour PD, Turner MR, Taylor TG and Munday KA :** Platelet aggregation in rats in relation to hyperuricemia induced by dietary single-cell protein and to protein deficiency. *Thromb. Haemost.* 1978; 39 (1978), pp. 346-359.
- Witlin AG, Saade GR, Matter F and Sibai BM :** Predictors of neonatal outcome in women with severe preeclampsia or eclampsia between 20 and 33 weeks gestation. *Am J Obstet Gynecol* 2002; 41: 34-35.
- Ylikorkala O and Makela UM:** Prostacyclin and thromboxane in Gynecology and obstetrics. *AmJ Obstet Gynecol* 2002; 182: 607-611.
- Zeek PM and Assali NS:** Vascular changes in deciduas associated with eclamptogenic toxemia of pregnancy. *Am J Clin Pa thol* 1985; 152: 318-329.

**Zinaman M, Rubin J and Lindheimer MD:** serial plasma oncotic pressure levels and echoencephalography during and after delivery in sever preeclampsia. *Lancet*, 1992; 20: 1099-1102.