

REFERENCES

- 1- **Abraham, G.E. (1977):**
Metabolism of testosterone, J. Clin. Invest., 48 : 696.
- 2- **Ahmad, S. (1991):**
Enalapril and reversible alopecia. Arch. Intern. Med., 151 : 404.
- 3- **Arndt, G.J. (1987):**
Jojoba, Cosm. Toil., 102 : 66.
- 4- **Auerbach, R. and Orentlich, N. (1968):**
Alopecia and ichthyosis secondary to allopurinol, Arch. Dermatol., 98 : 104.
- 5- **Azmy, N.H.; Sallam, J.H.; El-Sheimy, S. and Madour, T.M. (1984):**
Iron deficiency as a cause of chronic diffuse falling of hair. Egypt. J. Dermat. Vener, 4 : 135.
- 6- **Backburn, G.L.; Bistran, B.R. and Hoag, C. (1976):**
Hair loss with rapid weight loss.
JAMA. 236 : 252.
- 7- **Bailey, M.A. (1969):**
Effect of malnutrition on hair shaft. In : Hair growth, Montagna, W. and Dobson, R. Pergamon, Oxford, P : 109.
- 8- **Bardelli, A. (1989):**
Telogen effluvium and minoxidil. J. Am. Acad. Dermatol., 21 : 572.
- 9- **Barman, J.M.; Osborne, D. and Dobson, R.I. (1969):**
Embryology of the skin. In : Advances in Biology of Skin, Montagna, W. and Dobson, R.I. Pergamon, Oxford, P : 211.
- 10- **Barth, J.H.; Cherry, C.A. and Wojnarowska, F. (1989):**
Spironolactone is an effective and well tolerated systemic antiandrogen therapy for hirsute women. J. Clin. Endocrinol. Metab., 68 : 966.

- 11- Benedict, L.M.; Abell, E. and Jegasothy, B. (1991):**
Telogen effluvium associated with eosinophilia myalgia syndrome.
J. Am. Acad. Dermatol., 25 : 112.
- 12- Bergfield, W.F. (1987):**
Androgenic alopecia, Dermatol. Clin., 5, 3 : 491.
- 13- Bernstein, G.M. (1988):**
Postfebrile telogen effluvium in critically ill patients, Crit - Care Med., 16 : 98.
- 14- Bertolino (1994):**
Differential expression of type I hair keratins. J. Invest. Dermatol., 94 : 297.
- 15- Bloomer, H.A. and Kyrle, L.H. (1959):**
Myxedema, a reevaluation of clinical diagnosis is based on eight cases. Arch. Int. Med., 104 : 234.
- 16- Blum, I. (1980):**
Increased hair loss as a side effect of bromocriptine treatment. N. Engl. J. Med., 303 : 1418.
- 17- Buhl, A.E. (1990):**
Minoxidil sulfate is the active metabolite that stimulate hair follicles. J. Invest. Dermatol., 95 (5) : 557.
- 18- Bradfield, R.B. (1969):**
Hair root adaptation to marasmus in Andean Indian children. Lancet, 2 : 1395.
- 19- Britton, K.E.; Valerie, B.L. and Ekins, R.P. (1975):**
A strategy for thyroid function tests. Br. Med. J., 350 : 25.
- 20- Brodin, M.B. (1987):**
Drug - related alopecia. Dermatol. Clin., 5 (3) : 571.
- 21- Brown, T.D. (1987):**
Aphase I clinical trial of recombinant DNA gamma interferone. J. Clin. Oncol., 5 : 790.

- 22- **Callen, J.P. (1995):**
Hair disorders in systemic disease. In : Dermatologic Signs of Internal Disease. Jorizzo, J.L. Greer, K.E. (eds). JB Lippincott. Philadelphia, P : 1.
- 23- **Caserio, R.J. (1987):**
Diagnostic techniques for hair disorders : clinical hair manipulations and clinical findings. *Cutis*, 40 : 442 -8.
- 24- **Cavallo, E. (1990):**
Resting metabolic rate, body composition and thyroid hormones. *Horm. Metab. Res.*, 22 : 632.
- 25- **Champan, R.S. (1967):**
Diffuse thinning of hair in iodide - induced hypothyroidism. *Br. J. Dermatol.*, 79 : 103.
- 26- **Chan - Lam, D.; Fitzsimons, E.J. and Douglas, W.S. (1987):**
Alopecia after immunoglobulin infusion. *Lancet*, 1 : 1436.
- 27- **Ciotta, L.; Cianci, A. and Merletta, E. (1994):**
Treatment of hirsutism with flutamide and low - dosage oral contraceptive in polycystic ovarian disease patients. *Fertile. Steril.*, 62 : 1129.
- 28- **Comaish, J.S. (1971):**
Metabolic disorder and hair growth. *Br. J. Dermatol.*, 84 : 83.
- 29- **Comaish, T.S. (1985):**
The thyroid and hair growth, *semin. Dermatol.*, 4 : 4.
- 30- **Corea, L. and Lupattelli, L. (1972):**
Studies in hair growth in disorders of parathyroid gland, *Folia endocrin*, 25 : 347.
- 31- **Cormia, F.E. (1967):**
Alopecia from oral contraceptives. *JAMA*, 201 : 141.
- 32- **Cornin, E. (1980):**
Hair tonics, stimulants and restorers. In : Contact dermatitis, Cornin, E. (ed.) 1st edition. Churchill livingstone Edinburgh, P : 132.

33- Cotterill, J.A. (1981):

Dermatological non disease : a common and potentially fatal disturbance of cutaneous body image. *B. J. Dermatol.*, 104 : 661.

34- Cotterill, P.C. and Unger, W.P. (1992):

Hair transplantation in females. *J. Dermatol., Surg. Oncol.*, 6 : 477.

35- Cusan, L.; Dupont, A. and Gomez, J. (1994):

Comparison of flutamide and spironolactone in the treatment of hirsutism. *Fertile. Steril.*, 61 : 281.

36- Dahlin, P.A. (1984):

Telogen effluvium - Hair loss after spinal cord injury. *Arch. Phys. Med. Rehab.*, 65 : 485.

37- Daniel, W.W. (1987) :

Biostatistics. Wayne W. Daniel, John Wiley Sons, In : A foundation for analysis in the health sciences. 4th edition New York, Chichester, Brisbane, Toronto, P : 531.

38- Dawber, R. (1971):

Frequency, hair loss and the pill, *Br. Med. J.*, 4 : 234.

39- Dawber, R. P. (1981):

Common baldness in women. *Int. J. Dermatol.*, 20 : 647.

40- Demis, D.J. (1991):

Introduction to disease of the hair In : *Clinical Dermatology*, Demis D. J. (ed). J.B. Lippincott, New York. P : 1.

41- Demis, D.J.; Thiers, B.H. and Smith, E.B. (1991):

Schematic diagram of an anagen hair, *Clin. Dermatol.* Philadelphia, J.B. Lippincott : 3.

42- Detwiler, S.P.; Carson, J.L. and Woosley, J.T. (1994):

Bubble hair. *J. Am. Acad. Dermatol.*, 121 : 197.

43- DeVillez, R.L. and Dunn, J. (1986):

Female androgenetic alopecia - the 3 α , 17 B and rostandiol, G/SHBG ratio as a possible marker for female pattern baldness. *Arch. Dermatol.*, 122 : 1011.

- 44- Dinarello, C.A. (1988):**
New concepts on the pathogenesis of fever, *Rev. Infect. Res.*, 10 : 168.
- 45- Dupré, A. Bonafé, J. and Carriere, J. (1979):**
The hair in acrodermatitis enteropathica A disease indicator ?
Acta. Dermatol. Venereol., 59 : 177.
- 46- Durr, N. (1982):**
Biology of scalp hair growth. *Clin. Plast. Surg.* 9 : 1970.
- 47- Dworkin, R.H. (1981):**
Fixed Oils. *Lancet.*, 1 : 1153.
- 48- Ebling, F. (1987):**
The biology of hair. *Dermatol. Clin.*, 5 (3), 479.
- 49- Eckert, J. (1975):**
Diffuse hair loss and psychiatric disturbance. *Acta Dermatol. Venereol.*, 55 : 147.
- 50- Eckert, J. (1985):**
Diffuse hair loss and psychiatric disturbance. *Acta. Dermatol. Venereol.*, 55 : 147.
- 51- Eckert, J.; Church, R.; Ebling, F. and Munro, D.S. (1967):**
Hair loss in women. *Br. J. Dermatol.*, 79 : 543.
- 52- Ehrmann, D.A. and Rosenfeld, R.I. (1990):**
Hirsutism beyond the steroidogenic block. *N. Engl. Med.*, 323 : 909.
- 53- Erenus, M. Gurbuz, O.; Durmusoglu, F. (1994):**
Comparison of the efficacy of spironolactone versus flutamide in the treatment of hirsutism. *Fertile. Steril.*, 61 : 613.
- 54- Famuyiwa, O.O. and Bella, A.F. (1990):**
Thyrotoxicosis in Nigeria. *Trop. Geogr. Med.*, 42 : 248.
- 55- Feingold, K. and Elies, P. (1987):**
Endocrine - skin interactions. *J. Am. Acad. Dermatol.*, 17 : 921.

56- Fich, A. and Eliakim, R. (1988):

Doses sulfasalazine induce alopecia ? J. Clin. Gastroenterol., 10 : 466.

57- Field, L.M. (1980) :

Toxic alopecia caused by pyridostigmine bromide. Arch. Dermatol., 116 : 1103.

58- Fiedler, V.C. (1988):

Amphetamines and hair loss. JAMA, 260 (2) : 184.

59- Fiedler, V.C. (1994):

Diffuse alopecia, In : Disorders of Hair Growth, Olsen, E.A., McGraw - Hill, New York, P : 243 - 49.

60- Fiedler, V.C. and Abdul Hafeez (1994):

Telogen Hair loss, In : Disorders of hair growth, Olsen, E.A. McGraw - Hill, New York, P : 250.

61- Flesh, P. (1963):

Inhibition of keratinizing structures by systemic drugs Pharmacol. Rev., 15 : 653.

62- Fox, C. (1985):

Hair products patent review and update. Cosm. Toil, 100 : 57.

63- Fox, C. (1993) :

Hair care : literature and patent review, cosm. Toil., 108 : 29.

64- Fraunfelder, F.T.; Meyer, S.M. and Mcrackar, S. J. (1990):

Alopecia possibly secondary to topical ophthalmic B - blockers JAMA, 263 (11) : 1493.

65- Freedberg, I.M. (1971):

Skin. In : Review of Body System. Werner S.C. and Ingbore, S.H. (eds). 3rd edit., Harpar and Row publishers. New York, P : 722.

66- Freeiman, D. (1971):

Alopecia In : Human Hair Growth In heath And Disease, Freeiman, D. (Ed). Chales, C. Publ., Bannerstone House, U.S.A., P : 206.

- 67- Freinkel, P. and Freinkel, N. (1972):**
Hair growth and alopecia in hypothyroidism. Arch. Dermatol., 166 : 349.
- 68- Freinkel, P. and Freinkel, N. (1987):**
Cutaneous manifestations of endocrine disorders, In : Dermatology In General Medicine, Fitzpatrick, T. Eisen, A. and Wolff, K. (Eds). McGraw - Hill, New York, P : 2063.
- 69- Garcia - Muret, M.P.; Sitgas, D.; Tuneu, L. and DeMoragas, J.M. (1990):**
Telogen effluvium associated with albendazole therapy. Int. J. Dermatol., 29 : 669.
- 70- Gauci, L. (1980):**
Changes in hair pigmentation associated with cancer chemotherapy. Cancer Treat. Rep., 64 : 193.
- 71- Giacometti, L. (1965):**
In : Advances in Biology of Skin. Quoted from Textbook of Dermatology, Rook, A. Wilkinson, D.S. and Ebling, F.J.(eds) Blackwell scientific publication, Oxford, P : 1734.
- 72- Giacometti, L. and Montagna, W. (1969):**
The innervation of human hair follicles. Montagna W. and Richard L. (Eds) In : Advances In Biology of Skin, Vol. IX, Pergamon, Oxford : 393.
- 73- Gorez, G. (1990):**
Cicatricial alopecias. In : Hair And Hair Diseases. Orfanos, C. and Happle, R (eds). Springer - Verlag, Berlin, P : 611.
- 74- Graves, K.; Kestenbaum, T. and Kalivas, J. (1980) :**
Hereditary acrodermatitis enteropathica in an adult. Arch. Dermatol., 116 : 562.
- 75- Griffiths, W.A. (1973):**
Diffuse hair loss and oral contraceptives. Br. J. Dermatol., 88 : 31.
- 76- Grossman, K.L. (1994):**
Anagen hair loss. In : Disorders of Hair Growth. Olsen, E.A., McGraw-Hill, New York, P : 225.

77- Gucklhorn, I.R. (1970):

A review of antimicrobial agents, including cetrimide used in cosmetics. *Mfg, Chem.*, 41 : 28.

78- Hale, P. (1983):

Hormones and hair growth. *Dermatol. Clin.*, 5 : 479.

79- Hamilton, J.B. (1951):

Patterned loss of hair in man : Types and incidence *Ann. New York. Acad. Sc.*, 53 : 708.

80- Hansen, L.S.; Coggle, J.E. and Wells, J. (1984):

The influence of the hair cycle on the thickness of mouse skin. *Anat. Rec.* 210 : 573.

81- Hansen, S.; Kuylen, K. and Bjakam, A. (1989):

Hair loss and scaling with proguanil, *Lancet*, 1 : 225.

82- Hashimoto, K. (1988) :

The structure of human hair. *Clin. Dermatol.*, 6 : 21.

83- Hard, S. (1963):

Non anemic iron deficiency as an etiological factor in diffuse loss of hair of the scalp in women. *Acta. Dermatol. Venereol.*, 43 : 562.

84- Harris, K.A. (1990):

Human hair follicle aromatase activity in females with androgenetic alopecia. *J. Invest. Dermatol.*, 94 : 575.

85- Harry, R. (1962):

Hair tonics and lotions. In : *The Principles And Practice of Modern Cosmetics.* Wilkinosn, T.B. and Mclaughling T.P. (eds). Fifth edition, Leonard Hill Limited P : 340.

86- Hassett, J.M. (1988) :

Post - febrile telogen effluvium in critically ill patients. *Crit care Med.*, 16 (1) : 98.

87- Headington, J.T. (1988) :

Androgenetic alopecia, histologic studies of human salp. *Clin. Dermatol.*, 6 : 188.

88- Heimer, W.L. (1993):

Evaluation of hair loss. J. Am. Acad. Dermatol., 28 : 669.

89- Heymann, W. (1992):

Cutaneous manifestations of thyroid disease. J. Am. Acad. Dermatol., 26, 885.

90- Hong, L.S. (1988):

3, B-Hydroxysteroid dehydrogenase activity in sebaceous glands of scalp of male pattern baldness. J. Invest. Dermatol., 91 : 105.

91- Hordinsky, M.K. (1987):

General evaluation of the patient with alopecia, Dermatol. Clin., 5 (3) : 483 -9.

92- Hordinsky, M.K. (1993):

The antiandrogens, when and how they should be used. Dermatol. Clin, 1 : 65.

93- Hori, Y. (1976) :

Alopecia in hypothyroidism. In : Biology of Diseases of Hair, Kobori, T. Montagna, W. (eds), University of Tokyoopress, Tokyo, P : 279.

94- Hunt, S.J.; Caserio, R.J. and Abell, E. (1991):

Primary systemic amyloidosis causing diffuse alopecia by telogen arrest. Arch. Dermatol., 127 : 1067.

95- Inaba, M. (1990):

Male pattern baldness : sebaceous gland hypothesis. Cosm. Toil. 105 : 77.

96- Itami, S.; Kurata, S. and Sonoda, T. (1991):

Characterization of 5 a - reductase in cultured human dermal papilla cells from beard and occipital hair. J. invest. Dermatol., 96 : 57.

97- Jayne, C. (1991):

Bonding techniques. In : Hair Additions : The fourth dimension, Frangie, C (ed). Milady publishing, New York, P : 127.

98- Jayne, C. (1991):

Braiding and sewing tracks. In : Hair Additions : the fourth dimension. Frangie, C. (ed), 1st edit., Milady publishing, New York, P : 88.

99- Jayne, C. (1991):

Introduction. In : Hair Additions : Frangie, C (ed), 1st edit., Milady publishing, New York, P : 24.

100- Jee, J.Y. and Hsu, M.I. (1991):

Alopecia syphilitica, a simulator of alopecia areata : hisopathology and differential diagnosis. J. Cutan. Pathol., 18 (2) : 97.

101- Jelinek, J.E. (1970):

Cutaneous side - effects of oral contraceptives.
Arch. Dermatol., 101 : 181.

102- Johnson, A.; Latham, M. and Roe, D. (1976):

An evaluation of the use of changes in hair root morphology in the assessment of protein - calorie malnutrition. Am. J. Clin. Nutr., 29 : 502.

103- Johnson, W. (1975):

Chronic diffuse alopecia. Br. J. Dermatol., 77 : 115.

104- Katz, H.I. (1987):

Long term efficacy of topical minoxidil in male pattern baldness.
J. Am. Acad. Dermatol., 16 : 711.

105- Karyn, L.; Grossman and Joseph, C. (1994) :

Anagen hair loss In : Disorders of Hair Growth, Olsen, E.A., McGraw - Hill New York, P : 223.

106- Kaufman, J. P. (1976):

Teolgen effluvium secondary to starvation diet. Arch. Dermatol., 112 : 731 : 731.

107- Kessels, A.G.; Cardynalas, R.L. and Rogres, R. (1991) :

The effectiveness of the hair restorer "DABAO" in males with alopecia androgenetica : A clinical experiment. J. Clinical Epidemiology, 44 : 439.

108- Kollar, E. (1970):

The induction of hair follicles by embryonic dermal papillae. J. Invest. Dermatol., 55 : 374.

109- Kopf, A. and Orentreich, N. (1957):

Alkaline phosphatase in alopecia areata. Arch. Dermatol., 76 : 288.

110- Klein, A.W.; Rudolph, R.I. and Leyden, J.J. (1973):

Telogen effluvium as a sign of Hodgkin disease. Arch Dermatol., 108 : 702.

111- Kleniewska, D. (1970):

Rosercinol. Br. J. Dermatol., 83 : 543.

112- Kligman, A.M. (1961):

Pathologic dynamics of human hair loss, Arch. Dermatol., 83 : 175 - 198.

113- Kuster, W. and Happle, R. (1984):

The inheritance of common baldness. J. Am. Dermatol., 11 : 921.

114- Laskey, J.; Webb, I. And Schulman, H.M. (1988):

Evidence that transferrin supports cell proliferation by supplying iron for DNA synthesis. Exp. Cell. Res., 176 : 87.

115- Lauharanta, J.; Kanerva, L. and Geiger, J. (1988):

Clinical and ultrastructural effects of acitretin in Darrier's disease. Acta Derm. Venereol, 68 : 492.

116- Leaker, B. and Whiteworth, J.A. (1984):

Alopecia associated with captopril treatment. Aust. NZ. J. Med., 214 : 866.

117- Leonard, C. (1996):

Evaluation of hair loss. Curr. Probl. Dermatol., 3 : 101.

118- Lesse, S. (1978):

Tranlycypromine, A study of 1000 patients with severe agitated depression, Am. J. Psychother, 32 : 320.

119- Leung, A.Y. (1977):

The use of Aloe vera in cosmetics. Cosmet. Ind. 120 : 34.

- 120- Levy, L.K. (1950):**
Loss of hair following use of propyl thiouracil. JAMA, 143 : 860.
- 121- Lewis, J.R. (1978):**
Valproic acid. JAMA, 340 : 2190.
- 122- Lovatt, G.E. (1974):**
Allopurinol and drug induced alopecia, Br. J. Dermatol., 91 : 115.
- 123- Ludwig, E. (1977):**
Classification of the type of androgenic alopecia arising in the female sex. Br. J. Dermatol., 97 : 247 - 49.
- 124- Macoul, K.L. (1975):**
Pilocarpine. Arch. Ophthal., 93 : 587.
- 125- Maguire, H.C. (1975):**
Diseases of the hair. In Textbook of Dermatology. Moschella, S.L., Pillsbury. D.H. and Hurley, H.J. (eds), 1st ed. W.B. Saunders Company. London, P : 1191.
- 126- Main - Waring, W. (1969):**
Androgen receptor in the cytoplasm of rat prostate. J. Endocrinol, 45 : 531.
- 127- Marshall, A. (1971):**
Alopecia and leucodopa Br. M. J., 2 : 47.
- 128- Marritt, E. (1984):**
Single hair transplantation for hair refinement : A practical solution. J. Derm. Surg. Oncol., 10 : 962.
- 129- Miller, J.A. (1981):**
Low sex hormone binding globulin in young women with diffuse hair loss. Br. J. Dermatol., 106 : 331.
- 130- Miller, J.A.; Darby, C.R. and Karkavitas, K. (1982):**
Low sex hormone binding globulin level in young women with diffuse hair loss. Br. J. Dermatol., 106 : 331.

131- Mitchell, A.J. (1984):

Alopecia : Pathogenesis and treatment.
J. Am. Acad. Dermatol., 11 : 763.

132- Modly, C.E. (1989):

Evaluation of alopecia. Cutis, 43 : 148.

133- Muting, D. (1969):

Wachstumsstörungen der Kopfhaare als folge von Hepatopathien.
Arch. Klin. Exp. Derm., 235 : 386.

134- Nestel, R. (1973):

Fixed Oils. Lancet, 2 : 131.

135- Oba, K. (1988):

Cosmetic products influencing hair growth.
Cosm. Toil., 103 : 69.

136- Olsen, E.A. and Weiner, M.S. (1987):

Topical minoxidil in male pattern baldness, the effect of
discontinuation of treatment. J. Am. Acad. Dermatol., 17 - 97.

137- Olsen, E. (1989):

Androgenetic alopecia. Dermatol. Clin., 15 : 40.

138- Olsen, E.A. (1989):

Interferon alpha in the treatment of cutaneous T. cell lymphoma
J. Am. Acad. Dermatol., 20 : 395.

139- Olsen, E.A. (1990):

Five year follow-up of men with androgenetic alopecia treated with
topical minoxidil. J. Am. Acad. Dermatol., 22 : 643.

140- Olsen, E.A. (1992):

Terfenadine associated hair loss (replay to letter to editor) N. Med.
Soc. J., 53 : 390.

141- Olsen, E.A. (1994):

Clinical tools for assessing hair loss. In : Disorders of Hair
Growth, Olsen, E.A. McGraw - Hill. Inc. New York, P : 59 - 69.

142- Orfanos, C.E. (1990):

Androgenetic alopecia : clinical aspects and treatment. In : Hair And Hair diseases. Orfanos, C.E. and Happle, R. New York, Springer - Veuag, P. 485 - 527.

143- Papadopoulos, S. and Harden, R.M. (1966):

Hair loss in patients treated with carbimazole, Br. Med. J., 2 : 1502.

144- Pariser, D.M.; Caserio, R.J. and Eaglstein, W.H. (1986):

Techniques for diagnosing skin and hair disease. Dermatol. Clin., 5 : 493.

145- Paus, R. (1991):

Hair growth inhibition by heparin in mice : A model system for studying the modulation of epithelial cell growth by glycosaminoglycans. Br. J. Dermatol., 124 : 415.

146- Peck, G.L. (1979):

Prolonged remissions of cystic and conglobate acne with 13 c is retinoic acid. N Engl. J. Med., 300 (7) : 1028.

147- Penneys, N.S. (1991):

Immunohistochemical distribution of aromatase and 3 B - hydroxysteroid dehydrogenase in human hair follicle and sebaceous gland. J. Cutan. Pathol., 19 : 309.

148- Pinski, J.B. (1991):

Hair transplantation and bald scalp reduction. Dermatol. Clinics, 9 : 151.

149- Poulsen, J. (1981) :

Hair loss, depigmentation of hair, ichthyosis and blepharoconjunctivitis produced by dixyrazine ACTA derm. Venereol., 61 : 85.

150- Radwanski, E.; Perentesis, G.; Symchowicz, S. and Zampaglione, N. (1989):

Single and multiple dose pharmacokinetic evaluation of flutamide in normal geriatric volunteers. J. Clin. Pharmacol., 29 : 554.

- 151- Randall, V. and Ebling, F. (1991):**
Seasonal changes in human hair growth, Br. J. Dermatol., 124 : 146.
- 152- Randall, V.; Hamada, K. and Messenger, A. (1992):**
Mechanism of androgen action in cultured papilla cells. Invest. Dermatol.; 98 : 86.
- 153- Reynolds, J. (1982):**
Fixed oils. In : Martindale, The Extrapharmacopia, Reynolds, J. Pharmaceutical Press, London, Twnty eight edit., P : 694.
- 154- Riella, M.C. (1975):**
Essential fatty acid deficiency in human adults during total parenteral alimentation. J. Pediatr, 81 : 894.
- 155- Rietschel, R.L. and Duncan, S.H. (1987):**
Safety and efficacy of topical minoxidil in the management of androgenetic alopecia. J. Am. Acad. Dermatol., 16 : 677.
- 156- Rosenbaum, E.E. (1979):**
Chrysotherapy. Arch. Intern. Med., 139 : 1316.
- 157- Rosenfield, R. (1985):**
Endocrine influences on hair growth. J. Ped. 107 : 415.
- 158- Rosenfield, R. and Furlanetto, R. (1985):**
Physiologic induction of puberty. J. Ped., 107 : 415.
- 159- Rudzki, E. (1970):**
Mercuric chloride, Br. J. Dermatol., 83 : 543.
- 160- Rushton, D.H. (1990):**
Biochemical and trichological characterization of diffuse alopecia in women. Br. J. Dermatol., 123 : 187.
- 161- Rushton, D.H. (1993):**
Management of hair loss in women. Dermatol. Clin., 20 - 47 : 52.
- 162- Rushton, D.H. (1993):**
Treating reduced hair volume in women.
Cosm. Toil., 108 : 59.

- 163- Rushton, D.H.; Kingsley, I. And Futterweit, W. (1991):**
Quantitative assessment of spironolactone treatment in women with diffuse androgen - dependant alopecia. J. Soc. Cosmet. Chem., 42 : 317.
- 164- Rushton, D.H.; Ramsay, I.D. and James, K.C. (1990):**
Biochemical and trichological characterization of diffuse alopecia in women. Br. J. Dermatol., 123 : 187.
- 165- Rushton, D.H. and Ramsay, I.D. (1992) :**
The importance of adequate serum ferritin levels during oral cyproterone acetate and ethinyl estrodiol treatment of diffuse androgen - dependant alopecia in women. Clin. Endocrinol., 36 : 421.
- 166- Sadjadpour, K. (1974):**
Methysergide alopecia. JAMA, 229 : 539.
- 167- Saito, R.; Hori, Y. and Kuribayashi, S. (1976):**
Alopecia in hypothyroidism. In : Biology And Disease Of The Hair, Kobri, T and Montagna, N. (eds) 1st ed. Univesity Park Press, London, P : 315.
- 168- Salmon, D. (1982):**
Chemical hyperthyroidism : serum triiodothyroinine levels in clinically euthyroid individuals treated with leveothyroxine. Arch. Of Int. Med., 142 : 571.
- 169- Sawaya, M.E. (1990):**
Advances in clinical reserch, Dermatol. Clinic., 15 : 38.
- 170- Sawaya, M.E. (1992) :**
Clinical up dates in hair, Dermatol. Clinic, 15 : 37.
- 171- Sawaya, M.E. (1997):**
Clinical updates in hair; Dermatol. Clinic, 15 : 39.
- 172- Schatter, A. and Shanon, Y. (1989):**
Crohn's ileocolitis presenting as chronic diffuse hair loss J.R. Soc. Med., 82 : 303.

- 173- Schaumburg, G. (1990):**
Histology of the skin. In Histopathology of the skin. Lever, W. and Schawnbarg, G. (Eds). Seventh edit., Vol. 1, J.B. Lippincott Co., Philadelphia P : 9.
- 174- Schell, H. (1991) :**
Cell cycle kinetics of human anagen scalp hair bulbs in thyroid disorders. Dermatologica, 182 : 23.
- 175- Sciarra, F.; Toscano, V.; Concolino, G. and DisSilverio, F. (1990):**
Antiandrogens : Clinical applications. J. Steriod Biochem. Mol. Biol., 37 : 349.
- 176- Scott, M.I. and Scott, A.M. (1992):**
Effects of anabolic androgenic steroids on the pilosebaceous unit. Cutis, 50 : 113.
- 177- Scott, O.P. Calahan, M.G. and Faulkner, R.M. (1976):**
Wigs and hair : In : Cosmetology. Scatt, O.p., Callahan, M.G. and Fanlkner R.M. (eds). 1st edit., Englewood Cliffs, New Jersey, P : 141.
- 178- Simpson, N.B. (1991):**
Hair patterns - Hirsuties and baldness In : Disease of the hair and scalp, Rook, A Dawber, R. Blackwell Scientific publication, London, P : 96.
- 179- Skolnik, P.; Eaglstein, W. and Libott, V. (1977):**
Human essential fatty acid deficiency, Arch. Dermatol., 113 : 939.
- 180- Smith (1993):**
Diagnostic and predictive value of horizontal sections of scalp biopsy specimens in male pattern androgenetic alopecia. J. Cutan. Path., 17 : 325.
- 181- Smith, A.J.; Hoornite, S.J. and Donker, A. (1983):**
Zinc deficiency during captopril treatment. Nephron, 34 : 196.
- 182- Soloman, A.R. (1994):**
The transversely sectioned scalp biopsy specimen. Adv. Dermatol., 9 : 127.

- 183- Steiger, U.; Cotting, J. and Reichen, J. (1990):**
Aloendazole treatment of echinococcosis in humans. Effects on microsomal metabolism and drug tolerance, Clin. Pharmacol. Ther., 47 : 347.
- 184- Spencer, L.V. (1987):**
Hair loss in systemic disease. Dermatol. Clin., 5 (3) : 568.
- 185- Sperling, L. (1991):**
Hair anatomy for clinician J. Am. Acad. Dermatol., 25 : 5.
- 186- Sperling, L.C. (1993):**
Androgen biology as a basis for the diagnosis and treatment of androgenic disorders in women. J. Am. Acad. Dermatol., 28 : 669.
- 187- Sperling, L.C. (1995):**
The histopathology of non - scarring alopecia. J. Cutan. Pathol., 22: 97.
- 188- Sperling, L.C. (1996):**
Evaluation of hair loss, Curr. Probl. Dermatol., 3 : 115.
- 189- Sperling, L. C. (1996):**
Evaluation of individual forms of alopecia.
Curr. Probl. Dermatol., 3 : 108.
- 190- Sperling, L.C. and Lupton, G.P. (1995):**
The histopathology of non - scarring alopecia, J. cutan. Pathol., 22 : 97.
- 191- Steck, W.D. (1979):**
The clinical evaluation of pathologic hair loss. Cutis., 24 : 293.
- 192- Steinert, P.M. (1988) :**
The molecular and cellular biology of intermediate filaments.
Annu. Rev. Biochem., 57 : 593.
- 193- Stengel, F. (1978):**
Diagnosing the common alopecias.
Am. Fam. Physician, 18 : 17.

- 194- Stimson, W.H. (1961):**
Vitamine A intoxication in adults. Report of a case with a summary of literature. N. Engl. J. Med., 265 : 369.
- 195- Sulzberger, M.B; Witten, V.H. and Kopf, A.W. (1960):**
Diffuse alopecia in women. Arch. Dermatol., 81 : 550.
- 196- Sunn, T. T. (1992):**
Interaction of trichohyalin with intermediate filaments : Three immunoligically defined stages of trichohyalin maturation. J. Invest. Dermatol., 98 : 24.
- 197- Tepper, L.B. (1976):**
Absorption and fate of castor oil. Pharm. Ther., 1 : 131.
- 198- Testuo - Makino (1991):**
Determination of zinc in body fluids : Chemica Clinic. Acta, 197 : 209.
- 199- Thompson, J.S. (1989):**
Alopecia after ileal pouch -anal anastmosis. Dis. Colon. Rectum, 32 : 557.
- 200- Tullio, C. (1985):**
Cimetidine - induced alopecia clin. Pharmacol., 4 : 145.
- 201- Unger, W. (1992):**
Punch hair transplants. Clinics in Dermatol., 10 : 335.
- 202- Van Scott, E.J. (1957):**
Detection of radiation effect on hair roots of the human scalp. J. Invest. Dermatol., 29 : 205.
- 203- Venning, V.A. (1988):**
Patterned androgenic alopecia. J. Am. Acad. Dermatol., 18 : 1073.
- 204- Virginia, C. (1987):**
Minoxidil. Dermatol. Clin., 5 (3) : 267.
- 205- Wagner, K.D. and Teicher, M.H. (1991):**
Lithium and hair loss in childhood. Psychosomatics, 32 : 355.

- 206- *Warnock, J.K. (1991):*
Psychotropic medication and drug related alopecia.
Psychosomatics, 32 : 149.
- 207- *Wastell, H.; Bartett, K. and Dale, G. (1988):*
Biotinidase deficiency. A survey of 10 cases. Arch. Dis. Child., 63 : 1244.
- 208- *Watanakunakorn, G.; Hodges, R.E. and Evans, T.C. (1965):*
Myxedema. Arch. Int. Med., 116 : 183.
- 209- *Weinblatt, M.E. (1985):*
Toxicity of low dose methotrexate in rheumatoid arthrities. J. Rheumatol., 12 : 35.
- 210- *Weisman, K. (1977):*
Possible markers of zinc deficiency.
Acta. Dermato-venereologica, 57 : 88.
- 211- *Weisman, K. (1978):*
Acquired zinc deficiency dermatoses in man. Arch. Dermatol., 114 : 1509.
- 212- *Wheeler, G.E. and Barrows, G.H. (1981):*
Alopecia universalis, A manifestation of occult amyloidosis and multiple myeloma. Arch Dermatol., 117 : 815.
- 213- *Whiting, D.A. (1993):*
The diagnostic and predictive value of horizontal sections of scalp biopsies in male pattern androgenetic alopecia. J. Am. Acad. Dermatol., 28 : 755.
- 214- *Whiting, D.A. (1994):*
Disorders of the hair shaft. In : Clinical Dermatology, Demis, D.J. Vol. I, Philadelphia, J.B., Lippincott, P. : 29.
- 215- *Wilburn, M. (1951):*
Hair loss and pigmentation due to thiouracil derviatives. JAMA, 147 : 379.

- 216- *Williams, M.; Packman, S. and Cowan, M. (1983):*
Alopecia and periorificial dermatitis in biotin - responsive multiple carboxylase deficiency. *J. Am. Acad. Dermatol.*, 9 : 97.
- 217- *Windhorst, D.B. and Nigra, T. (1982):*
General clinical toxicology of oral retinoids *J. Am. Acad. Dermatol.*, 6 : 675.
- 218- *Winkelman, R.K.; perry, H.O.; Achor, R.W. and Kirby, T.J. (1963):*
Cutaneous syndromes produced as side effects of triparanol therapy. *Arch. Dermatol.*, 87 : 140.
- 219- *Woodroffe, R.C. (1973):*
Uses of triclosan. *J. Appl. Bact.*, 36 : 387.
- 220- *Wouwe, J. (1989):*
Clinical and laboratory diagnosis of acrodermatitis enteropathica. *Eur. J. Pediat.*, 149 : 2.
- 221- *Younis, I. (1990):*
Incidence of skin diseases over a 5 year period in Cairo, *Med. J. Cairo Univ.*, 57 : 215.
- 222- *Zviak, C. (1986):*
Hair structure, function, and physicochemical propertis. In : *The science of Hair Care*. Zviak, C. Marcel Dekker (eds), New York, P: 44.