


REFERENCES



REFERENCES

- 1- *Akil, H.; Mayer, D. and Liebeskind, J.C. (1976):*
Antagonism of stimulation-produced analgesia by naloxone, a narcotic antagonist. *Science*, 191: 961.
- 2- *Altschuler, R.A.; Mosinger, J.I. and Parakkal, M.H. (1982):*
Immuno cytochemical localization of enkephalin like immuno reactivity in the retina of guinea pig. *Proc. Natl. Acad. Sci. U.S.A.*, 79:2398.
- 3- *Atweh, S.F. and Kuhar, M.J. (1983):*
Distribution and physiological significance of opioid peptide in the brain. *Br. Med. Bull.*, 39(1): 47.
- 4- *Ayres, J.; Rees, J. and Lee, T. (1982):*
I.V. naloxone in acute respiratory failure. *Br. Med. J.*, 284: 929.
- 5- *Babson, A.L. (1991):*
The immulite automated immunoassay system. *J. Clin. Immunoassay*, 14: 83.
- 6- *Barker, J.N.; Karabin, G.D. and Stoof, T.J. (1991):*
Detection of interferon-gamma mRNA in psoriatic epidermis by polymerase chain reaction. *J. Dermatol. Sci.*, 2: 106.
- 7- *Barna, M.; Bos, J.D. and Kapsen Berg, M.L. (1997):*
Effect of calcitriol on production of T-cell derived cytokines in psoriasis. *Br. J. Dermatol.*, 136: 536.
- 8- *Bata- Csorgo, Z.; Hammerberg, C. and Voorhees, J.J. (1993):*
The localization of the primary hyperproliferative population in psoriasis. *J. Exp. Med.*, 178: 1271.
- 9- *Black, A.K. and Gratton, C.E.H. (1992):*
The mediators involved in acute and chronic urticaria and their therapeutic implications. In recent advances in Dermatology. Edited by Champion, R.H. and Rye, R.J. Churchill Livingstone- Edinburgh, London, P: 119.

-
- 10- **Bernstein, J.E. (1983):**
Neuropeptides and the skin. In Gold Smith, L.A. (ed).
Biochemistry and physiology of the skin. Oxford University
Press New York, P: 816.
- 11- **Bonifazi, R.; Carofalo, E.L. and Pisani, V. (1985):**
Role of some infectious agents in AD Acta. Dermatol.,
Venereol (Stockh),114: 98.
- 12- **Bos, J.D. (1988):**
The pathomechanisms of psoriasis, the skin immune system,
and cyclosporin. Br. J. Dermatol.,118: 141.
- 13- **Braathen, L.R. (1989):**
Langerhans cells and atopic dermatitis. Acta. Dermatol.
Venereol, (Stockh), 144: 55.
- 14- **Bruijnzeel-Koomen, C.A.; Mudde, G.C. and Bruijnzeel, P.L. (1992):**
New aspects in the pathogenesis of AD. Acta. Derm.
Venereol, 144: 58.
- 15- **Bruijnzeel-Koomen, C.A.; Mudde, G.C. and Bruijnzeel, P.L. (1989):**
The presence of IgE molecules on epidermal langerhans cells
in AD and their significance for its pathogenesis. Allergy
Immunol. (Paris),21: 219.
- 16- **Bruni, J.F.; Uanuugt, D. and Meites, J. (1977):**
Effect of naloxone, morphine and methionine enkephalins on
serum prolactine, L.H., FSH, TSH, and GH. Life Sci.,21;
461.
- 17- **Byrom, N.A. and Timlin, D.M. (1979):**
Immune status in AD: a survey. Br. J. Dermatol., 100: 491.
- 18- **Clement-Jones, V. and Besser, G.M. (1983):**
Clinical prespective of opioid peptides. Br. Med. Bull,39(1):
95.
- 19- **Chandra, R.K. and Baker, M.C. (1983):**
Numerical and functional deficiency of suppressor T-cells
before development of atopic enzyme. Lancet,II; 1393.
-

- 20- **Champion, R.H. and Parish, W.F. (1992):**
Atopic dermatitis. In: Textbook of Dermatology. Edited by
Champion, R.H. Burton, J.L. and Ebling, F.J.G. Blackwekk
Scientific publications. London, 1 Boston, Edinburgh,
Melbourne, Paris, Berlin, Vienna. Vol. 1, P.: 589.
- 21- **Christopher, E.; Henseler, T. (1989):**
Patient subgroups and the inflammatory pattern in psoriasis.
Acta. Derm. Venereol. (Stockh).,69: 88.
- 22- **Coates, P.J.; Doniach, I.; Hale, A.C. and Rees, L.H. (1986):**
The distribution of α MSH, β -MSH and B-END in the adult
human pitutary gland. J. Endocrinol.,111; 335.
- 23- **Cooper, K.D. (1994):**
Atopic dermatitis, recent trends in pathogenesis and therapy.
J. Invest. Dermatol.,102: 128.
- 24- **Cooper, K.D.; Wuepper, K.D. and Hanifin, J. (1983):**
Immunoregulation in AD Functional analysis of T-B cells
interactions and the enumeration of C receptors bearing T-
cells. J. Invest. Dermatol., 80: 139.
- 25- **Dekloet, E.R.; Palkouits, M. and Mezy, M. (1981):**
Opiocortein peptides, localization sources. Pharmacol. Ther.,
12; 321.
- 26- **Delitala, G.; Grossman, A. and Besser, G.M. (1983):**
The different effects of opiate peptide on anterior pitutary
hormone. Neuro-endocrinology,37; 275.
- 27- **Delitala, G.; Tomasi, P.; Palermo, H. and Besser, G.M. (1989):**
Opioid stimulates growth hormone release independently of
growth hormone releasing hormone. J. Clin. Endocrinol
Metaboli, 69; 356.
- 28- **Delperte, G.; DeCarli, M. and Romagnan, S. (1993):**
Human IL-10 is produced by both type 1 helper (TH1) and
type 2 helper (TH2) T-cells clones and inhibit their antigen
specific proliferation and cytokine production. J. Immunol.,
150: 253.

-
- 29- **Dewing, S.B. (1971):**
Remission of psoriasis associated with cutaneous nerve section. *Arch. Dermatol.*, 104: 220.
- 30- **Elder, J.T.; Nair, R.P. and Voorhees, J.J. (1994):**
Epidemiology and the genetics of psoriasis. *J. Invest. Dermatol.*, 102: 245.
- 31- **Farber, E.M. and Nall, M.L. (1982):**
Epidemiology in psoriasis-research. *Hawaii. Med. J.*, 41: 430.
- 32- **Farber, E.M.; Nickoloff, N.J. and Franki, J.E. (1986):**
Stress, symmetry and psoriasis possible role of neuropeptides. *J. AM. Acad Dermatol.*, 14: 305.
- 33- **Fisher, A.; Comly, M. and Do, R. (1984):**
Two pools of B. end like immunoreactivity in blood, plasma, erythrocytes, *life Sci.*, 34: 1839.
- 34- **Frank, J. and Criffin, T. (1989):**
Stress and immunity. *Veta. Immunol. Immunopathol.*, 20: 263.
- 35- **Gabrilovac, J.; Martin-Kleiner, I. and Osmak, M. (1990):**
Modulation of mouse natural kille cell activity by leukenkephalin in vivo and vitro period: *Biol.*, 92, 79.
- 36- **Gasior-Chrzan, B. and Falk, E.S. (1990):**
Lysozyme and IgA values in patients with AD. *Acta. Dermatol. Venereol (Stcokh)*, 70: 448.
- 37- **Georgala, S.; Schulpis, K.H. and Stratigas, J. (1994):**
Raised B-END serum level in children with AD and pruritus. *J. Dermatol. Sci.*, 8: 125.
- 38- **Gibbins, I.L.; Wattedant, D. and Conventry, B. (1987):**
Two immunohistochemical identified populations of calcitonin gene-related peptides. *Brain, Res.*, 4: 143.
- 39- **Gisolomoni, G. and Giannetti, A. (1989):**
Neuropeptidie. *J. Ital Dermatol, venereol*, 67: 185.
-

- 40- *Glinski, W.; Brodeck, A.; Glinskaferenz, M. and Kowalski, D. (1994):*
Increased concentration, of B-endorphin in sera of patients with psoriasis and other inflammatory dermatosis. Br. J. of Dermatology, 131: 200.
- 41- *Glinski, W.; Brodeck, A.; Glinska-Ferenz, M. and Kowlski, D. (1995):*
Increased concentration of beta endorphin in the sera of patients with severe atopic dermatitis. Acta. Derm. Venerol. (Stockh)., 75: 9.
- 42- *Goldstein, A.; Fischli, W. and Lowney, L.I. (1981):*
Procine pituitary dynorphin: complete amino acid sequence of biologically active hepatopeptide. Proc. Natl. Acad. Sci. USA., 78: 7219.
- 43- *Grabbe, J.; Zuberbier, t. and Wagnepfeil, S. (1993):*
Skin prick test to common allergens in adult with AD and rhinitis. Dermatology, 186: 113.
- 44- *Griffiths, T.W.; Friggiths, C.E.M. and Voorhees, J.J. (1995):*
Immunopathogenesis and immunotherapy of psoriasis. Dermatologic clinics.,13: 739.
- 45- *Grossman, A. and Rees, L.H. (1983):*
Neuroendocrine of opioid peptides. Br. Med. Ball, 39: 88.
- 46- *Gudelsky, G.A. and Porter, J.C. (1979):*
Morphine and opoid peptides induced inhibition of the release of dopamine from tuberoinfundibular neurones. Life Sci., 25: 1697.
- 47- *Guillemin, R. (1977):*
Endorphins, brain peptides that act like opiates. N. Engl. J. Med., 296: 266.
- 48- *Hamid, Q.; Bonguniewicz, M. and Leung, D.Y. (1994):*
Differential in situ cytokine gene expression in atopic dermatitis. J. Clin. Invest.,94 (2): 870.

- 49- **Hanifin, J.M. and Rajka, G. (1980):**
Diagnostic features of atopic dermatitis. *Acta Dermatol. Venerol* (Stockh), 60: 44.
- 50- **Hanifin, J.M. (1990):**
Immunologic aspects of AD. *Dermatol. Clin.*, 8: 747.
- 51- **Hanifin, J.M. (1993):**
Atopic dermatitis. In: *Text book of Dermatology*. Edited by Moshella, S.L. and Hurbey, H.T. Saunders Company, Philadelphia, London, Toronto. Vol., 3. P: 441.
- 52- **Hardas, B.D.; Zhao, X.; Zhang, J.; Shol, S. and Elder, J.T. (1996):**
Genetics of psoriasis, coordinate over expression of clustered genes in psoriasis. *J. Invest. Dermatol.*, 106(4): 753.
- 53- **Harrison, P.V. (1990):**
Nalmefen and pruritus (letter). *J. Am. Acad. Dermatol.*, 23: 530.
- 54- **Henseler, T. and Christophers-Enno, (1995):**
Disease concomitance in psoriasis. *J. Am. Acad. Dermatol.*, 32: 982.
- 55- **Hirooka, Y.R.; Chardson, S. and Prasad, J. (1978):**
Modulation of the hypothalamic hormone release. *Endocrinology*. (Suppl): 102: 25A.
- 56- **Holtt, V.R.; Pizewtocki and Aherz, A. (1978):**
Radiommuo-assay of beta-endorphin basal and stimulated levels in extracted rat plasma. *Archives of pharmacology*, 303 (2): 171.
- 57- **Holzer, P. (1988):**
Local effector functions of capsaicin: sensitive sensory nerve endings involvement of Tachykinines, CGRP and other NP. *Neuroscience*, 24: 739.
- 58- **Horrocks, C.; Holder, J.E. and Camp, R.D. (1997):**
Antigen independent expansion of T-cells from psoriatic skin lesions. *Br. J. of Dermatol.*, 137: 331.

- 59- **Howlett, J.A. and Rees, L.H. (1988):**
Endogenous opioid peptides. In Oxford text book of medicine (ed). Weatherall, D.J. Ledingham, J.G. and Warrell, D.A. Second edition EIBS. P: 10.
- 60- **Hughes, J.; Smith, T.W. and Kosterlitz, H.W. (1975):**
Identification of two related pentapeptides from the brain with potent opiate agonist activity. Nature, 258; 557.
- 61- **Hunt, S.J. and Jegasothy, B.V. (1992):**
The function of cell in inflammatory disease chapter 10. In Dermatology Ed. By Moschella, S.L. and Aurley, H.T. 3rd edition. Pub. By W.B. Saunders Company. Philadelphia, London, Toronto, Tokyo, P.; 250.
- 62- **Infortuna, M.; Bardero, G. and Santisi, G. (1989):**
The behaviour of lymphocyte population and the immunoglobulins in children with AD in whom asthma develops Minerva pediatri., 41: 253.
- 63- **Imayama, S.; Shimozone, Y.; Ohta, S. and Yoneyama, K. (1994):**
Reduced secretion of IgA and skin surface of patients with AD. J. Allergy Clin. Immunol., 94: 195.
- 64- **Imura, H.; Kato, Y. and Nakai, Y. (1985):**
Endogenous opioid and related peptides from molecular biology to clinical medicine. J. Endocrinol, 107: 147.
- 65- **Jankovic, B.D. and Maric, D. (1990):**
In vivo modulation of immune system by enkephalins. Int. J. Neuro. Sci., 51: 167.
- 66- **Jegasothy, B.V.; Ackerman, C.D. and Todo (1992):**
A new therapeutic agent for severe psoriasis. Arch. Dermatol., 128: 181.
- 67- **Jones, H.E.; Levis, C.W. and MC-Gerity, J. (1973):**
Allergic contact sensitivity in AD. Arch. Dermatol., 107: 217.
- 68- **Judd, A.M. and Hedge, G.A. (1983):**
Direct pituitary stimulation of thyrotropin. Secretion by opioid peptides. Endocrinol, 113: 706.

-
- 69- *Julliard, J.H.; Shibasaki, T. and Ling, N. (1980):*
Measurement of B-endorphin in human plasma. *Science*,
208: 183.
- 70- *Kadunce, D.P. and Krueger, G.G. (1995):*
Pathogenesis of psoriasis. *Dermatologic Clinics*, 13: 723.
- 71- *Kahn, B.D.; Chang, J.Y. and Sehgal, S.N. (1992):*
Selective T-cell immuno suppressant agent. *Clin. Haematol.*,
11: 743.
- 72- *Kapsenberg, M.L.; Wierenga, E.A. and Vander Heijen, F.L. (1992):*
Atopic dermatitis and CD4+ allergen specific Th2
lymphocytes. *Eur. J. Dermatol.*, 2: 601.
- 73- *Kawana, S. and Nishiyama, s. (1995):*
Possible role of leukotriens B4 and CD4 in a flare of
synovitis. *Dermatology*, 190 (1): 35.
- 74- *Kemmet, D. and Tidman, M.T. (1991):*
The influence of the menstrual cycle and pregnancy on
atopic dermatitis. *Br. J. Dermatol.*, 125: 59.
- 75- *Kenneth-Blaylock, V. (1985):*
Atopic dermatitis in: *Dermatologic immunology and allergy*.
Edited by Stone. J. C.V. Mosby company, Toronot,
Principetion, P: 515.
- 76- *Kimura, J.Y.; Ohta, N. and Ishii, A. (1990):*
Lymphocyte response to house dust mite antigens in atopic
non atopic individuals. *Immunology*, 70: 385.
- 77- *Krueger, G.G. (1989):*
A perspective of psoriasis as an aberration in skin modified
to expression by inflammatory repair system. In Norris DA
(ed): *Immune mechanisms in cutaneous disease*.
(immunology series). New Yokr. Marcel Dekker, 1989, P.
425.
- 78- *Krulich, L.; Ciachetti, A. and Marchlewska, A. (1977):*
Regulation of TSH secretion in rates. *Endocrinology*, 100:
496.
-

-
- 79- **Kuntzman, R.; Jacobson, M. and Scheneidman K. (1964):**
Similarities between oxidative drug metabolising enzymes and steroid hydroxylases in liver microsomes. *J. Pharmacol. Exp. Ther.*, 146: 280.
- 80- **Labella, F.S. (1975):**
Opiate specific displacement of steroid hormones from microsomes. *Life sci.*, 16: 1783
- 81- **Lavker, R.M. and Sun, T. (1983):**
Epidermal stem cells. *J. Invest. Dermatol.*, 81: 1215.
- 82- **Leehey, D.J.; Gollapudi, P. and Reid, R.W. (1991):**
Naloxone increase water, electrolyte excretion after water loading in patients with cirrhosis and ascitis. *J. Lab. Clin. Med.*, 118: 484.
- 83- **Leung, D.Y.M.; Travers, J.B. and Norris, D.A. (1995):**
The role of superantigens in skin diseases.
J. Invest. Dermatol., 105: 375.
- 84- **Lightman, S.L. and Jacobs, H.S. (1979):**
Naloxone: Non steroidal treatment for post-menopausal flushing. *Lancet*, 2: 1071.
- 85- **Luger, T.A. and Schwarz, T. (1991):**
Therapeutic use of cytokines in dermatology.
J. Am. Acad. Dermatol., 24: 915.
- 86- **Mains, R.E.; Eipper, B.A. and Ling, N. (1977):**
Common precursor to corticotropins and endorphins.
Proc. Natl. Acad. Sci. U.S.A., 74: 3014.
- 87- **Marone, G.; Lembo, G. and Byala, F. (1986):**
Human basophil release ability changes in patients with AD.
J. Invest. Dermatol., 87: 19.
- 88- **Mayer, D.J. and Liebeskind, J.C. (1974):**
Pain reduction by focal electrical stimulation of the brain. An anatomical and behavioral analysis. *Brain Res.*, 68: 73.
-

- 89- Meites, J.; Bruni, J.F. and Smith, A.F. (1979):
Relation of endogenous opioid peptides and morphine to neuroendocrine functions. Life Sci., 24: 1325.
- 90- Melmed, S.; Morley, J.E. and Kurtzman, G. (1979):
Opiates and dopamine enhance prolactin release from cultured pituitary tumor cells. Clin. Res., 27: 257A.
- 91- Morley, J.E. (1981):
The endocrinology of opiate and opioid peptides. Metabolism, 30: 195.
- 92- Mousa, A.M.; Kamel, M.A.; Ragab, M.S.; Zaher, H.A.; Abou El-Ela, M.M. and Shaaban, F. (1992):
Serum IgE and IgG levels in AD. Their relation to severity, family history and presence of atopic disease. Egypt, J. Dermatol., venereol., 12: 15.
- 93- Mozzanica, N.; Cattaneo, A. and Finizi, A. (1994):
Plasma neuropeptide level in psoriasis. Acta. Dermatovenereologica, 186: 67.
- 94- Nakao, K.; Nakai, Y. and Oki, S. (1978):
Presence of immuno reactive B-endorphine in normal human plasma. J. Clin. Invest, 62: 1399.
- 95- Naukkarinen, A.; Nickoloff, B.J. and Farber, E.M. (1989):
Quantification of cutaneous sensory nerves and their substance P contents in psoriasis. Invest. Dermatol., 92: 126.
- 96- Nickoloff, B.J. (1991):
The cytokine network in psoriasis. Arch. Dermatol., 127: 1835.
- 97- Nickoloff, B.J.; Fivenson, D.P. and Modlin, R. (1991):
Preferential but not unique cutaneous endothelial cell expression of ELAM-1. Clin. Res., 39: 153.
- 98- Olsen, G.A.; Olson, R.D. and Kastin, A.J. (1986):
Endogenous opiates. Peptides, 7: 907-33.

-
- 99- Ott, N.L.; Gleich, G.J. and Peterson, E.A. (1994):**
Assessment of eosinophil and neutrophil participation in AD.
J. Allergy Clin. Immunol., 94: 120.
- 100- Pasternak, G.W. (1981):**
Opiate enkephalin and endorphin analgesia. Neurology, 31:
1311.
- 101- Paterson, S.J.; Robson, J.E. and Kosterlitz, H.W. (1983):**
Classification of opiate receptors. Br. Med. Bull, 39: 31.
- 102- Perez, S.E.A.; Silveria, J.E.N. and Silvanetto, C.R. (1990):**
Participation of opiate pathways on the lateral hypothalamic
area in the control of renal electrolytes and water excretion.
Braz. J. Med. Biol. Res., 23: 1343.
- 103- Pico, A.; Caciado, I. and Varela, (1986):**
Interaction of thyrotropin releasing hormone and the
enkephalin analogue DAMM E. on Pitutary hormone
secretions. Clin. Endocrinol, 25; 501.
- 104- Pincelli, G.; Fantini, F. and Massini, P. (1990):**
Neuropeptides in the skin from patients with atopic
dermatitis. Br. J. Dermatol., 122: 745.
- 105- Polak, J.M.; Sullivan, S.N. and Bloom, S.R. (1977):**
Enkephalin like immunoreactivity in human gastrointestinal
tract. Lancet, 1: 972.
- 106- Rahe, R. (1995):**
Life change events. Comprehensive text book of psychiatry
VI volume 2. Edited by Kaplan, H. and Sadock, B. P. 1545.
- 107- Rajka, G. (1990):**
Grading of the severity of atopic. Dermatitis. Acta.
Dermatol. Venerol., (Stockh.) 144: 13.
- 108- Ramsay, B. and Lawrence, G.M. (1991):**
Measurement of involved surface in patients with psoriasis.
Br.J. Dermatol., 124: 565.
-

- 109- Rees, J.; Friedmann, P.S. and Mathews, J.N.S. (1990):**
Contact sensitivity to dinitrochlorobenzene, Impaired in atopic dermatitis. Arch. Dermatol., 126: 1173.
- 110- Reynolds, D.V. (1969):**
Surgery in the rats during electrical analgesia induced by focal brain stimulation. Science, 164: 444.
- 111- Ring, J. (1991):**
The pathophysiology of AD. In hand book of AD Edited by Ruzicka, T.; Ring, J. and Przyhill. B. Springer, P: 330.
- 112- Romagnani, S. (1992):**
Human TH₁ and TH₂ subsets regulation and differentiation 2nd role in protein and immunopathology. Int. Arch. Allergy. Immunol., 98: 279.
- 113- Rubin, P.; Blaschke, T.F. and Guilleminault, C. (1981):**
Effect of naloxone on blood pressure falling during sleep. Circulation, 63: 117.
- 114- Rupprecht, M.; Hornstein, O.P. and Koch, H.U. (1995):**
Corticotropin, cortisol, beta-END, responses to corticotropin-releasing hormone in patients with atopic eczema. Psychoneuroendocrinology, 20: 543.
- 115- Rupprecht, M.; Salzer, B. and Raum, B. (1997):**
Physical stress-induced secretion of adrenal and pituitary hormones in patients with atopic eczema. Exper. and Clini. Endocrin and Diabetes, 105: 39.
- 116- Ruzicka, E.; Samochocki, A. and Litewska, D. (1991):**
Hand book of atopic eczema. Quoted from Ring and Bieber (1992).
- 117- Scharrer, B. (1991):**
Neuro-immunology. The importance and role of comparative approach. Adv. Neuroimmunology, 1: 1.
- 118- Scott, P. and Kaufmann, S.H.E. (1991):**
The role of T-cell subsets and cytokines in the regulation of infection. Immunol. Today, 12: 346.

119- Seckl, J.R. and Lightman, S.L. (1987):

Effect of naloxone on oxytocin and vassopressin release during vagino cervical stimulation of the goat. *J. Endocrinol.*, 115: 317.

120- Shanker, G. and Shama, R.K. (1979):

B-endorphin stimulates Corticosterone synthesis in isolated rat adrenal cells. *Biochem. Res. Commun.*, 86: 1.

121- Shinizu, T.; Kristjansson, S. and Wennergren, G. (1994):

Leukotriene B₄ and C₄ generation by blood leucocytes after ex. vivo stimulation in children with AD. *Pediatr. Allergy Immunol.*, 5(2): 95.

122- Sillevis Smith, J.H.; Bos, J.D. and Krieg, S.R. (1986):

Antigen-presenting cells and T-cell subsets in AD. *Clin. Exp. Dermatol.*, 11: 159.

123- Slominski, A.; Wortsman, J. and Gorbani, A. (1993):

The carcinogenic risk of treatment with photchemotherapy. *J. Labor. And Clinic. Medicine*, 122: 658.

124- Snedecor, G. and Cochran, W.G. (1989):

Statistical methods, 8th ed. Ames, J.A. Lowastate University Press.

125- Straus, E.W. and Raufman, J.R. (1985):

Regulatory systems polypeptide in gastro-enterology (ed) Berk, J.E. Fourth edition W.B. Saunders company Philadelphia New York.

126- Swanbeck, G.; Inerot, A.; Martinsson, T. and Enlund, F. (1995):

Epidemiology of psoriasis. *Br. J. Dermatol.*, 133 (5): 768.

127- Teofoli, P.; Lotti, T. and Guarcello, V. (1993):

Detection of mRNA encoding pro-opio-melanocortin in the A431 cell line. Abstract book of third congress of the European Academy of Dermatology and Venerology, Sept. 26-30, Copenhagen: 352.

128- Theeuwes, M. and Morhenn, V. (1995):

Allelic instability in the miotsis model and the inheretence of psoriasis. *J. Am. Acad. Dermatol.*, 32 (1): 44.

- 129- *Thestrup-Pedersen, K. (1989):*
Immunology of atopic dermatitis and psoriasis. *Acta. Dermatol. Venerol. (Stockh.)*, 69: 77.
- 130- *Thomas, J.A.; Shahid-Salles, K.S. and Donovan, M.P. (1977):*
Effects of narcotics on the reproductive system.
Adv. Sex. Horm. Res., 3: 169.
- 131- *Thomas, P.; Pfutzner, W. and Przybilla, B. (1995):*
Frequency and activity of IgE secreting peripheral.
Blood B-cells in AD. *Science*, 271: 733.
- 132- *Thomas, P.; Semer, H. and Rinez, J. (1989):*
In vitro IgE secretion in AD. *Acta. Dermatol. Venerol. (Stockh.)*, 144: 110.
- 133- *Torellolotti, H.D.; Giuseppe-Hautmann and Emiliano, P. (1995):*
Neuropeptides skin and the skin. *J. Am. Acad. Dermatol.*, 33: 482.
- 134- *Urisu, A.; Kondo, Y.; Wada, E. and Hariba, F. (1994):*
Specific IgE antibodies and serum eosinophil. Cationic protein in AD patients. *Acta. Paediatr. JPN.*, 36(2): 146.
- 135- *Uyemura, K.; Yamamura, M. and Fivenson, D.F. (1993):*
The cytokine network in lesional and lesion free psoriatic skin is characterised by a T-helper type I cell-mediated response. *J. Invest. Dermatol.*, 101: 701.
- 136- *Vale, W.; Rivier, C.; Yang, L. (1978):*
Effects of purified hypothalamic corticotropin-releasing factor on the secretion of adrenocorticotrophin and B-endorphin-like immuno. activities in vitro. *Endocrinology*, 103: 1910.
- 137- *Venkova, K.; Radomirov, R. and Pencheva, N. (1990):*
Effect of leucin enkephalin on mechanical activity of longitudinal muscles of small intestine of cat.
Neuropharmacology, 28: 1183.
- 138- *Wakita, H.; Sakamoto, T. and Tokura, Y. (1994):*
E-selectin and vascular cell adhesion molecule-1 for infiltration of T-lymphocytes in AD.
J. Cut. Pathol., 21(1): 33