

References

REFERENCES

- Adashek J.A.; Lagrew D.C.; Iriye B.K.; Carr M.H.; Porto M and Freeman R.K (1996): The influence of ultrasound examination at term on the rate of cesarean section. *Am. J. Obstet Gynecol.* 174:328.
- American collage of Obstetricians and Gynecologists (1994): Management of diabetes mellitus in pregnancy. Technical Bulletin No.92.
- American collage of Obstetricians and Gynecologists (1999): Dietary therapy for gestational diabetes: How long is long enough? 93:978-982.
- American Diabetes Association (1995): Medical management of pregnancy complicated by diabetes, 2nd ed. Jovanovic-Peterson L (ed) Alexandria, Virginia, ADA.
- American Diabetes Association Inc. (2002): Gestational Diabetes Mellitus. *Diabetes Care* 25: S94 – S96.
- Anoon S.S.; Rizk D.E. and Ezimokhai M. (2003): Obstetric outcome of excessively overgrown fetuses. *J. Perinat Med.* 31 (4): 295-301.
- Archibong E.I.; Sobande A.A. and Assindi A.A. (2003): Antenatal intrauterine fetal death ; a prospective study in a tertiary hospital in South-Western Saudi. Arabia. *J. Obstet Gynecol.* 23 (2): 170-3.
- Bajoria R.; Ward S. and Chatterjee R. (2003): Brain natriuretic peptide and evodothelin-1 in the pathogenesis twins. *Am. J. Obstet Gynecol.* 189 (1): 189-94.
- Banaczek Z. and Penza G. (1995): Evaluation of a seven years period of managing pregnancies complicated by diabetes mellitus. *Ginekol Pol.* 66 (2): 76-9.

- Bansal R.K.; Ecker J.L.; and Lares R.K (1996): Blood glucose monitoring in gestational diabetes. *N Engl Med* 334:598.
- Bar-Hava I.; Scarpelli S.A.; Barnhard Y.; and Divon M.Y. (1994): Amniotic fluid volume reflects recent glycemic status in gestational diabetes mellitus. *Am. J. Obstet Gynecol.* 171: 952.
- Barnhart Y.; Bar-Hava I. and Divon M.Y. (1995): Is polyhydramnios is an ultrasonographically normal fetus an indication for genetic evaluation?. *Am. J. Obstet Gynecol.* 173: 1523-1527.
- Bartha J.L.; Martinez-Del-Fresno P. and Comino-Delgads R. (2003): Early diagnosis of gestational diabetes mellitus and prevention of diabetes-related complication. *Eur. J. Obstet Gynecol. Repord Biol.* 109 (1): 41-4.
- Beand R.W. and Koet J.J. (1982): Is gestational diabetes a clinical entity?. *Diabetologia* 23: 307-12.
- Beard R.W. and Lowy C. (1982): commentary: the British Surgery of diabetic pregnancies. *Br. J. Obstet Gynecol.* 89:783.
- Ben-Chetrit A.; Hokner- Clnikier D. and Rom M. (1990): Hydramnios in the third trimester of pregnancy: Achange in the distribution of accompanying fetal anomalies as a result of early ultrasonographic prenatal diagnosis. *Am. J. Obstet Gynecol.* 162: 1344-1345.
- Ben-Haoush A.; Yogev Y.; and Had M. (2004): Epidemiology of gestational diabetes mellitus and its association with type 2 diabetes. *Diabet Med.* 21 (2): 103-13.
- Berkowitz K.; Reyes C.; Sadaat P. and Kjos S.5 (1996): Comparison of fetal lung maturation in well dated diabetic and non diabetic pregnancies. *Am. J. Obstet. Gynecol.* 174-373.

- Biggio T.R.; Wenstrom K.D.; Dubard M.B. and Cliver S.P. (1999): Hydramnios prediction of adverse perinatal outcome. *Obstet Gynecol.* 94 (Spt 1): 773-7.
- Blackwell S.C.; Hassan S.S.; Berry S.M.; Treadwell M.C.; Zador I. and Wolfe H.M (2003): Abnormal amniotic fluid volume as a screening test prior to targeted ultrasound. *Med. Sci. Monit* 9 (11): 119-22.
- Bo S.; Menato G.; Gallo M.L.; Bardelli C.; Lezo A.; Signorile A.; Gambino R.; Cassader M.; Massobrio M. and Pagano G. (2004): Mild gestational hyperglycemia, the metabolic syndrome and adverse neonatal outcomes. *Acta Obstet Gynecol Scand.* 83(4): 335-40.
- Bottom S.F.; Welch R.A. and Zader I.E. (1986): Limitation of using maximum vertical pocket and other sonographic evaluation of amniotic fluid volume to predict fetal growth: Technical or physiologic? *Am. J. Obstet Gynecol.* 155: 154-6.
- Boyd E.; Metzger B.E.; Donald R. and Coustan M.D. (1997): Summary and recommendations of the Fourth International Workshop-Conference on Gestational Diabetes Mellitus. *Diabetes* 46 (Suppl. 4): 158-161.
- Brace R.A. (1986): Amniotic fluid volume and its relationship to fetal fluid balance: Review of experimental data. *Semin perinatal.* 10: 103-112.
- Brace R.A. (1995): Progress toward understanding the regulation of amniotic fluid volume : water and solute fluxes in and through the fetal membranes. *Placenta.* 16: 1-18.

- Brace R.A. and Resnick R. (1999): Dynamics and disorders of amniotic fluid. In: Creasy R.K.; Resnick R.; editors. Maternal-fetal medicine. 4th ed. Philadelphia: WB Saunders; 1999. P. 640.
- Brace R.A. and Wolf E.J. (1989): Normal amniotic fluid volume changes throughout pregnancy. *Am. J. Obstet Gynecol.* 161: 382-388.
- Brace R.A.; Wlodek M.E. and Cock M.L. (1994): Swallowing of lung liquid and amniotic fluid by the ovine fetus under normoxic and hypoxic conditions. *Am. J. Obstet Gynecol.* 171: 1764-1770.
- Bradley R.J.; Brudenell J.M. and Nicolaides K.H. (1991): Fetal acidosis and hyperlacticaemia diagnosed by cordocentesis in pregnancies complicated by maternal diabetes mellitus: *Diabet. Med.* 8 (5): 464-8.
- Brady K.; Polzin W.H. and Kopelman J.N. (1992): Risk of chromosomal abnormalities in patient with idiopathic polyhydramnios. *Obstet Gynecol.* 79: 234-238.
- Bran Y.W.; Shnnon D.L. and Hunter M.A. (1983): Maternal diabetes and neonatal macrosomia I, postpartum maternal haemoglobin A1C levels and neonatal hypoglycemia. *Pediatrics* (4): 576-81.
- Branhart Y. Bar-Hava I. and Divon M.Y. (1995): Is poyhydramnios in an ultrasonographically normal fetus an indication for genetic evaluation ?. *Am. J. Obstet Gynecol.* 173: 1523-1527.
- Carlson D.E.; Platt L.D. and Medearis N. (1990): Quantifiable poyhydramnios: diagnosis and management. *Obstet Gynecol.* 75: 989-993.
- Chamberlain P.F.; Manning F.A. and Morrison I. (1984): Ultrasonic evaluation of amniotic fluid volume I: The relationship of marginal

- and decreased amniotic fluid volumes to perinatal outcome. *Am. J. Obstet Gynecol.* 150: 245-249.
- Chamberlein P.F.; Manning F.A.; Morrison I.; Harman C.R. and Lavge I.R. (1984): Ultrasound evaluation of amniotic fluid volume. *Am. J. Obstet Gynecol.* 150: 250-254.
- Combs C.A.; Singh N.B. and Khoury J.C. (1993b): Elective induction versus spontaneous labour after sonographic diagnosis of fetal macrosomia. *Obstet Gynecol.* 81:492.
- Conway D. and Langer O. (1996): Elective delivery for macrosomia in the diabetic pregnancy: A clinical Cost-benefit analysis *Am. J. Obstet Gynecol.* 174:331.
- Cooper M.J.; Enderlein M.A.; Tarnoff H. and Roge C.L. (1992): Asymmetric septal hypertrophy in infants of diabetic mothers, fetal echocardiography and the impact of maternal diabetic control. *Am. J. Dis. Child.* 146 (2): 205-9.
- Cordero L. Treuer S.H.; Landon M.B. and Gabbe S.C. (1998): Management of infants of diabetic mothers. *Arch Pediatr Adolesc Med.* 152 (3): 249-54.
- Cousins M.L. (1987): Pregnancy complications among diabetic women: Review 1965-1988. *Obstet Gynecol. Surv.* 42:140.
- Cowetle R.M. and Schwartz R. (1982): The infant of diabetic mother. *Pediatr. Clin of N. Am.* 29 (5): 1213-1231.
- Cruikshank D.P. and Pitkin R.M. (1983): Calcium metabolism in diabetic mothers, fetus and newborn infant *Am. J. Obstet Gynecol.* 145:1010.

- Cunningham F.G.; Macdonald P.C.; Gant N.F.; Leveno K.J.; Gilstrap L.C.; Mankins G.D. and Clarfe S.L. (1997): Diseases and abnormalities of the fetal membranes. In Williams Obstetrics. Textbook of Obstetric 20th edition P. 661.
- Cunningham F.G.; Macdonald P.C.; Gant N.F.; Leveno K.J.; Gilstrap L.C.; Mankins G.D. and Clarfe S.L. (1997): Management of gestational diabetes mellitus. In Williams Obstetrics. Textbook of Obstetric 20th edition P. 1208.
- Dacus J.V.; Meyer N.L.; Muram D.; Stilson R.; Phipps P. and Sibai B.M. (1994): Gestational diabetes: Postpartum glucose tolerance testing. Am. J. Obstet Gynecol. 171: 927.
- Damm P. ; and Molsted-Pedersen L. (1989): Significant decrease in congenital malformation in newborn infants of an unselected population of diabetic women. Am. J. Obstet Gynecol. 161: 1163-7.
- Dashe J.S.; Molntire D.D.; Ramus R.M.; Santo-Rames R. and Twickler D.M. (2002): Hydramnios: anomaly prevalence and sonographic detection. Obstet. Gynecol. 100 (1): 134-9.
- Daskas N.; Argyropoulou M.; Pavlou M. and Andronikous S. (2002): Congenital mesoblastic nephroma associated with poyhydramnios and hypercalcemia. Pediatric. Nephrol. 17 (3): 187-9.
- De Veciana M, Major C.A.; Morgan M.; Asrat T.; Toohey J.S.; Lien J.M. and Evans A.T. (1995): Postprandial versus preprandial blood glucose monitoring in women with gestational diabetes mellitus requiring insulin therapy : N Engl J. Med. 333: 1237.

- Di Cianni R.; Volp L.; Lencioni C. and Del Prato S. (2003): Intermediate metabolism in normal pregnancy and in gestational diabetes. *Diabetes Metab. Res. Rev.* 19 (4): 259-70.
- Dicker D.; Feldberg D. and Samuel N. (1998): Spontaneous abortion in patients with IDDM. The effect of pre-conception diabetic control *Am. J. Obstet. Gynecol.* 158: 1161-4.
- Donald R.; Coustan D.R.; Marshall W. and Carpenter M.D. (1985): Detection and treatment of gestational diabetes. *Clinical Obstet Gynecol.* 28 No. 3 September.
- Donald R.; Coustan D.R.; and Philip F. (1988): Diabetes mellitus. Medical complication during pregnancy 344-58.
- Expert Committee on the Diagnosis and Classification of Diabetes Mellitus (2002): Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 25 (Suppl.1): S5-S20.
- Fadda G.M.; Antona D. D.; Ambrosini G.; Cherchi P.L.; Nardelli G.B.; Capobianco G. and Dessole S. (2001): Placental and fetal pulsatility in gestational diabetes mellitus. *J. Reprod Med.* 46 (4): 365-70.
- Flack N.J.; Sepulveda G. and Bower S. (1995): Acute maternal hydration in third trimester oligohydramnios : effect on amniotic fluid volume, uteroplacental perfusion, and fetal blood flow and urine output. *Am. J. Obstet Gynecol.* 173: 1186-1191.
- Fontain P. (2003): Gestational diabetes. *Rev. Prat.* 53 (17): 1984-9.
- Franz M.J.; Horton E.S.; Bantle J.P.; Beebe C.A.; Brunzell J.D.; Coulston A.M.; Henry R.R.; Hoogwerf B.J. and Stacpoole P.W. (1994):

- Nutrition principles for the management of diabetes and related complications. *Diabetes care* 17: 490-518.
- Garner P.R (1995b): Type I diabetes and pregnancy. Correspondence. *Lancet* 346:966.
- Garner P.R.; M.E.; Alton D. and Dudley K.H. (1990): Pre-eclampsia in diabetic pregnancies. *Am. J. Obstet Gynecol.* 163: 505 – 8.
- Gilbert W.M. and Brace R.A. (1989): The missing link in amniotic fluid volume regulation : intra-membranous absorption. *Obstet Gynecol.* 74: 748-754.
- Gilbert W.M. and Brace R.A. (1993): Amniotic fluid volume and normal flows to and from the amniotic cavity. *Semin Perinatol.* 17): 150 - 157.
- Golan A.; Wolman I.; Sailer Y. and David M.P. (1993): Hydramnios in singleton pregnancy: Sonographic prevalence and etiology. *Gynecol. Obstet Invest.* 35: 91-3.
- Golan A.; Wolman I. and Sagi J. (1994): Persistence of polyhydramnios during pregnancy- its significance and correlation with maternal and fetal complications. *Gynecol. Obstet Invest* 37: 18-20.
- Greenberg L.R.; Moore T.R. and Murphy H. (1995): Gestational diabetes mellitus: Antenatal variables as predictors of postpartum glucose intolerance. *Obstet Gynecol.* 86: 97.
- Hallak M.; Kirshon B.; Smith E.O. and Cotton D.B. (1993): Amniotic fluid index. Gestational age-specific values for normal human pregnancy. *J. Reprod Med.* 38: 853.

- Harding R. (1994): Development of the respiratory system. In : Thorburn G.D.; Harding R. eds. Textbook of fetal physiology. Oxford University Press. 140-167.
- Hedderson M.M.; Ferrara A. and Sacks D.A. (2003): Gestational diabetes mellitus and lesser degrees of pregnancy hyperglycemia: association with increased risk of spontaneous preterm birth. *Obstet Gynecol.* 102 (4): 850-6.
- Hedriana H.L. and Moore T.R. (1994): Accuracy limits of ultrasonographic estimation of human fetal urinary flow rate. *Am. J. Obstet Gynecol.* 171: 989-992.
- Hendricks S.K.; Conway L. and Wang K. (1991): Diagnosis of polyhydramnios in early gestation: indication for prenatal diagnosis. *Prenat Diagn.* 11: 649-654.
- Hernandez V. and Mared Zarate A. (2001): Fetal weight at weight as predisposing risk factor for type 2 diabetes in adulthood. *Gynecol. Obstet Mex.* 69: 390-8.
- Hibbard (1988): Definition, diagnosis and classification in WHO, Expert committee on diabetes mellitus. (1988) Technical report series 646. WHO, Geneva, PP. 8-12 Quoted by Hibbard (1988). P. 287.
- Hill L.M.; Breckle R. and Thomas M.L. (1987): Polyhydramnios ultrasonographically detected prevalence and neonatal outcome. *Obstet Gynecol.* 69: 21-25.
- Hoddick W.K.; Callen K.S. and Filly P.W. (1984): Ultrasonic determination of quantitative amniotic fluid volume in intrauterine growth retardation: Reassessment of 1-cm rule. *Am. J. Obstet Gynecol.* 149: 758-62.

- Hopp H.; Vollert W. and Ebert A. (1995): Diabetic retinopathy and nephropathy complications in pregnancy and labour. *Geburtshilfe Frauenheilkd* 55 (5): 275-9.
- Hoshi J.; Nishida H. and Takahashi N. (1991): Prenatal morbidity of infants of diabetic mothers. *Acta. Pediatr. JPN.* 33 (2): 159-65.
- Ilce Z.; Erdogan E.; Kara C.; Celayir S. and Sarimurat N. (2003): Pyloric atresia review from a single institution. *J. Pediatr. Surg.* 38 (11): 1581-4.
- Jaffe R. (2002): Identification of fetal growth abnormalities in diabetes mellitus. *Semin Perinatal.* 26 (3): 190-5.
- Jang P.R. and Brace R.A. (1992): Amniotic fluid composition changes during urine drainage and tracheoesophageal occlusion in feta sheep. *Am. J. Obstet Gynecol.* 167: 1732-1742.
- Jeong-Ah Kim; Jeong Yeon Cho; Young Ho Lee and Yi-Kyeong Chun (2003): Complications arising in Twin pregnancy: Finding of prenatal ultrasonography. *Korean J. Radiology* 4 (1): 54-60.
- Jensen M.L.; Rix M. and Schroder H.D. (1995): fetal akinesia-hypokinesia deformation sequence (FADS) in two siblings with congenital dystrophy. *Clin. Neuropathy* 14: 105-108.
- Jimenez-Moleon J. J.; Bueno-Cavanillas A.; Luna-del-Castillo Jde D.; Garcia-Martin M.; Lardelli-Clart P. and Galvez-Vargas R. (2002): Impact of different levels of carbohydrate intolerance on neonatal outcomes classically associated with gestational diabetes mellitus. *Eur. J. Obstet Gynecol. Reprod Biol.* 102 (1): 36-41.
- Jodi S.; Lawrence N.; Donald D. and Kenneth J. (2000): Correlation between amniotic fluid glucose concentration and amniotic fluid

- volume in pregnancy complicated by diabetes Am. J. Obstet Gynecol. 182: 901-4.
- Jolly M.C.; Sebire N.J.; Harris J.P.; Regan L. and Robinson S. (2003) : Risk factors for macrosomia and its clinical consequences: a study of 350,311 pregnancies. Eur J. Obst. Gynecol Repord Biol. 111 (1): 9.
- Jones C.W. (2001): Gestational diabetes and its impact on the neonate. Neonatal Netw. 20 (6): 17-23.
- Kalaidzhieva M.; Popivanova A.; Doicheva E.; Nikolov A. and Dimitrov A. (2003): Maternal insulin-dependent diabetes and congenital malformation in the newborn. Ahush Ginekol (Sofia) 42 (3): 3-5.
- Kalter H. (1990): Perinatal mortality and congenital malformations in infants born to women with IDDM-United states. Canada and Europe JAMA. 264, 4:437.
- Kanaka G.C.; Mastorakas G. and Chrousos G.P/ (2003): Endocrine-Related causes and consequences of intrauterine growth retardation. Ann NY Acad Sci. 997: 150-157.
- Kitzmiller J.L.; Brown E.R. and Phillippe M. (1981): Diabetic nephropathy and perinatal outcome. Am. J. Obstet Gynecol.141: 741.
- Kitzmiller J.L. Gavin L.A/ and Gin G.D. (1991): Preconception care of diabetes. Glycaemic control prevents congenital anomalies. JAMA. 13: 265 (6) : 731-6.
- Landy H.J.; Isada N.P. and Larsen J.W. (1987): Genetic implications of idiopathic hydramnios. Am. J. Obstet Gynecol. 157: 114-117.
- Langer O. Radriquez D.A.; Xenakis E.M.J.; McFarland M.B.; Berkus M.D. and Arrendondo F. (1994): Intensified versus conventional

- management of gestational diabetes. *Am. J. Obstet Gynecol* 170:1036.
- Langer I.; Conway D.L.; Berkus M.D.; Xenakis EM-J. and Gonzales O. (2000): A comparison of glyburide and insulin in women with gestational diabetes mellitus *N Engl J. Med.* 343: 1134-1138.
- Latini G.; De Felice C.; Parrini S.; Verroti A.; Di Maggio G. and Petraglia F. (2002): Polyhydramnios: a predictor of severe growth impairment in achondroplasia. *J. Pediatr.* 141 (2): 274-6.
- Magee M.S.; Walden C.E.; Benedetti T.J. and Knopp R.H. (1993): Influence of diagnostic criteria on the incidence of gestational diabetes and prenatal mortality. *JAMA* 26: 609.
- Major C.A.; Henry M.J.; De Veciana M. and Morgan M.A. (1998): The effect of carbohydrate restriction in patients with diet-controlled gestational diabetes. *Obstet. Gynecol.* 91: 600-604.
- Mami C. Paolata A.; Manganaro R.; Lo Monaco I. and Gemelli M. (2000): Diabetes and pregnancy. Maternal risk factors and neonatal morbidity. *Minerva Ginecol.* 52 (6): 235-41.
- Manning F.A.; Platt D.L. and Sipos L. (1980): Antepartum fetal evaluation: Development of a fetal biophysical profile. *Am. J. Obstet Gynecol.* 136: 787.
- Manning F.A.; Hill L.M. and Platt L.D. (1981): Quantitative amniotic fluid volume determination by ultrasound: Antepartum detection of intrauterine growth retardation. *Am. J. Obstet Gynecol.* 139: 254-8.
- Many A.; Hill L.M. and Lazebnik N. (1995): The association between polyhydramnios and preterm delivery. *Obstet Gynecol.* 86: 389-391.

- Maser R.E.; Lenhard M.J.; Henderson B.C.; Cobb R.S. and Hands K.E. (2004): Detection of episodes of gestational diabetes mellitus; A need for specific guidelines. *J. Diabetes complications* 18 (2): 86-90.
- Massin M.M.; Van Elmbt G. and Soyeur D. (1999): Reversible hypertrophic cardiomyopathy in congenital hyperinsulinism. *Acta Cardiol.* 54: 359-61.
- McElduff A. (2003): Shared care: Gestational diabetes Aus Fam Physician. 32 (3): 113-7.
- Metzger B.E. and Coustan D.R. (1998): Proceedings of the Forth International Work-shop-Conference on Gestational Diabetes Mellitus. *Diabetes Care* 21 (Suppl.): B1-B176.
- Metzger B.E.; Bybee H.M.; Freinkel N.; Phleps R.L.; Radvany R.M. and Vaisrub N. (1985): Gestational diabetes mellitus: correlation between the phenotypic and genotypic characteristics of the mother and abnormal glucose tolerance during the first year postpartum. *Diabetes* 34: 111.
- Molsted – Pederson L. and Pederson J.F. (1985): Congenital malformations in diabetic pregnancies *Acta Pediatr. Scand.* 320 (suppl.) 13-6.
- Moore T.R. and Clayle J.E. (1990): The amniotic fluid index in normal human pregnancy. *Am. J. Obstet Gynecol.* 162: 1168-73.
- National Diabetes Data Group (1979): Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes*, 28:1039. (Quoted by Donald R.) Causton Brom Book Medical Complications during pregnancy.
- Nelson R.L. (1983): Diabetes and pregnancy. primary care, 10 (2): 225-40.

- Nwsou E.C.; Leilch C.R and Marasse P.R. (1993): Longitudinal assessment of amniotic fluid index. *Br. J. Obstet Gynecol.* 100: 816-819.
- Odar E.; Wandobwa J. and Kiondo P. (2004) Maternal and fetal outcome of gestational diabetes mellitus in Malago Hospital, Uganda. *Afr Health Sci.* 4 (1) : 9-14.
- Ostlund I.; Hanson U.; Bjorklund A.; Hjertberg R.; Eva N.; Nordlander E.; Swahn M.L. and Wager J. (2003) Maternal and fetal outcomes if gestational impaired glucose tolerance is not treated. *Diabetes care.* 26 (7): 2107-11.
- Owen J.; Phelan S.T.; Landon M.P. and Gabbe S.G. (1995): Gestational diabetes survey. *Am. J. Obstet Gynecol.* 172: 615.
- Panting-Kemp A.; Nguyen T. and Castrol L. (2002): Substance abuse and poyhydramnios. *Am. J. Obstet Gynecol.* 187: 602-5.
- Pauer H.U.; Viereck V.; Krauss V.; Osmers R. and Krauss T. (2003): Incidence of fetal malformations in pregnancies complicated by oligo-and Poyhydramnios. *Arch Gynecol. Obstet.* 286 (1): 52-6.
- Pedersen J. and Jorgensen G. (1950): Hydramnios in diabetics: amount of amniotic fluid in relation to treatment. *Acta Endocrinal* 15: 333-41.
- Peevy K.J.; Landaw S.A. and Gross S.A. (1980): Hyperbilirubinaemia in infants of diabetic mothers. *Pediatrics* 66-417
- Phelan J.P.; Smith C.V.; Broussdrd P. and Small M. (1987): Amniotic fluid volume assessment with the four-quadrant technique at 36-42 weeks' gestation. *J. Reprod Med.* 32: 540-2.
- Pritchard J.A. (1966): Fetal swallowing and amniotic fluid volume. *Obstet Gynecol.* 28: 606-610.

- Queenan J.T. and Gandov E.C. (1970): Polyhydramnios : chronic versus acute. *Am. J. Obstet Gynecol.* 108:349.
- Reginold C.; Tsany M.B.; Jeanne B.S. and Colleen B. (1981): The infant of diabetic mother, today and tomorrow. *Clinical Obstet and Gynecol.* 24 1 P. 125-142.
- Robert A. and Brace R.A. (1997): Physiology of amniotic fluid volume and regulation. *Clinical Obstet and Gynecol.* 40 (2): pp 280.
- Robson S.C.; Robin John A.S. (1992): Intrapartum amniotic fluid index and its relationship to fetal distress. *Am. J. Obstet Gynecol.* 166: 78-82.
- Rochelson B.; Wagner J. and Shmoys S. (1992): The clinical significance of resolving polyhydramnios. *Ultrasound Obstet Gynecol.* 2 (5): 321-4.
- Rosenn B.; Miodounik M. and Combs A. (1991): pre-conception management of IDDM: Improvement of pregnancy outcome. *Obstet Gynecol.* 77: 846-51.
- Sarkar S.; Watman J.; Seigel W.M. and Schaeffer H.A. (2003): A prospective controlled study of neonatal morbidities in infants born at 36 weeks or more gestation to women with diet controlled gestational diabetes. *J. Perinatal.*, 23 (3): 223-8.
- Sarno A.P.; Ahn M.O. and Brar H.S. (1989): Intrapartum Doppler velocimetry, amniotic fluid volume and fetal distress. An initial report. *Am. J. Obstet Gynecol.* 161: 1508-14.
- Schoetzau A. and Hillebrand B. (1990): Neonatal morbidity of children of diabetic mothers. *Z Geburtshilfe Perinatal.* 194 (2): 58-64.
- Seeds A.E. (1980): Current concepts of amniotic fluid dynamics. *Am. J. Obstet Gynecol.* 138 (5): 575 - 586.

- Sepulveda W. Alcala J.L.; Schnapp C. and Bravo M. (2003): Prenatal outcome after prenatal diagnosis of placental chorioangioma. *Obstet Gynecol.* 102: 1028-33.
- Siddiqi B.; Rasenn B. and Mimounil F. (1991): Hypertension during pregnancy in insulin-dependant diabetic women. *Obstet Gynecol.* 77. 4P. 514.
- Spellacy W.N.; Buhi W.C.; Bradley B. and Holsinger K.K. (1973): Maternal, fetal and amniotic fluid levels of glucose, insulin and growth hormone. *Obstet Gynecol.* 41: 323-31.
- Steel, R.R.D. and Torrie J.H. (1982). *Principles of procedures of statistics.* Mc. Graw – Hill International book company 3rd Ed., London, P. 633.
- Stell J.M.; Johnstone F.D. and Hepburn D.A. (1991): Can pregnancy care of diabetic women reduce the risk of abnormal babies?. *BMJ.* 10-103 (6760): 1070-4.
- Stenninger E. Scholin J. and Aman J. (1991): Neonatal macrosomia and hyperglycemia in children of mothers with insulin treated gestational diabetes mellitus. *Acta.Pediatr. Scand.* 80 (11): 1014-8.
- Stoll C.; Alembic Y.; Dott B. and Roth M.P. (2001): Risk factors in congenital abdominal wall defects (omphalocele and gastro schisi): A study in a series of 265, 858 consecutive births. *Ann Genetic* 44 (4): 201-8.
- Stratton R.J. and Patterson R.M. (1993): DNA confirmation of congenital myotonic dystrophy in non-immune hydrops fetalis. *Prenat. Diagn.* B: 1027-1030.

- Sun D.J.; Lee J.N.; Long C.Y.; and Tsai E.M. (2003): Early diagnosis of fetal sacrocooygeal teratoma. *Kaohsiung J. Med Sci* 19 (6): 313-6.
- Taricco E.; Radaelli T.; Nobile de Santis M.S. and Cetin I. (2003): Fetal and placental weights in relation to maternal characteristics in gestational diabetes. *Placenta* 24 (4): 343-7.
- Tongsong T.; Sirichotiyakul S.; Wanapirak C. and Chnprapaph P. (2002): Sonographic features of trisomy 18 at mid pregnancy. *J. Obstet. Gynecol. Res.* 28 (5): 245-50.
- Tsang R.C.; Balmord J. and Braun C. (1981): The infant of diabetic mother: Today and tomorrow. *Clin-Obestet. Gynecol.* 24 (1): 135-147.
- Uvena-Celebrezze J.; Fung C.; Thomas A.J.; Hoty A.; Huston-Presley L.; Amini S.B. and Catalano P.M. (2002): Relationship of neonatal body composition to maternal glucose control in women with gestational diabetes mellitus *J. Marten Fetal Neonatal Med.* 12 (6): 396-401.
- Veille J.C.; Sivakoff M.; Hanson R. and Fanaroff A.A.(1992): Interventricular septal thickness in fetuses of diabetic mothers. *Obestet. Gynecol.* 79(1): 51-4.
- Verspyck E.; Mandelbrot L. and Dommergues M. (1993): Myasthenia gravis with polyhydramnios in the fetus of an asymptomatic mother. *Prent. DIAGN* 13: 539-542.
- Wax J.R.; Costigan R.N. and Callan N.A.. (1993): Effect of fetal movement on the amniotic fluid index. *Am. J. Obstet. Gynecol.* 168: 188-9.
- Weintrob N.; Karp M. and Hold. M (1996): Short and long-range complications in off springs of diabetic mothers. *Diabetes complications* 10 (5): 294-301.

- Weith P.A.; Hofmann H.; Winter R.; Purstner P. and Lichteneberger W. (1985): Amniotic fluid glucose values in normal and abnormal pregnancies. *Obstet Gynecol.* 65: 333-9
- Wendel G.D.; Maberry M.C. and Christmas J.T. (1989): examination of amniotic fluid in diagnosing congenital syphilis with fetal death. *Obstet. Gynecol.* 74: 967-970.
- Windness J.A.; Susa J.B. and Garcia J.F. (1981): Increased erythropoiesis and evaluated erythropoiesis in infants born to diabetic mothers in hyperinsulinemic rhesus fetuses. *J. Clin. Invest.* 67-637.
- Wladimiroff J.w.; and Campbell S. (1974): Fetal urine production rates in normal and complicated pregnancy. *Lancet.* 2: 151-154.
- Wood S.L.; Jick H. and Sauve R. (2003): The risk of stillbirth in pregnancies before and after the onset of diabetes. *Diabet. Med.* 20 (9): 703-7.
- Yankowitz J.; Weirner C.P. and Henderson L. (1996): Outcome of low-risk pregnancies with evidence of intraamniotic viral infection detected by PCR on amniotic fluid obtained at second trimester genetic amniocentesis. *J. Soc. Gynecol. Invest.* 3: 132.
- Yasuhi I.; Ishimaru T.; Hirai M. and Yamabe T. (1994): Hourly fetal urine production rate in the fasting and the post prandial state of normal and diabetic pregnant women. *Obstet Gynecol.* 84: 64.