

REFERENCES

Amos, N.N.; Stabinsky, S.A. et al. (1998): Uterine thermal therapy for the treatment of menorrhagia. Br. J. Obstet. Gynaecol. 105 : 517 – 523.

Abbott, J.A. and Garry, R. (2002): The surgical management of menorrhagia. Hum. Reported Update, 8 (1) : 68 – 78.

Abdel-Aleem, H.; El-Nashar, I. and Abdel-Aleem, A. (2001): management of severe postpartum hemorrhage with misoprostol. Intl. J. Gynecol. Obstet., 72 : 75 -6.

ACOG Practice Bulletin (2001): Clinical management guidelines for obstetrician – gynecologists. Use of botanicals for management of menopausal symptoms. Obstet Gynecol., 96 (6 suppl.): 1 – 11.

Akkad, A.A.; Habiba, M.A.; Ismail, N.; Abrams, K. and Al-Azzawi, F. (1995): Abnormal uterine bleeding on hormone replacement: The importance of intrauterine structural abnormalities. Obstet. Gynecol., 86 : 330 – 4.

Alexander, S.P.H. & Peters, J.A. (1998): Receptor and ion channel nomenclature supplement.. Trends Pharmacol. Sci. (Ninth edition. Pp) , Elsevier, Salna Ltd. 64 – 66.

Alisa B. Goldberg, M.D., Mara B. Greenberg, B.S., and Philip D.

Darney, M.D.(2001) Misoprostol and Pregnancy Volume
344:38-47January 4.

Amant, F.; Spitz, B.; Timmerman, D. et al. (1999) : Misoprostol
compared with methylergometrine for the prevention of
postpartum hemorrhage : a double – blind randomized trial.
Br. J. Obstet. Gynecol., 106 : 1066 – 70.

Anderson J.(1998) : Factors in fibroid growth. Baillieres clin
obstet gynaecol ; 12:225-43.

Apter, D. and Vihko, R. (1983): Early menarche, a risk factor for breast
cancer, indicates early onset of ovulatory cycles. J. Clin.
Endocrinal, 57 : 82 -6.

Baggish MS, Sze EHM.(1996): Endometrial ablation: a series of 568
patients treated over an 11-year period. Am J Obstet
Gynecol , 174:908-913.

Baldwin RM, Whalley PJ, Pritchard JA.(1961): Measurement
of menstrual blood loss. AJOG :81:739-42.

Bamigboye, A.A.; Hofmeyer, G.J. and Merrell, D.A. (1998): Rectal
misoprostol in the prevention of postpartum hemorrhage : a

placebo-controlled trial. Am. J. Obstet. Gynecol., 179 : 1043-6.

Bamigboye, A.A.; Merrell, D.A.; Hofmeyer, G.J. et al. (1998): Randomized comparison of rectal misoprostol with syntometrine for the management of third stage of labor. Acta Obstet. Gynecol. Scand., 77 (2) : 178 – 81.

Baughan, D.M. (1993): Changes in the management of patients with dysfunctional uterine bleeding. Fam Pract Recertification, 15 : 68 – 78.

Bawle, E.V.; Conard, J.V. and Weiss, L. (1998): Adult and two children with fetal methotrexate syndrome. Teratology, 57 : 51 – 55.

Bayer S and DeCherney A.H.(1993): Clinical manifestations and treatment of dysfunctional uterine bleeding. JAMA, 269:1823-8.

Beazley J.M. (1986) : natural labour and its active management. Dewhurst's text book of obstet and Gynaecol for post graduates 4th edition, whitfield C.R.;P. 324-46.

Bergstorm, Danniellsson, H. and Sjovall, J. (1962):The structure of prostaglandins E1, F1 and F2. Acta chem.. Scand., 16 : 501

– 510. Quoted from Bruce S.L., Paul, R.H.; Van Dorsten, J.P. (1982): Control of postpartum uterine atony by intramyometrial prostaglandin. *Obstet. Gynaecol.*, 59 : 47S – 50S.

Blanchard, K.; Clark, S.; Winikoff, B.; Gaines, G.; Kabani, G. and Shannon, C. (2001): Review of the current data on misoprostol for reproductive health indications. In submission *Obstetrics and Gynecology*.

Blanchard, K.; Winikoff, B. and Ellertson, C. (1999): Misoprostol used alone for the termination of early pregnancy : a review of the evidence. *Contraception*, 59 : 209 – 217.

Boie, Y.; Sawyer, N.; Slipetz, D.M.; Metters, K.M. and Abramovitz, M. (1995): Molecular cloning and characterization of the human prostanoid DP receptor.. *J. Biol. Chem.* 270 : 18910 – 18916.

Bonnar J.(1994): Dysfunctional uterine bleeding. Presentation at the Royal Society of Medicine, London, Feb 1994. London: The Royal Society.

- Bonnar J, Sheppard BL.(1996) :** Treatment of menorrhagia during menstruation: randomised controlled trial of ethamsylate, mefenamic acid, and tranexamic acid. BMJ ;313:579–582.
- Bond, G.R. and Van Zee, A. (1994):** Overdosage of misoprostol in pregnancy. Am. J. Obstet. Gynecol., 171 : 561 – 562.
- Bonduella M, Walker JJ, Calder AA.(1991):** A comparative study of danazol and norethisterone in dysfunctional uterine bleeding presenting as menorrhagia. Postgrad Med J ; 67:833-6.
- Bradley and Linda L (2005):**Abnormal uterine bleeding. Lippincott Williams & Wilkins, Inc. Volume 30(10), October, pp 38-49.
- Bravender T, Emans SJ.(1999):** Menstrual disorders: dysfunctional uterine bleeding. Pediatr Clin North Am ;46(3):545-53.
- Breches ME, Monk TG, Good nough LT.** A standardized method for the calculation of blood loss. Transfusion 1997; 37: 1071 - 1074.
- Brenner, P.F. (1996):** Differential diagnosis of abnormal uterine bleeding. Am. J. Obstet. Gynecol., 175 (3 pt 2): 766 – 9.
- Breches ME, Monk TG, Good nough LT.(1997):** A standardized method for the calculation of blood loss. Transfusion , 37: 1071 -1074.

Brill AI.(1995): What is the role of hysteroscopy in the management of abnormal uterine bleeding? Clin Obstet Gynecol ; 38:319–345.

Brodt – Eppley, J. and Myatt, L. (1998): Changes in expression of contractile FP and relaxatory EP2 receptors in pregnant rat myometrium during late gestation, at labor, and postpartum. Biol. Reprod., 59 : 878- 883.

Brooks, P.G.; Serden, S.P. and Davos, I. (1991): Hormonal inhibition of the endometrium for resectoscopic endometrial ablation. Am. J. Obstet. Gynecol., 164 : 1601 -8.

Bullen, B.A.; Skriner, G.S.; Beitins, I.Z.; Von Mering, G.; Turnbull, B.A. and McAuthur, J.W. (1985): Induction of menstrual disorders by strenuous exercise in untrained women. N. Engl. J. Med., 312 : 1349 – 53.

Burbank F, Hutchins FL. (2000): uterine artery occlusion by embolization or surgery for treatment of fibroids : a unifying hypothesis transient uterine ischemia. JAAGL, 7(4):S1-49.

Cameron IT.(1992): Medical management of menorrhagia. Curr Opin Obstet Gynecol ,2: 136-40

Cameron IT, Leask R, Kelly RW.(1987) : The effects of danazol, mafenemic acid, norethisterone, and a

progesterone impregnated coil on endometrial prostaglandin concentrations in women with menorrhagia. Prostaglandins 1987; 34:99-110.

Cameron, I.T. and Norman, J.E. (1995): Endometrial biochemistry in menorrhagia. Prog. Report Med., 2 : 267 – 279.

Cameron, I.T. and Norman, J.E. (1995): Endometrial biochemistry in menorrhagia. Prog. Report Med., 2 : 267 – 279.

Campbell, S. and Cameron, I.T. (1998): The origins and physiology of menstruation. In : Cameron, I.T., Fraser IS, Smith SK (eds) Clinical Disorders of the Endometrium and Menstrual Cycle. Oxford : Oxford University Press, 13 – 30.

Canavan, T.P. and Doshi, N.R. (1999): Endometrial cancer. Am. Fam. Physician, 59 (11) : 3069 – 77.

Celik H and Sapmaz E(2003): use of a single preoperative dose of . Fertile Steril may; 79(5): 1207-10.

Chimbira, T.H.; Anderson, A.B.M.; Naish, C. and Cope, E. (1980): Reduction of menstrual blood loss by danazol in unexplained menorrhagia : lack of effect of placebo. BJOG, 1152 -8.

Chuong CJ, Brenner PF.(1996): Management of abnormal uterine bleeding. Am J Obstet Gynecol 1996; 175:787–792.

Claessens, E.A. and Cowell, C.A. (1981): Acute adolescent menorrhagia.

Am. J. Obstet. Gynecol., 139 : 277 – 80.

Clark, A.; Black, N.; Rowe, P. et al. (1995): Indications for and outcome of total abdominal hysterectomy for benign disease : A prospective cohort study. Br. J. Obstet. Gynaecol. 102 : 611- 620.

Clark, T.J.; Mann, C.H.; Shah, N.; Khan, K.S.; Song, F. and Gupta, J.K. (2002): Accuracy of outpatient endometrial biopsy in the diagnosis of endometrial cancer: a systematic quantitative review. BJOG, 109 : 313 – 21.

Cole, S.K.; Billewicz, W.Z. and Thgomason, A.M. (1971): Sources of variation in menstrual blood loss. J. Obstet. Gynaecol. Br. Commonw, 78 : 933 -9.

Cohen MA.(1980). Anemia and menstrual loss. Obstet gynaecol surge; 35-597-601.

Cole, S.K.; Billewicz, W.A. and Thomson, A.M. (1971): Sources of variation in menstrual blood loss. J. Obstet. Gynaecol. Br. Common, W., 78 (10): 933 – 9.

Coleman, R.A.; Smith, W.L. and Narumiya, S. (1994): International union of pharmacology classification of prostanoid receptors

: properties, distribution, and structure of the receptors and their subtypes. Pharmacol., Rev. 46 : 205 – 229.

Collins, P. (1990): Misoprostol discovery development, and clinical application. Med. Res. Rev., 10 : 149.

Collins, P.W. (1990): Misoprostol : discovery, development, and clinical application. Med. Res. Rev., 10 : 149 – 79.

Cook, C.M.; Spurrett, M.M. and Murray, H. (1999): A randomized clinical trial comparing oral misoprostol with synthetic oxytocin or syntometrine in the third stage of labor. Aust NZ. J. Obstet. Gynecol., 39 : 4 : 414-9.

Coulter, A.; Klassen, A.; MacKenzie, I.Z.; McPherson, K. (1993): Diagnostic dilatation and curettage: is it used appropriately ? BMJ, 306 : 236 – 239.

Creinin, M.D. and Darney, P.D. (1994): Methotrexate and misoprostol for early abortion. Contraception, 48 : 339 – 48. [Erratum, Contraception 1994, 49 : 99.

Cunningham FG, McDonald PC, Gant NF (eds).(1997): Williams's Obstetrics, 20th Edition, Appelton Lang, Norwalk, Connecticut , 779.

Cynthia L, Kilbourn MD, Christine S, Richard MD(2001): Diagnostic considerations and management options of menorrhagia. Postgraduate Medicine Vol 109 NO 1.

Danielsson, K.G.; Marions, L.; Rodriguez, A.; Spur, B.W.; Wong, P.Y. and Bygdeman, M. (1999): Comparison between oral and vaginal administration of misoprostol on uterine contractility. Obstet. Gynecol., 93 : 275 – 280.

Darney PD(2001): Misoprostol: a boon to safe motherhood...or not? Lancet, 358:682-683.

Darney PD.(1991): Absorption kinetics of misoprostol with oral or vaginal administration. Obstet Gynecol., 90:88-92.

Davey, D.A. (1986): Dysfunctional uterine bleeding in : Dewhurst's textbook of Obstet. Gynaecol. For postgraduates, 4th editions, Edited by CR. Whitefield Blackwell, Scientific publication Oxford, London, P. 624 – 645.

Davey, D.A. (1995): Dysfunctional uterine bleeding. In whitfield CR (ed) Dewhurt's text book of Obstet and Gynaecol., 5th edition, London : Blackwell science, 590 – 608.

Davey, D.A. (1995): Hypertensive disorders of pregnancy. In Whifield, C.R. (ed.) Dewhurst's textbook of obstetrics and gynecology for

postgraduates. 6th edition. London : Blackwell Scientific Publications P.P. 175 – 214.

Davis A, Godwin A, Lippman J, et al. (2000): Triphasic norgestimate-ethinyl estradiol for treating dysfunctional uterine bleeding. Obstet Gynecol , 96:913-920.

Diab, K.M.; Ramy, A.R. and Yehia, M.A. (1999): The use of rectal misoprostol as active pharmacological management of the third stage of labor. J. Obstet. Gynaecol. Res., 25 (5): 327 – 32.

Dijkhuizen, F.P.; Mol, B.W.; Brolmann, H.A. and Heintz, A.P. (2000): The accuracy of endometrial sampling in the diagnosis of patients with endometrial carcinoma and hyperplasia : a meta – analysis. Cancer, 8 : 1765 – 72.

Dilley, A.; Drews, C.; Miller, C. Lally, C.; Austin, H.; Ramaswamy, D. et al. (2001): Von Willebrand disease and other inherited bleeding disorders in women with diagnosed menorrhagia. Obstet. Gynecol., 97 : 630 -6.

Dockeray CJ, Sheppard BL, Bonnar J.(1989): comparison of danazol and mafenemic acid in treatment of established menorrhagia. BJOG 1989; 840-4.

Dray, F. and Heaulme, M. (1984): Prostaglandins of the E series inhibit release of noradrenaline in rat hypothalamus by a mechanism unrelated to classical α_2 adrenergic presynaptic inhibition. *Neuropharmacology*, 23, 457 – 462.

Drudy, L.; Sheppard, B.L. and Bonnar, J. (1991): Mast cells in the normal uterine and in dysfunctional uterine bleeding. *Eur. J. Obstet. Gynaecol. Repord. Biol.*, 39 : 193 – 231.

Drudy, L.; Sheppard, B.L. and Bonnar, J. (1994): Histamine concentration in plasma of menstrual and peripheral blood in patient with normal menstrual blood loss and in dysfunctional uterine bleeding. *J. Obstet. & Gynaecol.*, 14 : 435 – 437.

Duffy S.(1995): The tissue and thermal effects of electrosurgery in the uterine cavity. *Baillieres Clin Obstet Gynaecol* , 9:261-277.

Effective health care. The management of menorrhagia. No. 9.London: Churchill Livingstone, 1995.

Elaine N, Unemori, Mark E et al(1999): Relaxin stimulate expression of endothelia growth factor in normal human endometrial cells in vitro and is associated with menometrorrhagia in women. *Human reproduction* Vol 14, No. 3, 800-806.

El-Makhzangy IM, Hegab HM, Galal HM, Tappozada M.

(2003) : the effect of rectal misoprostol on menstrual blood loss in menorrhagia. Egypt J Fertil Steril 2003; 7(3):55-61.

El-Refaey, H. and Templeton, A. (1994): Early abortion induction by a

combination of mifepristone and oral misoprostol : a comparison between two dose regimens of misoprostol and their effect on blood pressure. Br. J. Obstet. Gynaecol., 101 : 792 – 796.

El-Refaey, H. and Templeton, A. (1995): Induction of abortion in the

second trimester by a combination of misoprostol and mifepristone: a randomized comparison between two misoprostol regimens. Hum Reprod., 10 : 475 – 478.

El-Refaey, H.; Calder, L.; Wheatley, D.N. et al. (1994): Cervical

priming with prostaglandin E1 analogues : gemeprost and misoprostol. Lancet, 343 : 1207 -9.

El-Refaey, H.; Hinshaw, K. and Templeton, A. (1993): The abortifacient

effect of misoprostol in the second trimester. Hum. Reprod., 8 : 1744 -6.

El-Refaey, H.; Nooh, R.; O'Brien, P. et al. (2000): The misoprostol third stage of labour study : a randomized controlled comparison between orally administered misoprostol and standard management. Br. J. Obstet. Gynaeco., 1207 : 1104 – 10.

El-Refaey, H.; O'Brien, P.; Morafa, W.; Walder, J. and Rodeck, C. (1997): Use of oral misoprostol in the prevention of postpartum hemorrhage. Br. J. Obstet. Gynaecol., 104 : 336 – 339.

El-Refaey, H.; O'Brien, P.; Morafa, W. et al. (1996): Misoprostol for third stage of labor. Lancet, [letter to the editor]; 347 : 1257.

El-Refaey, H.; Rajasekar, D.; Abdalla, M. et al. (1995) : Induction of abortion with mifepristone (Ru 486) and oral or vaginal misoprostol. N. Engl. J. Med., 332 : 983 -7.

Everett RB.(1999): Five-year review of endometrial ablation with the SideFire laser fiber. J Am Ass Gynecol Laparosc , 6:65-69.

Ewenstein, M.B. (1996): The pathophysiology of bleeding disorder presenting as abnormal uterine bleeding. An. J. Obstet. Gynaecol., 175 : 770 – 777.

Fayez, J.A. (1982): Dysfunctional uterine bleeding. Am. Fam. Physician, 25 : 109 – 15.

Ferenczy, A. and Gelfand, M.M. (1986): Hyperplasia versus neoplasia : two tracks for the endometrium. Contemp. OB/Gyn., 28 : 79 – 96.

Fleming et al. (1989): Recent review of menorrhagia. Journal of midwifery & Women's health, Vol. 47, No. 3, May/June.

Fleming, H.; Oster, A.G.; Pickel, H. and Fournelle, D.W. (1989): Arterio-venous malformations of the uterus. Obstet. Gynecol., 73 : 204 – 214.

Foote, E.F.; Lee, D.R.; Karim, A.; Keane, W.F. and Halstenson, C.E. (1995): Disposition of misoprostol and its active metabolite in patients with normal and impaired renal function. J. Clin. Pharmacol., 35 : 384 – 389.

Franks, S. (1995): Polycystic ovary syndrome [published correction appears in N. Engl. J. Med., 333 : 1435.

Farquhar C.(1992): management of dysfunctional uterine bleeding. Drugs ; 44:578-89.

Fraser, I. (1994): Menorrhagia – aprognostic approach to the understanding of causes and the need for investigations. Br. J. Obstet. Gynaecol., 101 : 3 – 7.

Fraser, I.S. (1981): Efficacy of mefenamic acid in patient with menorrhagia. J. Obstet. Gynaecology, 53 : 453 – 457.

Fraser, I.S. (1985): The dysfunctional uterus: dysmenorrheal and dysfunctional uterine bleeding. In : Shearman RP (ed) clinical reproductive endocrinology. Edinbergh : Chuchill Livingstone, 579 – 589.

Fraser, I.S. (1992): Hysteroscopy & Laparoscopy : In women with menorrhagia. Am. J. Obstet. Gynaecol., 162 : 1264 – 1269.

Fraser, I.S. (2000): Benefits and risks of steroidal contraception. In : Salamonsen L, editor. Hormones and women's health: the reproductive years. Amsterdam: Harwood Academic Publishers, 161 – 172.

Fraser IS, McCarron G.(1991): Randomized trial of 2 hormonal and 2 prostaglandin-inhibiting agents in women with a complaint of menorrhagia. Aust N Z J Obstet Gynaecol , 31: 66–70.

Fried A.(1995): transvaginal sonography of the uterus. In : Desai S, Allahbadia G(eds) infertility and transvaginal sonography: current coccepts, New Delhi: Jaypee Brothers : 208-211.

Galle PC, McRae MA.(1993) : Abnormal uterine bleeding.

Postgrad Med , 93:73-76.

Gannon MJ, Day P, Hammmadie N, Johnson N.(1996): A

new method for measuring menstrual blood loss and

its use in screening women before endometrial

ablation. BJOG ; 103(10):1029-33.

Gillespie, A. and Nichols, A. (1994): The value of hysteroscopy. Aust. N.

Z. J. Obstet. Gynaecol., 34 : 85 – 87.

Gimpelson RJ, Rappoid HO.(1988): A comparative study between

panoramic hysteroscopy with directed biopsies and dilatation

and curettage: a review of 276 cases. Am J Obstet Gynecol.

1988;158:489-492.

Glasser, M.H.; Zimmerman, J.D. (2003): HydroThermAblator system

for management of menorrhagia in women with submucous

myomas: 12 – to 20 – month follow-up. J. Am. Assoc.

Gynecol. Laparosc., 2003, No., 10 (4): 521-7.

Goldberg, G.; Tsalacopoulos, G. and Davey, D. (1982): A comparison of

endometrial sampling with the accurette and Vabra aspirator

and uterine curettage. S. Afr. Med. J., 61 : 114 – 116.

Goldhaber MK, Armstorng MA, Goldith IM et al(1993):Role of endometrial histopathology in menorrhagia of unknown origin. Am J Epidemiol; 138:508-521.

Goldstein, S.R.; Zeltser, I.; Horan, C.K.; Synder, J.R. and Schwartz, L.B. (1997): Ultrasonography – based triage for perimenopausal patients with abnormal uterine bleeding. Am. J. Obstet. Gynecol., 177 : 102 – 8.

Gonzalez, C.H.; Marques-Dias, M.J.; Kim, C.A. et al. (1998): congenital abnormalities in Brazilian children associated with misoprostol misuse in first trimester of pregnancy. Lancet, 351 : 1624 – 1627.

Goodman, A. (2000): Abnormal genital tract bleeding. Clin. Cornerstone, 3 : 25 – 35.

Gulmezoglu, A.M.; Villar, J.; Ngoc, N.N.; Piaggio, G.; Carroli, G.; Adetoro, L.; Abdel-Aleem, H.; Chang, L.; Hofmeyr, G.J.;Lumbiganon, P.; Unger, C.; Prendiville, W.; Pinol, A.; Elbourne, D.; El-Refaey, H. and Schulz, K.F. (2001): For the WHO collaborative group to evaluation misoprostol in the management of the third stage of labour : the WHO multicentre double – blind randomized controlled trial to

evaluate the use of misoprostol in the management of the third stage of labour. *Lancet.*, 358 : 689 – 695.

Gulmezolglu, A.M.; Forna, F.; Villar, J. and Hofmeyr, G.J. (2002):

Prostaglandins for prevention of postpartum haemorrhage (Cochrane Review). In : the Cochrane library, issue 2. Oxford, UK.

Hallberg, L. and Nilsson, L. (1964): Determination of menstrual blood loss. *Scand. J. clin. Lab. Invest.*, 16 : 244 -8.

Hallberg, L.; Hgdahl, A.M.; Nilsson, L. and Rybo, G. (1966):

Menstrual blood loss – a population study. *Acta Obstet. Gynaecol. Scand*, 45 : 320 – 51.

Hawe JA, Phillips AG, Chien PF, et al.(1999): Cavaterm thermal balloon ablation for the treatment of menorrhagia. *Br J Obstet Gynaecol* , 106:1143-1148.

Hedin KE, Duerson K, Clapham DE.(1993) : Specificity of receptor–G protein interactions: Searching for the structure behind the signal. *Cellular Signaling.* ; 5:505–518.

Hickey, M., Higham, J., and Fraser, I.S. (2001) : Progestogens versus oestrogens and progestogens for irregular uterine bleeding associated

with anovulation (Cochrane Review). The Cochrane Library (Issue 1).

Oxford: Update Software.

Higham, J.M.; O'Brien, P.M.S. and Shaw, R.W. (1990): Assessment of menstrual blood loss using a pectorial chart. BJOG, 97 : 734-9.

Higham JM, Shaw RW. (1991): Risk-benefit assessment of drugs used for the treatment of menstrual disorders. Drug Saf ; 6:183–191.

Hill, N.C.; Oppenheimer, L.W. and Morton, K.E. (1989): The etiology of vaginal bleeding in children. A 20-year review. Br. J. Obstet. Gynaecol., 96 : 467 – 70.

Hillard, P.A. (1995): Heavy uterine bleeding in adolescents. Contemp. Ob/Gyn., 1 : 21 – 32.

Hofmeyer, G.J. and Bamigboye, A.A. (1999): In reply [letter to the editor]. Am. J. Obstet. Gynecol., 6 (pt 1): 1601 -2.

Hofmeyer, G.J.; Nikodem, V.C.; DeJager, M. et al. (1998): A randomized placebo controlled trial of oral misoprostol in the third stage of labor. Br. J. Obstet. Tynaecol., 105 (9) : 971 -5.

Hofmeyr, G.J. (1998): Misoprostol in obstetrics and gynaecology – unregistered, dangerous and essential. S. Afr. Med. J., 88 : 535 – 536.

Hofmeyr, G.J. and Gulmezoglu, A.M. (2003): Vaginal misoprostol for cervical ripening and induction of labour (Cochrane Review). In : The Cochrane Library Chichester, UK: John Wiley & sons, Ltd, (4).

Hofmeyr, G.J.; Gulmezoglu A.M. (2003): Vaginal misoprostol for cervical ripening and induction of labor; Cochrane database Syst. Rev. (1) : CD00941.

Hofmeyr, G.J.; Walraven, G.; Maholwana, B.; Gulmezoglu, A.M. and Villar, J. (2004): Misoprostol to treat postpartum haemorrhage : a systematic review of evidence relevant to efficacy, dosage and route of administration. Br. J. Obstet. Gynaecol., in press, P.D. Darney (Dept. of Obs. & Gyn. San Francisco, USA). Misoprostol : a boon to safe motherhood...or not? The Lancet, Vol. 358, Sept. 1, p. 682.

Hofmeyr, G.J.; Walraven, G.; Maholwana, B.; Gulmezoglu, A.M. and Villar, J. (2001): Misoprostol to treat postpartum haemorrhage : a systematic review of evidence relevant to

efficacy, dosage and route of administration. Goldberg A, Greenberg M, Darney P. Misoprostol and Pregnancy. New England. Journal of Medicine, 3441 : 38 – 47.

Hofmeyr, G.J.; Walraven, G.; Maholwana, B.; Gulmezoglu, A.M. and Villar, J. (2004): Misoprostol to treat postpartum haemorrhage : a systematic review of evidence relevant to efficacy, dosage and route of administration. Br. J. Obstet. Gynaecol., in Press.

Hofmeyr, G.J.; Walraven, G.; Maholwana, B.; Gulmezoglu, A.M. and Villar, J. (2004): Misoprostol to treat postpartum haemorrhage : a systematic review of evidence relevant to efficacy, dosage and route of administration. Br. J. obstet. Gynaecol., in Press. P.D Darney (Dept. of Obs. & Gyn. San Francisco, USA).

Hofmeyr, G.J.; Walraven, G.; Maholwana, B.; Gulmezoglu, A.M. and Villar, J. (2004): Misoprostol to treat postpartum haemorrhage : a systematic review of evidence relevant to efficacy, dosage and route of administration. Br. J. Obstet. Gynaecol., in Press.

Hornstein MD, Surrey ES, Weisberg GW, Casino LA. Leuprolide (1998): acetate depot and hormonal add-back in endometriosis: a 12-month study. Lupron Add-Back Study Group. *Obstet Gynecol* , 91:16-24.

Hourihan, H.M.; Sheppard, B.L. and Bonnar, J. (1986): A morphometric study of the effect of oral norethisterone and levonorgestrel in endometrial blood vessels. *Contraception*, 34 : 603 – 612.

House Committee on Government Reform and Oversight (1996): Off-Label drug use and FDA review of supplemental drug applications: hearing before the Subcommittee on Human Resources and Intergovernmental Relationship. 104th Congress, 2nd session, September, 12, 53 : 94.

Inman, W.H. (1991): Report on current PEM studies : drugs for peptic ulcer or reflux. *Prescription Event. Monitoring News*, 7 : 32 – 4.

Irons, D.W.; Sriskandabalan, P. and Bullough, C.H. (1994): A simple alternative to parenteral oxytocics for the third stage of labor. *Int. J. Gynecol. Obstet.*, 46 (1) : 15 – 8.

Itoh, S.; Lu, R.; Bao, Y. et al. (1996): Structural determinants of substrates for the prostaglandin transporter PGT. *Mol. Pharmacol.*, 50, 736 – 742.

J. A. Arosh, S. K. Banu, P. Chapdelaine and M. A. Fortier (2003):Published online ahead of print September 18, 2003, 10.1210/en.2003-1007 . *Endocrinology* Vol. 145, No. 1 407-417.

Janssen, C.A.H.; Scholten, P.C. and Heintz, A.M.P. (1995): A simple visual assessment technique to discriminate between menorrhagia and normal menstrual blood loss. *Obstet. Gynaecol.*, 85 : 977 – 82.

Jing Song (2000): Use of misoprostol in obstetrics and gynaecology; review article – obstet & Gyn. Survey, 55 (8) : 503 – 10.

Johnstone, M. (1972): The cardiovascular effects of oxytocicdrugs. *Br. J. Anaesth*, 44 : 826 – 35.

Jones DED, Shackelford DP, Brame RG.(1999): Supracervical hysterectomy: back to the future? *Am J Obstet Gynecol* , 180:513-515.

Kadir, R.A.; Economides, D.; Sabin, C.A.; Owens, D. and Lee, C.A. (1998): Frequency of inherited bleeding disorders in women with menorrhagia. *Lancet*, 351 : 485 – 9.

- Karim A, Keane WF, Halstenson CE.(1995):** Disposition of misoprostol and its active metabolite in patients with normal and impaired renal function. *J Clin Pharmacol*;35:384-389.
- Kasonade JM. (1975) :** Effect of ethamsylate and amniocaproic acid on menstrual blood loss in women using intrauterine devices. *BMJ* 1975; iv:21-2.
- Kim, Y.M.; Tejani, N.; Chayen, B. et al. (1986) :** Management of the third stage of labor with nipple stimulation. *J. Reprod. Med.*, 31 (11) : 1033 -4.
- Kiriyama, M.; Ushikubi, F.; Kobayashi, T.; Hirata, M.; Sugimoto, Y. and Narumiya, S.(1997):** Ligand binding specificities of the eight types and subtypes of the mouse prostanoid receptors expressed in Chinese hamster ovary cells. *Br. J. Pharmacol.* 122 : 217 – 224.
- Klein, N.A. and Soules, M.R. (1998):** Endocrine changes of the perimenopause. *Clin. Obstet. Gynecol.*, 41 : 912 – 20.
- Krassas, G.E. (2000):** Thyroid disease and female reproduction. *Fertil. Steril.*, 74 : 1063 – 70.
- LaBerge PY, Fortin CA, Cooper JM, Fulop T. (1999):** 3-D bipolar global endometrial ablation: the technology. *Proceedings from*

the World Congress of Alternatives to Hysterectomy; February 26-28, , Miami, FL, USA. The American Association of Gynecologic Laparoscopists Publications Corp. 193.

Langer RD, Pierce JJ, O'Hanlan KA, Johnson SR, Espeland MA, Trabala JF, et al.(1997) : Transvaginal ultrasonography compared with endometrial biopsy for the detection of endometrial disease. N Engl J Med ;337:1792-8.

Liddell H.(1993) : menorrhagia. NZ Med J 1993;106:255-7.

Lethaby, A.; Farquhar, C.; Sarkis, A.; Roberts, H.; Jepson, R. and Barlow, D. (2003): Hormone replacement therapy in postmenopausal women: endometrial hyperplasia and irregular bleeding. Cochrane Database Syst. Rev., 4 : CD000402.

Lethaby A, Augood C, Duckitt K.(2004) : Nonsteroidal Anti-Inflammatory Drugs for Heavy Menstrual Bleeding (Cochrane Review). The Cochrane Library, Issue 3. Chichester, UK: Wiley, 2004

Livingstone, M. and Fraser, I.S. (2002): Mechanisms of abnormal uterine bleeding. Hum. Reprod. Update, 8 : 60 – 7.

Lindoff C, Rybo G, Astedt B (1993): Treatment with tranexamic acid during pregnancy, and the risk of thrombo-embolic complications. *Thromb Haemost* , 70:238-240

Long, C.A. (1996): Evaluation of patients with abnormal uterine bleeding. *Am J. Obstet. Gynecol.*, 175 : 784 -6.

Lumbiganon, P.; Hofmeyr, J.; Gulmezoglu, A.M.;Pinol, A. and Villar, J. (1999): Misoprostol dose – related shivering and pyrexia in the third stage of labour. *Br. J. Obstet. Gynaecol.*, 106 : 304 – 308.

Lumsden, M. and Smith, S. (1992): Menstruation and menstrual abnormality in : textbook of Gynaecol. Ed., by Robert W Shaw, Soutter WP and Santon, SL, 1st edition. Churchill livingstone.

Lyons, C.; Beharry, K.; Akmal, Y.; Attenello, F. and Nageotte, M.P. (2003): In vitro response of prostaglandin E2 receptor (EP3) in the term pregnant rat uterus and cervix to misoprostol. *Prostaglandins Other Lipid Mediat*, 70 : 317 – 329.

MacKenzie, I.Z. and Bibby, J.G. (1978): Critical assessment of dilatation and curettage of 1029 women. *Lancet*, 2 : 566 – 568.

March, C.M. (1998): Bleeding problems and treatment. Clin. Obstet. Gynecol., 41 : 928 – 39.

Mark, A.S.; Hricak, H.; Heinnrichs, L.W. et al. (1987): Adenomyosis and leiomyoma – differential diagnosis with MR imaging. Radiology; 163 : 527.

Matsumoto T, Sagawa N, Yoshida M, Mori T, Tanaka I, Mukoyama M, Kotani M, Nakao K (1997) : The prostaglandin E2 and F2 α receptor genes are expressed in human myometrium and are down-regulated during pregnancy. Biochem Biophys Res Commun 238:838–841

McKone, E.; Sheppard, B.L. and Bonnar, J. (1991): Uterine histology and menstrual blood loss. Irish, J. Med. Sci., 160 : 359 – 360.

Miller RM, Frank RA.(1992): Zoladex (Goserlin) in treatment of benign gynecological disorders :an overview of safety and efficacy. BJOG ;99:37-41.

Milsom I, Anderson K, Andersch B, Rybo G. A. (1991): comparison of flurbiprofen, tranxemic acid, and a levonorgestrel releasing intrauterine contraceptive device in the treatment of idiopathic menorrhagia. AJOG ; 164:879-83.

- Mihm, L.M.; Quick, V.A.; Brumfield, J.A.; Connors, A.F. Jr. and Finnerty, J.J. (2002):** The accuracy of endometrial biopsy and saline sonohysterography in the determination of the cause of abnormal uterine bleeding. *Am. J. Obstet. Gynecol.*, 186 : 858 – 60.
- Mishell, D.R. Jr. (2001):** Abnormal uterine bleeding. In :Stenchever MA, Droegmueller W, Herbst, AL, Mishell, DR Jr, editors. *Comprehensive gynecology*, 4th ed. St Louis (MO): Mosby: 1079.
- Munro MG.(2001) :** Abnormal uterine bleeding in the reproductive years. Part II—medical management. *J Am Assoc Gynecol Laparosc* 2000;7:19–32.
- Munro MG.(2000):** Medical management of abnormal uterine bleeding. *Obstet Gynaecol Cli North Am* , 27:287-304.
- Narumiya, S. (1994):** Prostanoid receptors: Structure, Function, and distribution. (In) cellular generation, transport, and effects of eicosanoids : biological roles and pharmacological intervention (ed. Goetzl, E.J. Lewis, R.A. and Rola –

Pleszczynski, M. pp) 126 – 138 New York : The New York Academy of Sciences.

Narumiya, S.; Sugimoto, Y. and Ushikubi, F. (1999): Prostanoid receptors: structures, properties, and functions. *Physiol, Rev.*, 79 : 1193 – 1226.

Neese, R.E. (1989): Abnormal vaginal bleeding in perimenopausal women. *Am. Fam. Physician*, 40 : 185 – 92.

Ng, P.S.; Chan, A.S.M.; Sin, W.K. et al. (2001): A multicentre randomized controlled trial of oral misoprostol and I.M. syntometrine in the management of the third stage of labour. *Hum. Reprod.* 16 (1) : 31 -5.

Nilsson L, Rybo G(1997): Treatment of menorrhagia. *Am J Obstet Gynecol.* ;110:713-720.

Noyes RW, Hertig AW, Rock J(1950): Dating the endometrial biopsy. *Fertil Steril*; 1:3-25.

O'Brien, P.; El-Refaey, H.; Gordon, A.A.A. et al. (1998): Rectally administered misoprostol for the treatment of postpartum hemorrhage unresponsive to oxytocin and ergometrine : a descriptive study. *Obstet Gynecol.*, 92 (2) : 212 -4.

Olive DL. (1991): The prevalence and epidemiology of luteal phase deficiency in normal and infertile women. *Clin Obstet Gynaecol*; 34:157-66.

Oriel, K.A. and Schrager, S. (1999): Abnormal uterine bleeding. *Am. Fam. Physician*, 60 : 1371 – 80.

Orioli, I.M. and Castilla, E.E. (2000): Epidemiological assessment of misoprostol teratogenicity. *Br. J. Obstet. Gynaecol.*, 107 : 519 -523.

Park KH, Kim JY, Shin JS, KnownJY, Koo JS, Jeong KA et al. (2003): Treatment and outcome of uterine artery embolization and laparoscopic uterine artery ligation for uterine myoma. *Yonsei Med J*, 44(4): 964-702.

Pastuszek, A.L.; Schuler, L.; Speck – Martins, C.E. et al. (1998): Use of misoprostol during pregnancy and Mobius syndrome in infants. *N. Engl. J. Med.*, 338 : 1881 – 1885.

Petrozza, J. and Poley, K. (1999): Dysfunctional uterine bleeding. In : Curtis MG, Hopkins MP, eds. *Glass's office gynecology*. 5th ed. Baltimore: Williams & Wilkins, 241 – 64.

Pickles, V.R.; Hall, W.J.; Best, F.A. and Smith, G.N. (1965): Prostaglandins in endometrium and menstrual fluid from normal and dysmenorrhoeic subjects. *J. Obstet. Gynaecol. Br. Commonw*, 72 : 185-1991.

Pittrof R, Majid S, Murray A.(1994): Transcervical endometrial cryoablation for menorrhagia. Int Obstet Gynaecol , 47:135-40.

Raisz, L.G. and Woodiel, F.N. (1989): Effect of alterations in the cyclopentane ring on bone resorptive activity of prostaglandin. Prostaglandins 37, 229 – 235.

Ramsey, P.S.; Ramin, K.D.; Bradle, S.S. (1999): Rectal misoprostol in the prevention of postpartum hemorrhage [letter to the editor]. Am. J. Obstet. Gynecol., 6)pt 1): 1601.

Rankin GLS, Veall N, Huntsman RG, LIdell J.(1998): measurement with Cr of red cell loss in menorrhagia. Lancet 1962;1:567-9.

Ramwell, P.W.; Foegh, M. Loeb, R. and Leovey EMK (1980): Synthesis and metabolism of prostaglandins, prostacyclins and thromboxane: the arachidonic acid cascade; seminars Perinatol., 4 : 3 .

Rangachari, P.K.; Betti, P.A.; Prior, E.T. and Roberts, L.J. (1995): Effects of a selective DP receptor agonist (BW 345 C) and antagonist (BW A868C) on the canine colonic epithelium;

an argument for a different DP receptor ? J. Pharmacol. Exp. Ther 275 : 611 – 617.

Rybo G.(1991) : Tranxemic acid thrapy is effective treatment in heavy menstrual bleeding; clinical update on safety. Ther Adv ; 4:1-8.

Rayburn, W.F. (1993): A physician's prerogative to prescribe drugs for off – label uses during pregnancy. Obstet. Gynecol., 81 : 1052 – 1055.

RCOG (1998) : The initial management of menorrhagia. Evidence-based clinical guidelines No. 1. London: Royal College of Obstetricians & Gynaecologists.

Reid RC, Wagner PL, Gambone JC.(1992) : Routin hysterectomy for large asymptomatic uterine liomyomata; a reappraisal. Obstet gynecol ;79(4):481-4.

Rueda (1991): Dysfunctional uterine bleeding a reappraisal Curr. Probl. Obstet. Gynaecol. Fertile. 14, 3 :79 – 96.

Rutherford TJ, Zreik TG, Troiano RN, et al.(1998): Endometrial cryoablation, a minimally invasive procedure for abnormal uterine bleeding. J Am Assoc Gynecol Laparosc , 5:23-28.

Rybo (1982): Variation in menstrual loss. Research and clinical forums, 4, 81 – 92.

Samulesson, B. (1963): The structure of prostaglandin E1. Am. J. Chem. Soc., 85: 1878. *Quoted from Bruce S.L.; Paul, R.H.; Van Dorsten, J.P. (1982):* control of postpartum uterine atony by intramyometrial prostaglandin. Obstet. Gynaecol., 59 : 47S – 50S.

Schultze (1975): Role of lysosomes in human endometrium. Am.J. Obstet Gynaecol. 123:650.

Schwarz BE (1983): The production and biologic effects of prostaglandins. Semin. Repord. Endocrinal ; 1:189.

Scott, J.C. and Massey, E. (1981): Menstrual pattern of myxedema. Am. J. Obstet. Gynaecol., 139 : 277 – 280.

Scott, J.R.; Cramer, D.W.; Peterson, H.B. et al. (1998): Reading the medical literature: applying evidence to practice. Washington, D.C.: American College of Obstetricians and Gynecologists, 1 – 8.

Senior, J.; Marshall, K.; Sangha, R. et al. (1993): In vitro characterization of prostanoid receptors on human

myometrium at term pregnancy. Br. J. Pharmacol., 108 : 501-6.

Shah AA, Grainger DA.(1996): Contemporary concepts in manganig menorrhagia. Medscape Women's Health, 1:12.

Shangold, M.; Rebar, R.W.; Wentz, A.C. and Schiff, I. (1990): Evaluation and management of menstrual dysfunctional in athletes. JAMA, 263 : 1665 – 9.

Sharma, S.C.; Sheppadrd, B.L. and Bonnar, J. (1990): Relationship of menstrual blood with uterine 6-keto-PGF1 and TXB2 levels in women with normal and dysfunctional uterine bleeding. Irish J. Med. Sci., 159 : 57.

Shaw RW.(1994) : Assessment of medical treatments for menorrhagia. Br J Obstet Gynaecol ; 101(Suppl 11):15–18.

Sheath & Allah – Buida (1999): Sheath S.S. adnexal pathology. In Sheath SS, Allahbadia. Diagnosis of menorrhagia. Oxford : Isis medical media, 58 : 124 – 134.

Sheppard, B.L.; Stack, M.; Jordan, M. and Bonnar, J. (1990): Plasmenogen activator in the human uerus in normal

menstruation and dysfunctional uterine bleeding. Irish J. Med. Sci., 159 : 151).

Sheth, S.S. (1991): Vaginal dimple: a sign to diagnose endometriosis. J. Obstet. Gynecol., 2 : 292.

Singh G, (1996): Gastrointestinal tract complications of nonsteroidal anti-inflammatory drug treatment in rheumatoid arthritis. A prospective observational cohort study. Arch Intern Med;156(14):1530-6.

Shwayder, J.M. (1999): Pathophysiology of abnormal uterine bleeding. Obstet. Gynecol. Clin. North. Am., 27 : 219 – 34.

Shwayder, J.M. (2000): Pathophysiology of abnormal uterine bleeding. Obstet. Gynecol. Clin. North. Am., 27 : 219 – 34.

Smith, W.L.; Garavito, R.M. and Dewitt, D.L. (1996): Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2 J. Biol. Chem. 271 : 33157 – 33160.

Smith S.K.; Abel M.H. (1982): The synthesis of prostaglandins from persistent proliferative endometrium. J.Clin. Endocrinal. Metab. 55:284-289.

Sowthern, E. (1972): Prostaglandin chemistry : In : The prostaglandins clinical application in human reproduction, 1st ed., Edited by future publishing Co. New York, USA, 26 – 30.

Speroff, L.; Glass, R. and Kase, N. (1999): Menopause and the perimenopausal transition. In : Speroff, L.; Glass, R.; Kase, N., editors. Clinical gynecologic endocrinology and infertility, 6th ed. Baltimore (MD): Lippincott Williams & Wilkins, 643 – 724.

Speroff, L.; Glass, R. and Kase, N. (1999): Anovulation and the polycystic ovary. In : Speroff, L., Glass, R., Kase, N., editors. Clinical gynecologic endocrinology and infertility, 6th ed. Baltimore (MD): Lippincott Williams & 6-Wilkins, 487 – 513.

Speroff, L.; Glass, R.H. and Kase, N.G. (1999): Clinical gynecologic endocrinology and infertility. 6th ed. Baltimore : Lippincott Williams & Wilkins, 201 – 38, 499, 575 -9.

Speroff, L.; Glass, R.H. and Kase, N.G. (1999): The endocrinology of pregnancy. In : Speroff L, Glass RH, Kase NG (ed) : Clinical gynaecologic endocrinologyand infertility 6th (ed); Williams& Wilkins. Blatimore, Mary Land, USA; P. 251.

Speroff, L.; Glass, R.H.; Kase, N.G. (1994): Dysfunction uterine bleeding. In : Clinical gynecologic endocrinology and infertility. 5th ed. Baltimore : Williams & Wilkins, 575 – 93.

Stabinsky, Seth A.; Einstein, Mark; Breen, James L.(1999): Modern treatments of menorrhagia attributable to dysfunctional uterine bleeding. Obstet Gynecol Surv. 1999 Jan;54(1):61-72. Review.

Stovall, T.G.; Ling, F.W. and Morgan, P.L. (1991): A prospective, randomized comparison of the Pipelle endometrial sampling device with the Novak curette. Am. J. Obstet. Gynecol., 165 (5 pt 1): 1287 – 90.

Studd, J.W.W. (1989): Hysterectomy and mwnorrhagia Baillieres Clin. Obstet. Gynaecol., 3 : 415 – 424.

Surbek, D.V.; Fehr, P.M.; Hosli, I. et al. (1999): Oral misoprostol for third stage of labor : a randomized placebo – controlled. Obstet. Gynecol., 94 (2): 255 – 8.

Syane, D.; Larson, B.; Andersson, K.E. and Forman, A. (1993): Endothelin -1 : immunocytochemistry, localization of binding sites, and contractile effects in human uteroplacental smooth muscle. Am. J. Obete. Gynaecol., 168 : 233 – 241.

Syrop C, Sahaklan V.(1992) : Transvaginal sonographic detection of endometrial polyps with fluid contrast augmentation. *Obstet Gynecol.* 79:1041-1043.

Tabor A, Watt HC, Wald NJ.(2002) : Endometrial thickness as a test for endometrial cancer in women with postmenopausal vaginal bleeding. *Obstet Gynecol* ;99:663-70.

Tang, O.S.; Schweer, H.; Seyberth, H.W.; Lee, S.W.H. and Ho, P.C. (2002): Pharmacokinetics of different routes of administration of misoprostol. *Human, Reproduction*, 17 : 332 – 336.

Tang, O.S.; Schweer, H.; Weyberth, H.W.; Lee, S.W.H. and Ho, P.C. (2002): Pharmacokinetics of different routes of administration of misoprostol. *Human Reproduction*, 17 : 332 – 336.

Taylor, P.T. Jr; Andersen, W.A.; Barber, S.R. et al. (1987): The screening papanicolaou smear contribution at endocervical brush. *Obstet. Gynaecol.*, 70 : 734 – 738.

Thomas EJ, Okuda KJ, Thomas NM.(1996): The combination of a depot gonadotrophin releasing hormone agonist and cyclical

hormone replacement therapy for dysfunctional uterine bleeding. BJOG, 103:18-21.

Thong, K.J. and Baird, D.T. (1992): Induction of abortion with mifepristone and misoprostol in early pregnancy. Br. J. Obstet. Gynaecol., 99 : 1004 – 1007.

Toppozada, M.K.; Anwar, M.Y.; Hassan, H.A. and El-Gazaerly, W.S. (1997): Oral or vaginal misoprostol for induction of labor. Int. J. Gynaecol. Obstet., 56 : 135 – 139.

Ulmann, A. and Silvestre, L. (1994): RU486 : the French experience. Hum Reprod. 9 : Suppl. 1 : 126 – 130.

USP Obstetrics and Gynecology Expert Advisory Panel Consensus on revision of misoprostol for prevention of postpartum hemorrhage, 6/2001.

Valle RF.(1981): Hysteroscopic evaluation of patients with abnormal uterine bleeding. Surg Gynecol Obstet 1981; 153:521-526.

Valle RD(1990): Hysteroscopic removal of submucous leiomyomas. J Gynecol Surg ;6:89-86

Van Dongen pw, De-Groot AN (1995): History of ergot alkaloid from ergotism to ergometrine. European journal of obstet and gynaecol Reproductive Biology; 60(2): 109-16.

Van den Bosch T, Vandendael A, Van Schoubroeck D, Wranz PA, Lombard CJ.(1995) : Combining vaginal ultrasonography and office endometrial sampling in the diagnosis of endometrial disease in postmenopausal women. Obstet Gynecol ;85:349-52.

Van Eijkeren M, Christiaens G, Haspels A, Sixma J.(1990): measured menstrual blood loss in women with a bleeding disorder or using oral anticoagulant therapy. AJOG ;162:1261-3.

Vargyas JM, Campeau JD, Mishell DA. (1987): Treatment of menorrhagia with meclofenamate sodium. Am J Obstet Gynecol ;157:944–950.

Vasilenko P, Crosignani PG, Mangoni C, Ferrari A, De Giorgi.(1998) : Treatment with gonadotropin releasing hormone agonist before hysterectomy for liomyoma: a multicentre, randomized controlled trial. BJOG 1998, 105 (11): 1148-54.

- Vercellini P, et al. Hysteroscopic myomectomy (1999):** long-term effects on menstrual pattern and fertility. *Obstet Gynecol* September ;94:341-7.
- Vessey M, Clarke J. MacKenzie I.(1979):** Dilatation and curettage in young women. *Health Bull* , 39: 59-62.
- Vessey, M.; Villard – Mackintosh, L.; McPherson, K. et al. (1992):** The epidemiology of hysterectomy : findings in a large cohort study. *Br. J. Obstet. Gynaecol.*, 99 : 402 – 407.
- Wall, D.M. and Roos, M.P. (1990):** Update on combination oral contraceptives. *Am. Fam. Physician*, 42 : 1037 – 48.
- Walley, R.L.; Wilson, J.B.; Crane, J.M.G. et al. (2000) :** A double – blind placebo controlled randomised trial of misoprostol and oxytocin in the management of the third stage of labour. *Br. J. Obstetrics Gynaecol.*, 107 : 1111 -5.
- Wellington, Keri ; Wagstaff and Antona(2003) :** Tranexamic Acid: A Review of its Use in the Management of Menorrhagia. *Adis Data Information BV Volume 63(13)*, pp 1417-1433
- Wentz, A.C. (1988):** Abnormal uterine bleeding in : *Novak’s Textbook of Gynaecology*. 11th edition. (Editor, Brown, C.L.) Williams & Wilkins, Baltimore, P. 378.
- Widrich T, Bradley LD, Mitchinson AR, Collins RL.(1996):** Comparison of saline infusion sonography with office

hysteroscopy for the evaluation of the endometrium. Am J Obstet Gynecol ;174: 1327-34.

Wiebe, E.R. (1997): Abortion induced with methotrexate and misoprostol: a comparison of various protocols. Contraception, 55 : 159 – 163.

Wing, D. (1999): Labour induction with misoprostol. Am. J. Obstet. Gynecol. 18 : 339 – 345.

Walthen PL, Henderson MC, Wintz CA.(1995) : abnormal uterine bleeding. Med Clin North Am ; 79:329-342.

Zieman, M.; Fong, S.K.; Benowitz, N.L.; Banskter, D. and Darney, P.D. (1997): Absorption kinetics of misoprostol with oral or vaginal administration. Obstet. Gynecol., 90 : 88 – 92.