



References

REFERENCES

- Agrawal R. (1996):* Pediatric anesthesia. In Duke J, Rosenberg SG (eds): Anesthesia Secrets. Philadelphia, Hanley & Belfus; P 369.
- Agro F, Brimacombe J, Carussiti M, Marchionni L. (1998):* The intubating laryngeal mask – clinical appraisal of ventilation and blind tracheal intubation in 110 patients. *Anaesthesia*; 53: 1084-1090.
- Agro F, Brimacombe J, Cassaiti M. (1998):* Use of a lighted stylet for intubation via the laryngeal mask airway. *Can J Anaesth*; 45 (6): 556-560.
- Agro F, Galli B, Ravussin P. (2002):* Preliminary results using the laryngeal tube for supraglottic ventilation. *Am J Emerg Med*; 20; 57-8.
- Akec O, Wadhwa A, Sengupta P, Durrani J, Hanni K, Wenke M, Yucel Y, Lenhardt R, Doufas AG and Sessler DI. (2004):* The new perilaryngeal airway (CobraPLA) is as efficient as the laryngeal mask airway (LMA) but provides better airway sealing pressures. *Anesth Analg*; 99 (1): 272-8.
- Akhtar TM, McMurray P, Kerr WJ. (1992):* A comparison of LMA with tracheal tube for intra ocular ophthalmic surgery. *Anesth*; 47: 688-671.
- Alan R, Tait, PHD, Uma A. (1998):* Use of the laryngeal mask airway in children with upper respiratory tract infections: A comparison with endotracheal intubation. *Anesth Analg*; 68: 706-11.
- Angeles Ayuso M, Xavier Sala, Merce Luis, Joan M, Carbo. (2003):* Predicting difficult orotracheal intubation in pharyngo-laryngeal disease: preliminary results of a composite index. *CAN J ANESTH*; 50: 1 PP 81-85.
- Annamoneni R, Hodzovic I, Wilkes AR, Lattop. (2003):* A comparison of simulated difficult intubation with multiple-use and single-use bougies in a manikin. *Anaesthesia*; 58: 45-9.

- Aoyama K, Yamamoto T, Takenaka I. (1998):* The jaw support device facilitates laryngeal exposure and ventilation during fiberoptic intubation. *Anaesth Analg*; 86: 432-434.
- Asui T, and Morris S. (1994):* The laryngeal mask airway, its features, effect and role. *Can. J. Anesthesia*; 41: 930-960.
- Asai T, Kawashima A, Hidaka I, Kawachi S. (2001):* Laryngeal tube: its use for controlled ventilation. *Masui*; 50: 1340-1.
- Asai T, Latta JP. (1995):* Use of the lighted stylet for tracheal intubation via the laryngeal mask airway (letter). *Br J Anaesth*; 75: 503-504.
- Asai T, Murao K, Shingu K. (2000):* Efficacy of the laryngeal tube during intermittent positive pressure ventilation. *Anaesthesia*; 55: 1099-102.
- Atikenhead A and Smith G. (1996):* Textbook of anesthesia, 3rd edition Churchill Livingstone. Anesthetic apparatus.
- Audenaert SM, Montgomery CL, Stone B. (1991):* Retrograde-assisted fiberoptic tracheal intubation in children with difficult airways. *Anaesth Analg*; 73: 660.
- Audu PB and Loomba N. (2004):* Use of cuffed oropharyngeal airway (COPA) for awake intracranial surgery. *J. Neurosurg Anesthesiol*; 16(2): 144-6.
- Barriot P, Riou B. (1988):* Retrograde technique for tracheal intubation in trauma patients. *Crit Care Med*; 16: 712-713.
- Basken PJF, Parr JA, Nolan JP. (1998):* The intubating laryngeal mask – results of a multicentre trial with experience of 500 cases. *Anaesthesia*; 53: 1174-1179.
- Bellhouse CP. (1987):* Management of the difficult airway. *Anaesth intensive Care*; 16:329-337.
- Benoumof JL. (1991):* Management of the difficult adult airway with special emphasis on awake tracheal intubation. *Anesthesiology*; 75: 1087

- Benumof JL, Cooper SD. (1996):* Quantitative improvement in laryngoscopic view by optimal external laryngeal manipulation. *J Clin Anaesth*; 8: 136-140.
- Benumof JL. (1991):* Management of the difficult adult airway: with special emphasis on awake tracheal intubation. *Anesthesiology*; 75: 1087-1110.
- Benumof JL. (1992):* Use of the laryngeal mask airway to facilitate fibroscope-aided tracheal intubation. *Anaesth Analg*; 74: 313-314.
- Bigeleisen PE. (2000):* Prophylactic placement of a cricothyrotomy catheter in a trauma patient with a suspected difficult airway. *Am J Anesthesiol*; 27 (2): 101-2.
- Bissinger U, Guggenberger H, Lenz G. (1995):* Retrograde-guided fiberoptic intubation in patients with laryngeal carcinoma. *Anaesth Analg*; 81: 408-410.
- Brain AIJ, Verghese C, Addy EV, Kapila A. (1997):* The intubating laryngeal mask I. Development of a new device for intubation of the trachea. *Br J Anaesth*; 79: 699-703.
- Brain E, and Duff MD. (1999):* Use of the laryngeal mask airway in otologic surgery. *Laryngoscope*; 109: 1033-1036.
- Brimacombe J, and Keller C. (1999):* Comparison of the flexible and standard laryngeal mask airways. *Can. J. Anesth*; 46: 6, PP 558-563.
- Brimacombe J, Berry (1994):* Active swallowing to aid LMA insertion in awake patients. *Anesth Analg*; 78: 1029.
- Brimacombe J, Berry A. (1993):* Laryngeal mask airway insertion: A comparison of the standard versus neutral position in normal patients with a view to its use in cervical spine instability. *Anaesthesia*; 48: 670-671.
- Brimacombe J, Berry A. (1995):* The incidence of aspiration with the laryngeal mask airway-a meta-analysis of published literature. *J Clin Anesth*; 7: 297-305.

- Brimacombe J, Brain AIJ, and Berry A. (1997):* Anesthetic management and neonatal resuscitations both in the LMA. A review and practical guide Ed by J. Brimacombe. AIJ Brain and A Berry (Ch. 6, 65-105) (Ch. 14, 174-199) and (Ch. 16, 216-226) Pub by W.B. Saunders Comp., London and Tokyo.
- Brimacombe J, Brain AIJ, Branagan H, Spry M, Schotfield J. (1996):* Optimal shape of the laryngeal mask cuff: the influence of three deflation techniques. *Anaesthesia*; 51: 673-676.
- Brimacombe J, Keleir C, Kurian S, Myles. (2000):* Reliability of epigastric auscultation to detect gastric insufflation. *Br J Anaesth*; 88: 127-9.
- Brimacombe J. (1996):* The laryngeal mask airway for access of the upper gastrointestinal tract. *Anesthesiology*; 84: 1009-1010.
- Captan RA, Benumof JL, Berry FA. (2003):* Practice guidelines for management of the difficult airway. A report by the American Society of Anaesthesiologists. Task force on management of the difficult airway. *Anesthesiology*; 78: 597-602.
- Casati A, Fanelli G, Casuletti E. (1999):* Do we need new supraglottic devices? Clinical appraisal of the cuffed oropharyngeal airway (COPA). *Minerva Anestesiol*; 65 (12): 823-830.
- Chadd GD, Grave DL, Phillips RM. (1992):* Extubation and reintubation guided by the laryngeal mask airway in a child with Pierre Robin syndrome. *Anesthesiology*; 76: 640-41.
- Childress WF. (1981):* New method for fiberoptic endotracheal intubation of anaesthetized patient. *Anesthesiology*; 55: 595-596.
- Chirstian K, Joseph B, And Keller K. (1999):* Pressures exerted against the cervical vertebrae by the standard and intubating laryngeal mask airway: A randomized, controlled, crossover study in fresh cadavers. *Anesth Analg*; 89: 1296-1300.
- Cohen SM. (1989):* The role of the reinforced tube laryngeal mask. *Anesthesiology*; 71: A937.
- Cook TM, Nolan JP. (2002):* The PorSeal laryngeal mask airway (letter). *Anaesthesia*; 57: 288-9.

- Cook TM. (2000): A new practical classification of laryngeal view. Anaesthesia; 55: 274-9.*
- Cooper RM. (1995): The difficult airway. II. Anaesthesiol Clin North Am; 13: 3.*
- Costa E Sliva L, Brimacombe J. (1996): Tracheal tube/laryngeal mask exchange for emergence. Anesthesiology; 85: 218.*
- Cros AM, Sesay M. (1998): COPA Ann Fr Anesth Reanim; 17 (5): 112-113.*
- Crosby ET, Cooper RM, Douglas MJ. (1998): The unanticipated difficult airway with recommendations for management. Can J Anaesth; 45 (7): 757-776.*
- Crosby E.T. (1999): The Trachlight – a few more lessons (letter). Can J Anaesth; 46 (3): 297-301.*
- Culling RD Mongan P, Castro T (1989): Lightwand guided rapid sequence orotracheal intubation (abstract) Anaesthesiology; 71: A994.*
- Davey A, Moyle JTB, Wurd CS. (1992): Anesthetic equipment, 3rd edn. WB Saunders, London.*
- David J, and Thomas JG. (2000): Airway management in: Text book of anesthesia Roland Miller. Fifths edition Churchill livingstone; Ch 39: 1414-1451.*
- Davis JM, Weeks S, Crone LA. (1989): Difficult intubation in the parturient. Can J Anaesth; 36: 668.*
- Divatia JV. Anesthetic management of tracheal surgery using the laryngeal mask airway. Anaesth Intens Care; 22: 59-73.*
- Dorges V, Ocher HM Wenzel V und Schmucher P. (2000): The laryngeal tube: A New Simple airway device. Anesth Analg; 90: 1220-2.*
- Dorges V, Ocher Hm Wenzel V, Schmucher P. (2000): The laryngeal tube: A new simple airway device. Anesth Analg; 90: 1220-2.*

- Dorsch JA, Dorsch SE. (1999):* Understanding anesthesia equipment, 4th ed. William and Wilkins. Endotracheal tubes, laryngoscopes, Face Masks, and airway.
- Dravid RM, Popat MT. (1999):* The cuffed oropharyngeal airway. *Anaesthesia*; 54 (4): 402-403.
- Eduardo F, Miguel V, and Francisco M. (1999):* Laryngo-pharyngeal complaints after use of the laryngeal mask airway. *Can J Anesth*; 46: PP 220-225.
- Elaine Marier, RN. (1998):* The respiratory system in human anatomy & physiology. Fourth edition. Benjamin/cummings science publishing; Ch 23: 799-847.
- El-Gamouri, El-Baz N, Ford E. (1988):* Training residents in fiberoptic intubation in the operating room (abstract). *Anaesthesiology*; 69: a805.
- Ellis H and Feldman S (1993):* Anatomy for anaesthetist. 6th ed. Black-Well includes a chapter on airway anatomy.
- Evans NR, Llewellyn RE, Gardner SV, James MF. (2002):* Aspiration prevented by the PorSealTM laryngeal mask airway: a case report. *Can J Anesth*; 49: 413-6.
- Ferson DZ, Rahlfs TF, and Jones RL. (1997):* Evaluation of the intubating laryngeal mask as a primary airway device and a guide for blind endotracheal intubation, *Anesth*; 87: 45-52.
- Frass M. (1995):* The combitube. In: Benumof JL (ed): Airway management: Principles of practice. Mosby, St Louis; pp. 444-454.
- Fukutome T. Correct positioning of the epiglottis for application of the Brain laryngeal mask airway. Anaesthesia; 50: 818-819.*
- Fung DM, and Devitt JH. (1995):* The anatomy, physiology, and innervation of the larynx. *Anesthesiol Clin North Am*; 13: 259.
- Genzwuerker HV, Vollmer T, and Ellinger K. (2002):* Fiberoptic tracheal intubation after placement of the laryngeal tube. *British Journal of Anesthesia*; 89 (5): 733-8.

- Giggs WM, Worthley LIG, Gilligan JE. (1990): A simple percutaneous tracheostomy technique. Surg Gynecol Obstet; 170: 543-545.
- Glaisyer HR, Yule C. (2005): The protex "soft seal" single use laryngeal mask – a preliminary study in pediatric anesthesia. Paediatr Anaesth; 15 (2): 110-4.
- Gorback MS. (1987): Inflation of the endotracheal tube as an aid to blind nasal endotracheal intubation (letter). Anaesth Analg; 66: 917.
- Gorman AP. (1998): COPA. Anaesthesia; 55 (4): 414.
- Graney DO. (1986): Basic science: Anatomy. In Cummings CW, Fredrickson JM, Flacker LA, et al (eds): Otolaryngology-Head and Neck Surgery. St. Louis, CV Mosby.
- Gundoppa N. (1992): Management of pediatric difficult airway. Seminars in Anesth; 3: 220-228.
- Hagberg CA and Westhofen P. (2003): A two-person technique for fiberscope-aided tracheal extubation / reintubation in intensive care unit. J. Clin Anesth; 15 (6): 467-70.
- Hugberg CA, Vartazarin TN, Chelly JE. (1999): The incidence of gastro-esophageal reflux with the esophageal-tracheal combitube as compared to the laryngeal mask airway: An analysis using tracheal and esophageal pH electrodes. Anesthesiology; 91: (3A): A 1336.
- Hagberg CA. (2000): Current concepts in the management of the difficult airway. ASA.; 163.
- Hames KC, Pandit JJ, Marfin AG, Popat MT, Yentis SM. (2003): Use of bougie in simulated difficult intubation-I. Comparison of single-use bougie with the fiberscope. Anaesthesia; 58: 846-5.
- Harrison CA, Wise CC. (1988): Retrograde intubation (letter). Anaesthesia; 43: 609.
- Hartman, T, Kernu, CG, Zoeggler A. (1993): The esophageal-tracheal combitube in pre-hospital cardiac arrest. Ann Emerg Med; 22 (8): 1263-8.

- Hawkins M, O'Sullivan E, Charters P. (1998):* Fiberoptic intubation using the cuffed oropharyngeal airway and Aintree intubation catheter. *Anaesthesia*; 53: 891-894.
- Higgins MS, Marco AP. (1992):* An aid in oral fiberoptic intubation (letter). *Anesthesiology*; 77: 1236.
- Howarth A, Brimacombe J, Keller C. (2002):* Gum elastic bougie-guided insertion of the PorSeal laryngeal mask airway. A new technique. *Anaesthesia and Intensive Care*; 30: 624-7.
- Hsu YW, Pan MH, Huang CJ. (1998):* Comparison of the COPA and laryngeal mask airway in spontaneous breathing anesthesia. *Acta Anaesthesiol Sin*; 36 (4): 187-192.
- Hung OR, Al-Qatari M. (1997):* Light-guided retrograde intubation. *Can J Anaesth*; 44:877-882.
- Inui M. (1995):* Comparison of cardiovascular responses to airway management using a new adapter, laryngeal mask insertion, or conventional laryngoscopic intubation. *J Clin Anesth*; 7: 14-18.
- Isubeau W, and Janae L. (2000):* Basic techniques for anesthesia. In: *Pediatric anesthesia*. Second edition; 165-211.
- Ishimura H, Minami K, Sata T. (1995):* Impossible insertion of the laryngeal mask airway and oropharyngeal axes. *Anesthesiology*; 83: 867-869.
- J. Roger Malthy, Micheal T, Neil C, Waston, David J, Lipert. (2003):* LMA-Calssic™ and LMA-ProSeal™ are effective alternatives to endotracheal intubation for gynecologic laparoscopy. *Canadian Journal of Anesthesia*; 50: 71-77.
- Jaeger JM, Durbin CG Jr. (1999):* Special purpose endotracheal tubes. *Respir. Care*; 44: 661.
- John RF, Hill S, and Hughes TJ. (1991):* Airway protection by the laryngeal mask. A barrier to dye placed in the pharynx. *Anesthesia*; 46: 366-367.
- Johnston DF, Wrigly SR, Robb PJ. (1990):* The laryngeal mask airway in pediatric anesthesia. *Anesthesia*; 45: 924-927.

- Jones DH, Cohle SD. (1993):* Unanticipated difficult airway secondary to lingual tonsillar hyperplasia. *Anaesth analg*; 77: 1285.
- Joshi GP, Morsion SG, Okon Kwo NA. (1996):* Continuous hypopharyngeal PH measurements in spontaneously breathing anesthetized out patients: laryngeal mask airway versus tracheal intubation. *Anesth Analg*; 82: 254-257.
- Kattan B, And Sanyder HS. (1991):* Lingual artery hematoma resulting in upper airway obstruction. *J Emerg Med*; 9: 421.
- Keller C, Brimacombe J, Kleinsasser A, Brimacombe L. (2002):* The laryngeal mask airway ProSeal™ as a temporary ventilatory device in grossly and morbidly obese patients before laryngoscopy-guided tracheal intubation. *Anesth Analg*; 94: 737-40.
- King TA, Adams AP. (1990):* Failed tracheal intubation. *Br J Anaesth*; 65: 400-414.
- Koga K, Kaku M, Sata T. (1998):* Effective use of the COPA. *Anaesthesia*; 53 (7): 715-716.
- Kovac AL. (1993):* The Augustine guide: A new device for blind orotracheal intubation. *Anaesthesiol Rev*; 20 (1): 25-29.
- Lamb K, James MEM, and Janicki PK. (1992):* The LMA for intraocular surgery effects on intraocular pressure and stress responses. *Br J. Anesthesia*; 69: 143-7.
- Latto IP. (1985):* Management of difficult intubation. In: Latto IP, Rosen M (editors): *Difficulties in tracheal intubation*. London; 99: 141, Bailliere Tindall.
- Lim SL, Tay DHB, Thomas E. (1994):* A comparison of three types of tracheal tube for use in laryngeal mask assisted blind orotracheal intubation. *Anaesthesia*; 49: 225-227.
- Lopez Gil M, Brimacombe J, Cebrain J. (1996):* The laryngeal mask airway in pediatric practice, prospective study of skill acquisition by resident anesthesiologists. *Anesthesiology*; 84: 807-811.

- Mallampatti SR, Gatt SP. (1985):* A clinical study to predict difficult tracheal intubation. A prospective study. *Can J Anaesth*; 32: 429-434.
- Martin PD, Cyna AM, Hunter WAH. (1993):* Training nursing staff in airway management for resuscitation – a clinical comparison of the face mask and laryngeal mask. *Anesthesia*; 48: 33-37.
- Mason RA. (1992):* Learning fiberoptic intubation: Fundamental problems. *Anaesthesia*; 47: 729-731.
- Mets S, Parmet JL, Levitt JD. (1996):* Failed emergency transtracheal ventilation through a 14 gauge intravenous catheter. *J Clin Anaesth*; 8: 58-62.
- Michael Ng, Hastings RH. (1998):* Successful direct laryngoscopy assisted by posture in a patient with ankylosing spondylitis. *Anaesth Analg*; 87: 1436-1437.
- Miller C. (2000):* Management of the difficult intubation in closed malpractice claims. *American Society of Anesthesiologists NEWSLETTER*; 64 (6): 13-9.
- Miller DM, Yaukhana I, Pearce AC. (2001):* The laryngeal mask and VBM laryngeal tube compared during spontaneous ventilation. A pilot study. *Eur J Anaesthesiol*; 18: 593-8.
- Miller R. (1994):* Tracheal intubation; *American Society of Anesthesiologists*; 11: 145-149.
- Minkowitch HS. (2000):* Airway Gadgest. In: Hagberg CA (editor): *Handbook of difficult airway management*. 1st edition, Philadelphia, Churchill Livingstone; P 149-167.
- Miriam H, Brain K, Anne H. (2000):* Airway complications in infants: Comparison of laryngeal mask airway and the facemask oral-airway. *Can. J. Anesth*; 47: 315-318.
- Molhoff T, Burgand G, and Prein T. (1995):* Low flow and minimal flow anesthesia and the laryngeal mask airway. *Anesthesiology*. 1995; 83: Aug. 499.
- Morgan E, Mikhail M and Murray M. (2002):* *Clinical anesthesiology*, 3rd edition. McGraw-Hill. Airway management. Chapter (5). P. 78-80.

- Morikawa S, Safar P, DeCarlo J. (1996):* Influence of the head-jaw position upon upper airway patency. *Anesthesiology*; 22: 265-279.
- Morris IR. (1988):* Functional anatomy of the upper airway. *Emerg Med Clin North Am*; 6: 339.
- Morton T, Brady S, Clancy M. (2000):* Difficult airway management in English emergency departments *Anaesthesia*; 55: 486-8.
- Mushambi MC, Iyer GA. (2002):* Gum elastic bougies. *Anaesthesia*; 57: 727.
- Nichols KP, Zornow MH. (1989):* A potential complication of fiberoptic intubation. *Anesthesiology*; 70: 562-563.
- Nishino T. (1993):* Swallowing as a protective reflex for the upper respiratory tract. *Anesthesiology*; 79: 588-601.
- Osborn IP, Cohen J, Soper RJ, Roth LA. (2002):* Laryngeal mask airway—a novel method of airway protection during ERCP: comparison with endotracheal intubation. *Gastrointest Endosc*; 56 (1): 122-8.
- Osborn NA, Jackson APF, Smith JE. (1998):* The laryngeal mask airway as an aid to training in fiberoptic nasotracheal endoscopy. *Anaesthesia*; 53: 1080-1083.
- Ovassapian A, Wheeler M. (2000):* Flexible fiberoptic intubation. In: Hagberg CA (editor): *Handbook of difficult airway management*. 1st edition, Philadelphia, Churchill Livingstone; P 83-114.
- Ovassapian A, Meyer RM. (1998):* Airway management. In: Longnecker DE, Tinker JH, Morgan GE (editors): *Principles and practice of Anaesthesiology*. 2nd edition. St. Louis, Mosby-Year Book; 1064-1102.
- Ovassapian A, Yelich SJ, Dykes MHM, Brunner EV. (1983):* Fiberoptic nasotracheal intubation – incidence and causes of failure. *Anaesth Analg*; 62: 692-695.
- Parr MJA, Gregory M, Baskett PJF. (1998):* The intubating laryngeal mask-use in failed and difficult intubation. *Anaesthesia*; 53: 343-348.

- Patel P and Verghese C. (2000):* Delayed extubation facilitated with the use of a laryngeal mask airway (LMA) in the intensive care unit (ICU). *Anaesthesia*; 55 (4): 396.
- Paul GB, Bruace FC, and Robert KS. (1997):* Airway management in: *Clinical anesthesia* third edition. Lippincott-Raven Philadelphia New York; Ch: 23: 573-594.
- Pearce AC. (1995):* Fiberoptic intubation. In: Atkinson RS, Adams AP (editors): *Recent advances in anaesthesia and analgesia* (19th edition), London, Churchill Livingstone; 5: 71.
- Rabitsch W, krafft P, Lackner FX, Frenzer R, Hofbauer R, Sherif C and Frass M. (2004):* Evaluation of the oesophageal-tracheal double-lumen tube (Combitube) during general anesthesia. *Wien. Klin. Wochenschr* 116 (3): 90-3.
- Randell T, Yli-Hankala A, Valli H, Lindgren L. (1991):* Topical anaesthesia of the nasal mucosa for fiberoptic airway endoscopy. *Br J Anaesth*; 66: 164-167.
- Reinhart DJ, Hansen K, and Odesseus K. (1994):* The laryngeal mask airway and trace gases in the operating room. *Anesthesiology*; 81: 555.
- Roberts JT. (1999):* Fiberoptic intubation and alternative techniques for managing the difficult airway. *American Society of Anaesthesiologists (ASA)*; 136
- Rocke DA, Murray WB, Rout CC, Grows E. (1992):* Relative risk analysis of factors associated with difficult intubation in obstetric anaesthesia. *Anesthesiology*; 77: 67.
- Ronald JR. (2000):* Anatomic airway considerations in anesthesia in: *Handbook of difficult airway management*. 1st ed. Churchill Livingstone; Ch 1: 1-13.
- Rooney R, Crummy EJ, McShane AJ. (1992):* Bacteraemia following nasopharyngeal airway insertion. *Anaesthesia*; 47: 1099.
- Rose DK, Cohen MM. (1994):* The airway: problems and predictors in 18,500 patient. *Can J Anaesth*; 41: 373-383.

- Rosenblatt WH, Murphy M. (1999):* The intubating laryngeal mask: use of a new ventilatory-intubating device in the emergency department. *Ann Emerg Med*; 33: 234-238.
- Rosenblatt WH. (2004):* The use of the LMA-ProSeal in airway resuscitation. *Anesth Analg*; 99 (1): 310-1.
- Russel SH, Hirsch NP. (1993):* Simultaneous use of two laryngoscopes. *Anaesthesia*; 18: 918.
- Samsooni GL and Young RJ. (1987):* Difficult tracheal intubation-a retrospective study. *Anaesthesia*; 42: 487.
- Samsoon GLT, Young JRB. (1987):* Difficult tracheal intubation a retrospective study. *Anaesthesia*; 42; 487.
- Santha TR. (1992):* Retrograde intubation using the subcricoid region. *Br J Anaesth*; 68: 109-112.
- Sarpellari M, Marson F, Nani R. (1998):* Translaryngeal tracheostomy (TLT). A variant technique for use in hypoxaemic conditions and in the difficult airway. *Minerva Anaesthesiol*; 64 (9): 333-337.
- Savva D. (1994):* Prediction of difficult tracheal intubation. *Br J Anaesth*; 73: 194.
- Sanchez AF, Morrison DE. (2000):* Retrograde intubation. In: Hagberg CA (editor): *Handbook of difficult airway management*. 1st edition, Philadelphia, Churchill Livingstone; P 115-148.
- Schumacher P, Stotz G, Schneider M, Urwyler A. (1992):* Laryngospasm during transtracheal high frequency jet ventilation. *Anaesthesia*; 47: 855-856.
- Schwab RJ, Gupta KB, Gefer WB. (1995):* Upper airway and soft tissue anatomy. *Am J Respir Crit Care Med*; 152: 1673.
- Sher M, Brimacombe J, and Laing D. (1995):* Anaesthesia for laser pharyngoplasty – a comparison of the tracheal tube versus reinforced laryngeal mask airway. *Anesth Intens Care*; 23:: 149-154.
- Sivarajan M, Stoler E, Kil HK, Bishop JM. (1995):* Jet ventilation using fiberoptic bronchoscopes. *Anesth Analg*; 80: 384-7.

- Smith BL. (1986):* Laryngeal cyst following, fiberoptic intubation. *Anaesthesia*; 41: 430-431.
- Smith I, and White PF. (1992):* The use of LMA as an alternative of a face mask during outpatient arthroscopy. *Anesthesiology*; 77: 850-855.
- Southerland AD, Sale JP. (1986):* Fiberoptic awake intubation – A method of topical anaesthesia and orotracheal intubation. *Can Anaesth Soc J*; 33: 502-504.
- Splinter WM, Smallman B, Rhine EJ. (1994):* Post-operative sore throat in children and the LMA. *Can. J. Anesth*; 41: 1081-83.
- Stauffer JL. (1999):* Complications of endotracheal intubation and tracheostomy. *Respir Care*; 44:828.
- Steven JE, and Burden G. (1994):* Reinforced LMA and magnetic resonance imaging. *Anesthesia*; 49: 79-80.
- Stoppacher R, Teggatz JR, Jentzen JM. (2004):* Esophageal and pharyngeal injuries associated with the use of the esophageal-tracheal combitube. *J Forensic Sci*; 49 (3): 586-91.
- Surendra Kumar D, Jones G. (2001):* Is your bougie helping or hindering you? *Anaesthesia*; 56: 1121.
- Sykes JM, and Isaacs RS. (1996):* Applied anatomic considerations in airway management. In Hanowell LH, Waldron RJ (eds): *Airway Management*. Philadelphia, Lippincott-Reven; P3.
- Taguchi M. Watanabe S, Asakura N. (1994):* End-tidal sevoflurane concentrations for laryngeal mask airway insertion and for tracheal intubation in children. *Anesht*; 81: 628-631.
- Takahatu O, Kubota M, Muniya K. (1997):* The efficacy of the "BURP" Maneuver during a difficult laryngoscopy. *Anaesth Analg*; 84: 419-421.
- Talke PO, Nguyen H. (1999):* Concept of easy fiberoptic intubation via a laryngeal mask airway (letter). *Anaesth Analg*; 88: 227-234.
- Tashayod M. (1967):* A new double-curve endotracheal tube for nasal intubation. *Brit J Anaesth*; 39: 823.

- Tintinalli JE, Claffey J. (1981):* Complications of nasotracheal intubation. *Ann Emerg Med*; 10: 142.
- Trevisanuto D, Grazzina N, Ferrarese P, Micaglio M, Verghese C, Zana V. (2005):* Laryngeal mask airway used as a delivery conduit for the administration of surfactant to preterm infants with respiratory distress syndrome. *Biol Neonate* 13; 87 (4): 217-220.
- Uchida T, Hikawa Y, Saito Y, Yasuda K. (1997):* The McCoy levering laryngoscope in patients with limited neck extension. *Can J Anaesth*; 44: 674-676.
- Vaughan RS. (1991):* Training in fiberoptic laryngoscopy. *Br J Anesth*; 66: 538-540.
- Wong MEK, Bradirek JP. (2000):* Surgical approaches to airway management for anaesthesia practitioners. In: Haberg CA (editor): *Handbook of difficult airway management*. 1st edition, Philadelphia, Churchill Livingstone; P 185-218.
- Yentis SM, Lee DJH. (1998):* Evaluation of an improved scoring system for the grading of direct laryngoscopy. *Anaesthesia*; 53: 1041-1044.
- Yoshitaka F, Yuhji S, Hiroyoshi T, and Hidenor T. (1998):* Cardiovascular response to tracheal extubation or removal of LMA in children. *Can. J. Anesthesic*; 45: 178-181.
- Zorror MH, Mitcheall MM. (1986):* Foreign body aspiration during fiberoptic assisted intubation. *Anaesthesiology*; 64: 303.