

REFERENCES

References

- Abernethy DR, Greenblatt DJ (1982):** Pharmacokinetics of drugs in obesity. Clin Pharmacokinet 7:108-24.
- Adams J.P. and Murphy P.G. (2000):** Obesity in anaesthesia and intensive care, 85:91-108.
- Agarwal N, Shibutani KJ, Sanfilippo JA, Del Guercio LRM (1982):** Hemodynamic and respiratory changes in surgery of the morbidly obese. Surgery, 92:226-34.
- Alauddin A, Meterissian S, Lisbona R, MacLean LD, Forse RA (1990) :** Assessment of Cardiac function in patients who were morbidly obese. Surgery, 108:809-20.
- Albrink MJ (1971):** Obesity. Cecil-Loeb textbook of medicine. Ed. by PB Beeson and W McDermott, 13th ed, p. 1452, Saunders, London.
- Alexander JK (1963):** Obesity and the circulation. Modern Concepts Cardiovasc Dis, 32:799-803.
- Alexander JK (1964):** Obesity and cardiac performance. Am J Cardiol, 14:860-5.
- Alexander JK, Dennis EW, Smith WG et al. (1962):** Blood volume, cardiac output, and distribution of systemic blood

flow in extreme obesity. Cardiovasc Res Center bull, 1:3944.

Alexander JK, Guthrie AE, Sakaguchi H, Crawford HW and Cole VW (1959): Lung volume changes with extreme obesity. Clinical Research, 7:171.

Alexander JK, Peterson KL (1972): Cardiovascular effects of weight reduction. Circulation, 45:310-8.

Alpert MA, Hashimi MW (1993): Obesity and the heart. Am J Med Sci, 306:117-23.

Alpert MA, Lamnbert CR, Terry BE et al. (1995): Influence of left ventricular mass on left ventricular diastolic filling in normotensive morbid obesity. Am Heart J, 130:1068-73.

Alpert MA, Singh A, Terry BE (1989): Effect of exercise on left ventricular function and reserve in morbid obesity. Am J Cardiol, 63:1478-82.

Alpert MA, Terry BE, Lambert CR, Kelly DL, Panayiotou H, Hukerji V, et al. (1993): Factors influencing left ventricular systolic function in nonhypertensive morbidly obese patients, and effect of weight loss induced by gastroplasty. Am J Cardiol, 71:733-737.

Amad KH, Brennan JC, Alexander JK (1965): The cardiac pathology of chronic exogenous obesity. *Circulation*, 32:740-5.

Anderson J, Rasmussen JP, Eriksen J (1977): Pulmonary function in obese patients scheduled for jejunio-ileostomy. *Acts Anaesthesiol Scand*, 21:346-51.

Anderson J (1972): Obesity. *British Medical Journal*, 1:560.

Anthony P. adams and Jereney N. cashnar(2000):Recent advances in anaesthesia and analgesia, chapter II, vol 21

Antoni jevic I, Mousa SA, Schafer M and Stein C (1995): Perineurial defect and peripheral opioid analgesia in inflammation. *J Neuro sci*, 15:165.

Aitkenhead A. R. and G.Smith (1990): textbook of anaesthesia, second edition. Churchill Livingstone (1990): p.337

Backman L, Freyschuss V, Hallberg D, Meicher A (1973): Cardiovascular function in extreme obesity. *Acta Med Scand*, 193:437-42.

Barash P. G., Cullen B. F., Steolting R. K., eds. (2001): Clinical anaesthesia, 4th ed. Philadelphia:Lippincott Williams&Wilkins, 2001:1035-1041.

Bardoczky GI, Yernault JC, et al. (1995): Large tidal volume ventilation does not improve oxygenation in morbidly obese patients during anesthesia. *Anesthesia and Analgesia*, 81(2):385-383.

Beebe DS, Ostanniny I, Komanduri, Ct al. (1998): Continual noninvasive blood pressur monitoring with the VasotractTM: Experience in the morbidly obese *Anesthesiology*, 89(3A):A565.

Belani KG and Buckley JJ (1995): The VasotractTM, a revolutionary instrument to monitor blood pressure. non-invasively in a continual fashion. *Anesthesiology*, 83:A489.

Bendixen H (1978): Morbid obesity. *Anesthesiology*, 6(10):1-15.

Bentley JB, Finley JM, Humphrey ER, Gandolfi AJ, and Brown Br (1983): Obesity and alfentanil pharmacokinetics. *Anesth Analg*, 62:S251.

Bentley JB. Vaughan RW, Cork RC (1980): Hepatorenal indices among general anaesthetics in obesity. *Anaesthesiology*, 53:S259.

Bentley JB, Vaughan RW, Gandolfi A, et al. (1982): Halothane biotransformation in obese and nonobese patients. *Anesthesiology*. 57:94.

Bentley JB, Vaughan RW, Miller MS, Calkins JM, Gandolfi AJ (1979): Serum inorganic fluoride levels in obese patients during and after enflurane anesthesia. *Anesth Analg*, 58:409-12.

Bentley JB, Borel JD, Gillespie TS, Vaughan RW, and Gandolfi AJ (1981): Fentanyl pharmacokinetics in obese and non-obese patients. *Anesthesiology*, 55:A177.

Bentley Robert W, Vaughan Randall, C Cork and Jay Gandolfi (1981): Does evidence of reductive halothane biotransformation correlate with hepatic binding of metabolites in obese patients? *Anaesthesia Analgesia*, 60:548.

Benumof JL (1991): Management of difficult airway, with special emphasis on awake tracheal intubation. *Anesthesiology*, 75:1087-1110.

Bharati S, Lev M (1995): Cardiac conduction system involvement in sudden death of obese young people. *Am Heart J*, 129:273-81.

Biring MS, Lewis MI, Liu JI, Mosenifar Z (1999): Pulmonary physiologic changes of morbid obesity. *Am J Med Sci*, 318:293-7.

Bjerkedal T (1957): Overweight and hypertension. *Acta Med Scand*, 159:13-26.

Blass NH (1981): Morbid obesity and other nutritional disorders. In: Katz J et al. (eds) *Anesthesia and Uncommon Diseases*. WB Saunders, Philadelphia.

Blouin RA, Kolpek JH, Mann HJ (1987): Influence of obesity on drug disposition. *Clin Pharm*, 6:706-14.

Boulanger BR, Milzman D, Mitchell K, Rodriguez A. (1992): Body habitus as a predictor of injury pattern after blunt trauma. *J Trauma Inj Inf Crit Care* , 33: 228—32

Bray GA (1987): Obesity and the heart. *Mod Concepts Cardiovasc Dis*, 56:67.

Bray GA and Gallagher TF (1972): Obesity: A serious symptoms. *Annual Internal Medicine*, 77:795.

Brodsky JB, Shulman MS, Kim SK, et al. (1986): epidural morphine following abdominoplasty. *Plast. Reconstr. Surg.*, 78:125.

Brown BR (eds) (1982): Anesthesia and the obese patient. In: Contemporary Anesthesia Practice, Vol. 5, FA Davis, Philadelphia.

Buckley FP (1990): Anesthetizing the morbidly obese patient. ASA Refresher Courses in Anesthesiology, 18:53-68.

Buckley FP (1992): Anesthesia and obesity and gastrointestinal disorders. In: Barash PG, Cullen BF, Stoelting RK, eds. Clinical Anesthesia. Philadelphia: J.b. Lipincott, 1169-83.

Buckley FP (1994): Anaesthesia for the morbidly obese patient. Can J Anaesth, 41:R94-R100.

Buckley FP, Robinson NB, Simonowitz DA, Dellinger EP (1983): Anaesthesia in the morbidly obese. A comparison of anaesthetic and analgesic regimens for upper abdominal surgery. Anaesthesia, 38:840-51.

Burwell CB, Robin ED, Whaley RD, et al. (1956): extreme obesity associated with alveolar hypoventilation: a pickwickian syndrome. Am J Med; 21:811-818

Catenacci AJ, Anderson JD, and Boersma L (1961): Anesthetic hazards of obesity. JAMA, 175:657.

Cheymol G (1993): Clinical pharmacokinetics of drugs in obesity : an update. Clin Pharmacokinet, 25:103-14.

Clark Venable, M.D. and Paul H. Ting, M.D. (2003): Obesity: Anesthetic Considerations, GAS Net. Jun 7th, 2003.

Cork RC, Vaughan RW, and Bentley JB (1981) : General anesthesia for morbidly obese patients - an examination of postoperative outcomes. Anesthesiology, 54:310-3.

Coussa M, Proietti S, Schnyder P, Frascarolo P, Suter M, Spahn DR, Magnusson L.(2004): Prevention of atelectasis formation during the induction of general anesthesia in morbidly obese patients. Anesth Analg. 2004 May; 98(5):1491-5, table of contents.

Damia G, Mascheroni D, Croci M, Tarenzi L (1988): Perioperative changes in functional residual capacity in morbidly obese patients. Br J Anaesth, 60:574-8.

Daniels SR, Morrison JA, Sprecher DL, Khoury P, Kimball TR (1999): Association of body fat distribution and cardiovascular risk factors in children and adolescents. Circulation, 99:641-5.

Duflou J, Virmani R, Rabin I, Burke A, Farb A, Smialek J (1995): Sudden death as a result of heart disease in morbid obesity. *Am Heart J*, 130:306-13.

Dumont L, Mattys M, Mardirosoff C, Vervloesem N, Alle JL, Massaut J (1997): Changes on pulmonary mechanics during laparoscopic gastroplasty in morbidly obese patients. *Acta Anaesthesiol Scand*, 41:408-13.

Fernandez Mere LA, Alvarez Blanco M. (2004): Obesity and bariatric surgery: anesthesia implications, *Nutr Hosp.* 2004 Jan-Feb; 19(1):34-44

Fisher A (1980): Obesity and its relation to anesthesia. *Anesthesiology*, 45:6330-6334.

Fisher A Waterhouse TD and Adams AP (1975): obesity: its relation to anaesthesia. *Anaesthesia*, 30:633.

Fleming PR (1975): Preoperative assessment of cardiovascular status. *General anaesthesia textbook*, Ed. by Cicel and JF Nunn, 3rd edition, p. 59, Part 11, Butterworths, London.

Frink EJ, Malan TP, Brown EA, Morgan S, Brown BR (1993): Plasma inorganic fluoride levels with sevoflurane anesthesia in morbidly obese and nonobese patients, *Anesth Analg*, 76:133-7.

Gallus AS, Hirsh J, O'Brien SE (1976): Prevention of deep venous thrombosis with small, subcutaneous doses of heparin. JAMA 235:1980.

Halaka K, Mustajoki P, Aittomaki J, Sovijarvi AR (1995): Effect of weight loss and body position on pulmonary functions and gas exchange abnormalities in morbid obesity. Int J Obes relat metab Disord, 19:343-6,686.

Hedenstierna G (1990): Gas exchange during anaesthesia. Br J Anaesth, 64:507-14.

Higuchi H, Satoh T, Arimura S, et al. (1993): Serum inorganic fluoride levels in mildly obese patients during and after sevoflurane anesthesia. Anesth Analg, 77(5): 1018.

Hills NH, Flug JJ, Jeyasingh K, Boardman L and Cobnan JS (1972): Prevention of deep vein thrombosis by intermittent pneumatic compression of calf. British Medical Journal, 1:131.

Hood DD, Dewan DM (1993): Anesthetic and obstetric outcome in morbidly obese parturients. Anesthesiology, 79:1210-18.

Hodgkinson R, and Husain FJ (1980): Obesity and the cephalad spread of analgesia following epidural administration of bupivacaine for cesarean section. *Anesth Analg.*, 59:89-92.

Hunter JD, Reid C, Noble D (1998): Anaesthetic management of the morbidly obese patient. *Hosp Med*, 59:481-3.

Javier Nieto F (1992): Childhood weight and growth rate as predictors of adult mortality. *Am J Epidemiol.*, 136:201-13.

Jeevanandam M, Rnmias L, Schilier WR (1991): Obesity and the metabolic response to severe multiple trauma in man. *J Clin invest*, 87:262-9.

Joris J, sottiaux T, Chicke JD, Luten I, and Lainy M (1994) : bi-level CPAP (BIPAP) reduces the postoperative restrictive pulmonary syndrome in obese patients after gastropasty. *Br J Anesth*, 72 (Suppl. 1):A111.

Jupiter JB. Ring D, Rosen H (1995): The complications and difficulties of management of nonunion in the severely obese. *J Orthopedics Trauma*, 9:363-70.

Juvun P, Marmuse JP, Delerme S, et al. (1999): Post-operative course after conventional or laparoscopic

gastroplasty in morbidly obese patients. Eur J Anaesthesiol., 16:400-3.

Kakkar VV, Howe CT, Nicolaides AN and Clarke MR (1970): Deep vein thrombosis of the leg: Is there a “high risk” group? American Journal of Surgery, 120:521.

Kaitman AJ, goldring RM (1976): Role of circulatory congestion in the cardio-respiratory failure of obesity. Am J Med., 60:645-55.

Kirby I and Howard EC (1987): Propofol in a morbidly obese patient. Anaesthesia, 42:1125-1126.

Kirkegaard-Nielsen H, Hebbo-hansen HS, et al. (1996): Anthropometric variables as predictors for duration of action of atracurium-induced neuromuscular blockade. Anesth Anaig, 83(3):1076-1080.

Kirkegaard-Nielsen H, LindhoLm P, et al (1998): Antagonism of atracurium-induced block in obese patients. Candian Journal of Anaesthesia, 45(1):39-41.

Ladergaard-Pederson MJ (1981): Recovery from general anesthesia in obese patients. Anesthesiology, 55:720.

Lean ME (1999): Obesity and cardiovascular disease: the wasted years. Br J Cardiol, 6:269-73.

Ledgerwood AM, Harrigan C, Saxe JM, and Lucas CE (1992): The influence of an anesthetic regimen on patient care, outcome, and hospital charges. *Am Surg*, 58:527-533.

Lee J, Larsen K, Buckley J, and Roberts R (1981): Pulmonary function and its correlation to the degree of obesity in 294 patients. *Anesthesiology*, 8(2):28-32.

Leith P, Sanborn K, and Brock-Utne JG (1997): Intraoperative epidural catheter malfunction in two obese patients. *Acts Anaesthesiol Scand*, 41:651-653.

Lesser GT, Deusch S and Matkowsky J (1971): Use of independent measurement of body fat to evaluate overweight and underweight. *Metabolism clinical and Experimental*, 20:792.

Lessing AJP and Dekock MJ (1968): Anaesthesia in a grossly obese patient. *British Journal of Anaesthesia*, 40:139.

Loos RJJ, Bouchard C (2003): Obesity —is it a genetic disorder? (Review). *J Intern Med* 2003; 254: 401-425.

Luce J (1980): Respiratory complications of obesity. *Chest*, 78:626-31.

MacGregor MI, Block AJ and Ball WC Jr (1970): Serious complications and sudden death in the pickwickian syndrome. Hopkins Medical Journal, 126:279.

Madsen P, Iversen H, and Secher Nil (1993): Central venous oxygen saturation during hypovolemic shock in humans. Scand J Clin Lab Invest, 53:67-72.

Magnusson L. and U. R. Spahn (2003): New concepts of atelectasis during general anaesthesia British Journal of Anaesthesia 91 (1): 61—72.

Maitra AM, Palmer SK, Bachhuber SR, et al (1979): Continuous epidural analgesia for cesarean section in a patient with morbid obesity. Anesth Analg., 58:348.

Mallampati SR, Gatt SP, Guigino LD, et al: A clinical sign to predict difficult tracheal intubation: A prospective study. Can Anaesth Soc J 32:429, 1985.

Mann GV (1974): The influence of obesity on health. N Engl J Med, 291 :(1)178, (11)226.

Magali Poulain, Mariève Doucet, Genevieve C. Major, Vicky Drapeau, Frédéric Sériès, Louis-Philippe Boulet, Angelo Tremblay, François Maltais(2006):The effect of obesity

on chronic respiratory diseases pathophysiology and
therapeutic strategies CMAJ APRIL 25, 2006 ;174(9) :1293

Manson JE, Colditz GA, Stampfer MJ, et al (1990): A prospective study of obesity and risk of coronary heart disease in women. New Engl J Med., 322:882-9.

Marik P, Varon J (1998): The obese patient in the ICU. Chest, 113:492-8.

Mayersohn M, Calkins JM, Perrier DG, et al. (1981): Thiopental kinetics in obese patients. Anesthesiology, 55:178A.

McCarrol SM, Bras P, and Edmiston LK (1990): A prospective study to identify the features associated with difficulty in laryngoscopy and intubation in the obese patient. Anesth Analg, 65:S106.

McCracken BH (1962): Etiological aspects of obesity. American Journal of Medical Sciences, 243:99.

McCulloch WJD, and Littlewood DG (1996): Influence of obesity on spinal analgesia with isobaric 0.5% bupivacaine. Br J Anaesth, 58:610.

Mckenzie JC (1967): Social and economic implications of minority food habits. Proceedings of the Nutrition Society, 26: 197.

Mikhail N, Golub MS Tuck ML (1999): Obesity and hypertension. Prog Cardiovasc Dis, 42:39-58.

Miller,eds. (2000): Anaesthesia , 6th edi. Churchill Livingstone, a Harcourt health sciences company(2000): sec 3, part A, chapter26.

Miller,eds. (2000): Anaesthesia , 6th edi. Churchill Livingstone, a Harcourt health sciences company(2000): sec 3, part C, chapter39.

Milligan KR, Cramp P, and Schatz L (1991): The effect of positioning and obesity on epidural analgesia spread. Anesth Analg, 72:S185.

Miliman RP, Meyer TJ, Eveloff SE (1992) : Sleep apnea in the morbidly obese. Rhode Island Med, 75:483-6.

Monte and Moya (1974): Evaluation of respiratory system. In: Introduction to the practice of anaesthesia. p. 261, London.

Munsch Y, and Sagnard p (1997): The anaesthetist's point of view in the surgical treatment of morbid obesity. Ann Chir, 51(20):183-188.

Murphy PG (2000): In: Hemmings HC Jr, Hopkins PM, eds. Foundations of Anaesthesia. Basic and Clinical Sciences. London: Mosby, 703-11.

Neuman GG, Baldwin CC, Pertini AJ, et al (1986): Perioperative management of a 430 kilogram (946 pound) patient with pickwickian syndrome. *Anesth Analg.*, 65:985.

Ninh T. Nguyen and Bruce M. Wolfe (2005): The Physiologic Effects of Pneumoperitoneum in the Morbidly Obese. *Ann Surg.* 2005 February; 241(2): 219–226.

Oberg B, and Poulsen TD (1996): Obesity: an anaesthetic challenge. *Acta Anaesthesiologica Scandinavica*, 40:191-200.

Pandazi A, Bourlioti A, Kostopanagiotou G.(2005): Bispectral Index (BIS) monitoring in morbidly obese patients undergoing gastric bypass surgery: experience in 23 patients. *Obes Surg.* 2005 Jan; 15(1):58-62.

Parker RK, Holtmann B, and White PF (1992): Patient controlled analgesia: Does a concurrent opioid infusion improve pain management after surgery? *JAMA*, 266:1947-1992.

Partinen M, Guilleminault C. (1990): Daytime sleepiness and vascular morbidity at seven-year follow-up in obstructive sleep apnea patients. *Chest*, 97:27-32.

Paul DR, Hoyt JL, and Boutros AR (1976): Cardiovascular and respiratory changes in response to change of posture in the very obese *Anesthesiology*, 45:73-8.

Pelosi P, Croci M, Ravgnan I, et al. (1998): The effect of body mass on lung volumes, respiratory mechanics, and gas exchange during general anesthesia. *Anesth Analg*, 87:645-60.

Pelosi P, Ravagnan I, Giurati G, Panigada M, buttoni N, Tredici S (1999): Positive end-expiratory pressure improves respiratory functions in obese but not in normal subjects during anesthesia and paralysis. *Anesthesiology*, 91:1221-31.

Pitkanen MT (1987): Body mass and spread of spinal anesthesia with bupivacaine. *Anesth Analg*, 66:127-131.

Pitkin RM (1976): Abdominal hysterectomy in obese women. *Surg Gynecol Obstet*, 142:532-536.

Pories WJ, Van Riz AM, and Johnson WD (1972): Complications following surgery for duodenal ulcer in obese patients. *Arch Surg.*, 105:438-440.

Postlethwait RW and Johnson WD (1972): Complications following surgery for duodenal ulcer in obese patients. *Archives of Surgery*, 105:438.

Prem KA, Mesheha NM, and McKeveley JL (1965): Operative treatment of adenocarcinoma of the endometrium in obese women. *Am J Obstet Gynecol.*, 92:16-22.

Randall C, Cork Robert, W Vaughan John, B Bentley (1981): General anaesthesia for morbidly obese patients. An examination of postoperative outcomes. *Anaesthesiology*, 54:310.

Rao CS (1996): Is it not dangerous to push epidural needles all the way to the hub? *Anaesthesia*, 51:513.

Rasmussen JP, Eriksen J, and Andersen J (1977): Evaluation of impedance cardiography during anaesthesia in extremely obese patients. *Acta Anaesthesiol Scand* 21:342-345.

Raucoles-Aime M, Brimaud D (1996): Diabetes mellitus: implications for the anesthesiologist. *Anesth*, 9:247-51.

Ray C, Sue D, Bray G, Hanien JE., Wasserman K (1983): Effects of obesity on respiratory functions. *Am Rev Respir Dis*, 128:501-6.

Rayfield EJ, Ault MJ, Keusch GT, Brothers MJ, Nechermias C, and Smith H (1982): Infection and diabetes: The case for glucose control. *Am J Med*, 72:439-450.

Robertson IK, and Etringham RJ (1985): Anaesthetic management of the morbidly obese. *Br J Hosp Med*, 42:224-228.

Rochester DF and Enson Y (1974): Current concepts in pathogenesis of the obesity hypoventilation syndrome. Mechanical and circulatory factors. *American Journal of Medicine*, 57:402.

Salem MR, Dalal FY, Zygmunt MP, et al. (1978): Does PEEP improve intraoperative arterial oxygenation in grossly obese patients? *Anesthesiology*, 48:280.

Samuelson PN, Merin RG, Tabes DR, et al (1976): Toxicity following methoxyflurane anesthesia: The role of obesity and effect of low dose anesthesia on fluoride metabolism and renal functions. *Can Anaesth Soc J*, 23:465-479.

Schwartz AE, Matteo RS, Ornstein E, Young WL, and Myers KJ (1991): Pharmacokinetics of sufentanil in obese patients. *Anesth Analg.*, 17:233-236.

Servin F, Farinotti R, Haberer JP, and Desmonts JM (1993):

Propofol infusion for the maintenance of anesthesia in morbidly obese patients receiving nitrous oxide. *Anesthesiology*, 78:657-665.

Shepard JW, et al. (1985): Relationship of ventricular ectopy to

oxyhemoglobin desaturation in patients with obstructive sleep apnea. *Chest*, 88:335.

Shields J (1972): Monozygomatic *twins* brought up apart and

brought together. Oxford University Press, London.

Silver S and Bauer J et al (1931): Obesity constitutional or

endocrine. *American Journal of Medical Sciences*, 18 1:769.

Silverstone JT and Solomon T (1965): Psychiatric and somatic

factors in the treatment of obesity. *Journal of Psychosomatic Research*, 9:249.

Sloan AW (1967): Estimation of body fat in young men. *Journal of*

Applied Physiology, 23:311.

Smith-Choban P Weireter U, Maynes C (1991): Obesity and

increased mortality in blunt trauma. *J Trauma*, 31:1235-7.

Snider RD, Kruse JA, Bander JJ, Dunn GB (1995): Accuracy of estimated creatinine clearance in obese patients with stable renal function in the intensive care unit. *Pharmacotherapy*, 15:747-53.

Soderberg M, Thomson D, and White T (1977): Respiration, circulation and anaesthetic management in obesity. Investigation before and after jejuno-ileal bypass. *Acta Anaesthesiol Scand*, 21:55-61.

Santesson J (1977): Oxygen transport and venous admixture in the extremely obese. Influence of anaesthesia and artificial ventilation with and without positive endexpiratory pressure. *Acta Anaesthesiol Scand*, 21:56-61.

Sprung J, Whalley DG, Falcone T, Wilks W, Navratil JE, Bourke DL.(2003): The effects of tidal volume and respiratory rate on oxygenation and respiratory mechanics during laparoscopy in morbidly obese patients. *Anesth Analg*. 2003 Jul; 97(1):268-74

Stoelting RK and Dierdrof SF(2002): Anaesthesia and co-existing disease, 4th ed.NewYork :Churchill Livingstone, 2002: page 446

Strube PJ, Hulands GB, Halsey MJ (1987): Serum fluoride levels in morbidly obese patients: Enflurane compared with isoflurane anaesthesia. *Anaesthesia*, 42:685-9.

Taviainen T, Tuominen M, and Rosenberg PM (1990): Influence of obesity on the spread of spinal analgesia after injection of plain 0.5% bupivacaine at the L3-4 or L4-5 interspace. *Br J anaesth*, 64:542-546.

Tsueda K, Debrand M, Zeok SS, et al (1979): Obesity supine death syndrome: reports of two morbidly obese patients. *Anesth Analg*, 58:345.

Tusman G, Bohm SH, Vazquez de Anda GF, do Campo JL, Lachmann B. (1999): ‘Alveolar recruitment strategy’ improves arterial oxygenation during general anaesthesia. *Br J Anaesth* 82: 8—13

Valensi P, Thi BN, Lormeau B, Panes J, Attali JR (1995): Cardiac autonomic function in obese patients. *Int J Obes Relat Metab Disord*, 19:113-18.

Valentine SJ, Marjot H, and Monk CR (1990): Preoxygenation in the elderly: A comparison of the four maximal breath and three-minute techniques. *Anesth Analg.*, 71:516-519.

Vaughan RW (1982): anesthetic management of the morbidly obese patient. In: Brown BR, Blitt CD, Giesecke AH, Vaughan RW, eds. *Anesthesia and the Obese Patient*. New York: Davis, 71-94.

Vaughan RW (1985): Anesthesia for the massively obese patient. ASA annual Refresher Course Lecture, No. 126. American Society of Anesthesiologists.

Vaughan RW and Roa NL (1973): A case of morbid obesity. *Anaesthesiology*, 39:107.

Vaughan RW and Wise L (1975): Choice of abdominal operative incision in the obese patient: A study usual blood gas measurement. *Annals of Surgery*, 181:829.

Vaughan RW and Wise L (1975): Postoperative arterial blood gas measurement in obese patients: effect of position on gas exchange. *Annals of Surgery*, 182:705.

Vaughan RW and Wise L (1976): Intraoperative arterial oxygenation in obese patients. *Ann Surg*, 184:35-42.

Vaughan RW, and Conahan TJ (1980): Cardiopulmonary consequences of morbid obesity. *Life Sci*, 26:2119.

Vaughan RW, Bauer S and Wise L (1975): Volume and PH of gastric juice in obese patients. *Anaesthesiology*, 43:686.

Vaughan RW, Bauer 8, and Wise L (1976): Effect of position (semi-recumbent versus supine) on postoperative oxygenation in markedly obese subjects. *Anaesth Analg*, 55:37-41.

Vaughan RW, Cork RC, and Hollander D (1981): The effect of massive weight loss on arterial oxygenation and pulmonary function tests. *Anesthesiology*, 54:325.

Vgontzas AN, Tan TL, Bixler EO, et al (1994): Sleep apnea and sleep disruption in obese patients. *Arch Int Med*, 154:37-41.

Vickers MD (1978): Obesity. In: *Medicine for Anaesthetists* Second edition, p. 389, Oxford, London.

Von UngernSternberg B. S., A. Regli, M. C. Schneider, F. Kunz and A. Reber (2004): Effect of obesity and site of surgery on perioperative lung volumes *British Journal of Anaesthesia* 92 (2): 202—7 (2004)

Wallace DH, Currie JM, Gilstrap IC, and Santos R (1992):
Indirect sonographic guidance for epidural anesthesia in
obese pregnant patients. Reg Anesth, 17:233-236.

Walton B, Simpson BR, Stranin L, et al (1976): Unexplained
hepatitis following halothane. BMJ, 1:1171.

Warlow CP, McNeil A, Ogston D and Douglas AS (1972):
Platelet adhesiveness, coagulation and fibrinolytic activity
in obesity. Journal of Clinical Pathology, 25:484.

Warner WA and Gantt LP (1968): The obese patient and
anaesthesia. JAMA, 205:102.

Wasan KM, Lopex-Berestein G (1993): The influence of serum
lipoproteins on the pharmacokinetics and
pharmacodynamics of lipophilic drugs and drug carriers.
Arch Med Res., 24:395-401.

Weinstein JA, Matteo RS, Ornstein E, et al. (1988):
Pharmacodynamics of vecuronium and atracurium in the
obese surgical patient. Anesth Analg., 67:1149.

Wylie and Churchill Davidson HC (1984): Difficult intubation.
In: A Practice of Anaesthesia, 5th edition, pp. 1137- 1139.

Xavier PI, and Sunger F (1993): Medical hazards of obesity. Ann.
Int Med, 119:655-660.