

References

REFERENCES

- **Abboud TK, Chulkin K, Nouihed R, Srlos F. (1983):** Epidural bupivacaine, chloroprocaine or lidocaine for caesarean section maternal and neonatal effects. *Anesth Analg*, 62:914.
- **Abboud, TK, Kin SH and Henriksen E. (1985):** Comparative maternal and neonatal effects of halothane and Euflurane for caesarean section. *Acta Anesthesiol Scand*, 29:663.
- Abouleish E, Abboud T, Lechevalier T, Alford K. (1994):** Rocuronium - for caesarean section. *Br J Anesth*; 73:336-341.
- **Aitkenhead, AR and smith, G. (1990):** Neuromuscular blockade. Textbook of anesthesia 2nd Ed. Churchill Livingstone, London:211.
- **Alahuhta S, Rasanen J, Jouppila P and Hollmen Al (1991):** Effect of extradural bupivacaine with adrenaline for caesarian section on uteroplacental and fetal circulation. *Br J Anesthesiology* ; 45(3):102-6.
- **Alahuhta S, Rasanen J, Jouppila p and Holmmen Al. (1993):** Uteroplacental and fetal circulation during extradural bupivacaine adrenaline and bupivacaine for caesarea section in hypertensive pregnancies with chronic fetal asphyxia. *Br J Anesth* ; 71:348.
- **Alapydin T, Karamanalis B and Pamukeun Z (1999):** Comparison of the maternal and neonatal effect of sevoflurane, isoflurane and halothane for elective caesarean section. *Br J Anesth*, 82:159.

-
- **Alon , Ball RH, Gillie MH and parer J.T. (1993):** Effects of propofol and thiopentone on maternal and fetal cardiovascular and acid-base variable in the pregnan ewe. *Anesthesiology*, 78:562.
 - **Anderson R,A., Baille TA, Axelson M, et al (1996):** stable isotope studies on steroid metabolism and kinetics in human pregnancy, *sterouds* 55: 443.
 - **Apgar, V., Burns, J.J., Brodie, B.B. and paper, E. M. (1952):** The transmission of meperidine across the human placenta. *Am J. Obstet, Gynecol.*, 64,1368.
 - **Aps C, Bell JA, lenkins BS and Reynolds F. (1976):** Logical approach to lignocaine therapy. *Br- Med*; 1:3.
 - **Aun CST (1999):** New IV agents. *Br J Anesthesiology*, 83:29-41.
 - **Baden JM and Brodsky JB. (1984):** Inhibition of fetal methionine synthetase by nitrous oxide. *Br J Anesth*; 56 :523.
 - **Bannardat JP, Maille JC and Colau JC.(1982):** Maternal and fetal concentration of morphine after intrathecal administratio during labor. *Br J Anesth*; 56:1351.
 - **Baraka A, Louis F and Dalleh R. (1990):** Maternal awareness and neonatal outcome after ketamine induction of anesthesia for caesarea section. *Can J Anesth* ;37: 641.
 - **Baraka A, Siddik S and Assaf B. (1999):** halothane for supplementation of general anesthesia during caesarean section. *Can J Anesth*; 46:95-96.
 - **Barrier G and Sureau G. (1982):** Effects of anesthtics and analgesic drugs on labor, fetus and neonat. *Clinics In obst and Gynecol* ;9:351.

- **Bartkowski RR, Witkowski TA, Azad S et al (1993):** Rocuronium onset of action: A comparison with atracurium and vecuronium. *Anesth Analg* 77: 574.
- **Bechman CR, Beiehl DR, Sitar D and Pacci W. (1986):** Isoflurane potency and cardiovascular effects during short exposure in the fetal lamb. *Can Anesth Soc J* ; 33:41.
- **Bevan DR, Bevan JC, Donati F. (1988):** Muscle relaxants in clinical Anaesthesia, Chicago, Year Book Medical.
- **Biehl DR, Cate J, Waele JG and Stair D. (1983):** Uptake of halothane by the fetal lamb in utero. *Can Anesth Soc J*; 30:24.
- **Bland BA, Lows EG and Downing JW. (1987):** Comparison of midazolam and thiopentone for rapid sequence anesthetic induction for elective caesarean section. *Anesth analgesia* ; 66:1165.
- **Blankenbaker WL, Difazia CA and Berry FA. (1975):** Lidocaine and its metabolites in the newborn. *Anesthesiology*, 42 : 325.
- **Blogy CE, Simpson BR and Bell JA. (1975):** Human placental transfer of A48 165. *Anesthesiology* ; 30:23.
- **Bokesch P, Raymond S, Strichartz G. (1987):** Dependence of lidocaine potency on pH and PCO₂. *Anesth Analg* 66: 9.
- **Bonadio W, Wagner V. (1988):** TAC (tetracaine, adrenaline, cocaine) for the repair of minor dermal lacerations (letter). *Pediatr Emerg Care* 4: 82.
- **Booth PN, Watson MS and Mc-Lead K (1977):** Pancuronium and the placental barrier. *Anesthesia*, 32 :320.
- **Booth, P.N., Watson, M. J. and McLeod, K. (1977).** Pancuronium and the placental barrier. *Anaesthesia*, 32, 320.

- **Bowman WC. (1993)** : Physiology and pharmacology of neuromuscular transmission, with special reference to the possible consequences of prolonged blockade. *Intensive Care Med* 19: S45.
- **Boyd JD, Hamilton WJ (1970)**: The Human placental cambridge, England, Heffer.
- **Boyes R. (1975)**: A review of the metabolism of amide local anaesthetic agents. *Br J Anaesth* 47: 225.
- **Brackbill, Y., Kane, J., Manniello, R. L. and Abramsion, D. (1974)**: Obstetric meperidine usage and assessment of neonatal status. *Anesthesiology*, 40,116.
- **Brandom BW, Woelfel SK, Cook D.R. (1989)**: Comparison of mivacurium and suxamethonium administered by bolus and infusion. *Br J Anaesth* 62 :488.
- **Branney SW, Haenel JB, Moore FA et al (1994)**: Prolonged paralysis with atracurium infusion: A case report. *Crit Care Med* 22:66.
- **Brazelton, T. B. (1961)**: Psychophysiological reactions in the neonate II. Effect of maternal medication in the neonate and his behaviour. *J. Pediat.*, 58,513.
- **Brazetton TB (1961)**: Psychophysiological reaction in the neonate Effect of maternal medication in the neaonate. *J paediat* 58:513.
- **Briggs SG, Freeman PK and Yoff S.J. (1990)**: Drugs in pregnancy and lactation ed. Balumore, William and Wilkins.
- **Bromage PR, Datta S and Dunford LA. (1974)**: Etidocaine and evaluation in epidural analgesia in obstetric. *Can Anesth Soc J*;21:535.

-
- **Brown WU, Bell GC and Lurie A.O. (1975):** Newborn blood levels of liodcaine and mepivacaine in the 1st postnatal day following maternal epidural anesthesia. *Anesthesiology*; 42:698.
 - **Buck MI, Reed M.D. (1991):** Use of non- depolarizing neuromuscular blocking agents in mechanically ventilated patents. *Clin pharm* 10:32.
 - **Caldwell JE, Castagnoli KP, Canfell PC (1988):** Pipecuronium and pancuronium: Comparison of pharmacokinetics and duration of action. *Br J Anaesth* 61 :693.
 - **Caldwell, J., Wakile, L. A., Notarianni, L.J., Smith, R. L., Lieberman, B. A., Jeffs, J., Coy, Y. and Beard, R.W. (1977):** Transphacental passage and neonatal elimination of pethidine given to mother in childbirth. *Brit. J. clin. Pharmacol.*, 4, 715p-716p.
 - **Campbell S., Pearce J.M.F, Jackett, Cohen- over book T, Hernadez (1986):** Qualitative assessment of uteroplacental blood flow: early screening test for high risk pregnancies, *obstet. Gynecol oz*, 649-653.
 - **Campbell Steal (1986):** New Doppler technique for assessing placental blood flow *lancet* 1483,1:635.
 - **Cannon JE, Fahey MR, castagnoli KP (1987):** Continiuons infusion of vecuronium: The effect of anaesthetic agents. *Anaesthiology* 67:503.
 - **Celleno D, Capagno G and Nisini R. (1989):** Neurobehavioral effects of propofol on the neonate following elective caesarean section. *Br J Anesth* ; 62:649.

- **Celleno, Caponga, Emanuelli, Varressei (1993):** Which induction drug C.S A comparison of thiopental sodium, propofol and Midazolam. J clin Anesth; Jul Aug 5(4):284-8.
- **Chang, Wang, Fan and Lieu (1997):** IV infusion of low dose propofol for caesarean section in CS before spinal anesthesia. Acta Anesthesiol Sin; Jun ;35 (2):79-84.
- **Chauvin M, Samii K and Schermann D. (1981):** Plasma morphine concentration after intrathecal administration of low dose of morphine. Br J Anesth; 53:1065.
- **Clark VT, smiley RM and Finster M. (1994):** Uterine hyperactivity after intrathecal injection of fentanyl for analgesia during labor a cause of fetal bradycardia? Anesthesiology;81:1083.
- **Cohen S.E. (1994):** Non obstetric surgery during pregnancy in obstetric anesthesia principle and practice Ed, Chestnut DH Mosby- year book, St Louis, 237.
- **Collins VJ. (1976):** Cyclopropan, principle of anesthesiology 2nd Ed. Henry Kimpton, London, 1563,1976.
- **Cook DR, Wingard LB, Taylor F.H. (1976):** Pharmacokinetics of succinylcholine in infants, children, and adults. Clin pharmacol Ther 20 : 493.
- **Cooper J, Jauniaux E, Gulbis B, Bromley L. (1999):** Placental transfer of fentanyl in early human pregnancy and its detection in the fetal brain. Br J Anesth; 82:929-31.
- **Corke C.B. (2001):** Drugs and obstetric anesthesia : Intravenous anesthetic agent, Drugs and anesthesia (pharmacology for anesthesiology), William and Wilkins, Baltimore.

- **Corke C.B. (2001):** Local anesthetics, Drugs and anesthesia (pharmacology for Anesthesiologist). William and Wilkins, Baltimore.
- **Corke C.B. (2001):** Passage of drugs across the placenta, Drugs and anesthesia (pharmacology for Anesthesiologist). William and Wilkins, Baltimore,.
- **Cosmi EV, Condorelli S and Scorpelli EM. (1974):** Fetal asphyxia induced by thiopental and methohexital in proceedings, fourth European Congress, perinatal medicine.
- **Cosmi E.V. (1976):** Drugs, anesthesia and fetus. In Scarpelli EM and Cosmi EV (eds). Review in perinatal medicine, Vol 1. University Park Press, Baltimore.
- **Costley EC and Way EL. (1965):** Respiratory sensitivity of the newborn infant to meprobamate and morphine. Clin pharmacol Ther 7:454.
- **Coupey M.S. (1997):** Barbiturates and pregnancy. Ped Revol Aug ;18(8):260-4.
- **Covino B, Vassallo H: Local Anesthetics (1976):** Mechanisms of Action and Clinical Use. Orlando, Fla, Grune & Stratton.
- **Covino B. (1984):** Pharmacokinetics of local anesthetic drugs. In Pyrs-Roberts C, Hug CJ (eds): Pharmacokinetics of Anesthesia. Oxford, Blackwell Scientific Publications, P 202.
- **Covino B. (1986):** Pharmacology of local anaesthetic agents. Br J Anaesth 58: 701.
- **Craft JB, Bolan JC, Cooldrakle L.A. (1983):** Placental passage and uterine effect of fentanyl. Anesth Analg; 62:894-8.
- **Crowford J.S. and Rudofsky S. (1965):** Placental transmission of pethidine. Br J Anesth; 37:929.

- **Cummings, A. J. and Whitelaw, A. G. L. (1981):** A study of conjugation and drug elimination in the human neonate. *Brit. J. clin. pharmacol.*, 12,511.
- **Dahlgren G, Christer H, Norman J. (1997):** Intrathecal sufentanil, fentanyl or placebo added to bupivacaine for caesarean section. *Anesth Analg* ;86:1288-93.
- **Dann WL, Hutchinson A and Cartwright D.P. (1987):** Maternal and neonatal responses to alfentanil administration before induction of general anesthesia for caesarean section. *Br J Anesth*; 59:1392-6.
- **Darrah W, Johnston J, Mirakhur R. (1989):** Vecuronium infusion for prolonged muscle relaxation in the intensive care unit. *Crit Care Med* 17:1297.
- **Data S, Sanjoy A and William F.D. (1995):** Clinical effects and maternal and fetal plasma concentration of epidural ropivacaine versus bupivacaine for epidural ropivacaine versus bupivacaine for cesarean section. *Anesthesiology*; 82:1346-1352.
- **Dawes, G.S. (1973):** Theory of fetal equilibration. In: fetal pharmacology, ed. L. Boreus, New York: Raven press.
- **Day D.M. (1978):** Nitrous oxide transfer across the placental and condition of newborn at delivery. *Br J Obstet Gyn*; 85:299,1978.
- **Demetriou M, Depoi J.P., Diakite B. (1982):** Placental transfer of ORG NC45 in women undergoing caesarean section. *Br. J. Anesth.* 54:643.
- **Derek L.J. (1982):** Intervillous space fundamentals of obstetrics and Gynecology, 3rd Ed. Vol 1 :22,.

- **Difenbach C, Mellinghoff H, Grond S, Buzello W. (1992):** Atracurium and vecuronium: Repeated bolus injection versus infusion. *Anesth Analg* 74:519.
- **Doody KM, Carr BR, Rainey WE, Byrd W, Murry BA, Strickler RC, Thomas JL, Mason J.I. (1990):** Hydroxysteroid dehydrogenase activity in glandular and extraglandular human fetal tissues. *Endocrinology* 126 : 2487.
- **Doucet O, Martini L, Loffon M, Jonville A.P. (1998):** Prolonged neuromuscular blockade with mivacurium in newborn. *Ann fr Anesth Reanim*; 17:725-7.
- **Downing JW, Buley RJR and Houlton P.C. (1979):** Etomidate for induction of anesthesia at caesarean section. *Br J Anesth*; 51:135.
- **Downing JW, Mahomed, MC, and Jeal D.E. (1974):** Anesthetic induction for caesarean section Althesin versus thiopentone. *Anesthesia*, 29:689.
- **Drabkova, J., Crul, J.F. and van der kleijn, E. (1973):** Placental transfer of C labelled succinylcholine in nearterm Macaca Mulatta monkeys. *Brit. J. Anaesth.*, 45, 1087.
- **Duvaldestin P, Mazze RI, Nivoche Y. (1981):** Occupational exposures to halothane results in enzyme induction in anesthetists. *Anesthesiology* 54: 57.
- **Ehrnebo, M., Augrell, S., Jalling, B and Boreus, L.O. (1971):** Age differences in drug binding by plasma proteins: studies on human foetuses, neonates and adults. *Europ.J. clin. Pharmacol.*, 3, 189-193.

-
- **Egan TD, Minto C.F. Herm (1996):** Remifentanil versus alfentanil comparative pharmacokinetics and pharmacodynamics in healthy adult male volunteers. *Anesthesiology*; 84:821-33.
 - **Egsele JH, Wright R and Rogge P. (1982):** Newborn and maternal fentanyl level at caesarean section. *Anesth Analg* ; 61 :179.
 - **Elton CD and MacDonald R. (2001):** Maternal and neonatal physiology, Textbook of Anesthesia, 4th Ed, Churchill Livingstone.
 - **Fernado E, Shevede K. and Eddi D. (1989) :** A comparative study of epidural alfentanil and fentanyl for labor pain relief. *Anesthesiology*; 71:A846.
 - **Finster, M., Mark, L.C., Morishima, H.O., Moya, F., perel, J. M., James. L. S. and Dayton, P.G. (1966):** Plasma thiopental concentration in the newborn following delivery under thiopental- nitrous oxide anesthesia. *Am. J, obstet. Gynecol.*, 95,621.
 - **Fleischer A. (1986):** uterine artery Doppler velocimetry in pregnant women with hypertension. *Am J obstet Gynecol*, 154:806.
 - **Flynn P.J. Frank M. and Hugher, R. (1984):** Use of atracurium in caesarean section *Br. J Anesth* 56:599.
 - **Fox H& Flston C.W. (1978):** The development and structure of the placenta. In: *Pathology of the placenta*. London. W.B. Saunders, pp1-37.
 - **Frink EJ, Ghantous H, Malan T.P. (1992):** Plasma inorganic fluoride with sevoflurane anesthesia: Correlation with indices of hepatic and renal function. *Anesth Analg* 74: 231.

-
- **Fuchi I and Nada K. (1979)** : Epidural analgesia in labor. Local anesthetic concentration in maternal and fetal blood. *Acta Med*;412: 375.
 - **Fuchi I and Nada K. (1980)**: Epidural analgesia in labor. Local anesthesia concentration in maternal and fetal blood. *Acta Med*;511:45.
 - **Furuhashi M, Seo H, Mizutani S, Narita O, Tomoda Y, Matsui N. (1989)**: Expression of low density lipoprotein receptor gene in human placenta during pregnancy. *Mol endocrinol* 3: 1252.
 - **Gallagher T.F. (1984)**: Relationship between amniotic fluid volume and maternal plasma volume expansion. *Am J obstet Gynecol*, 146:505.
 - **Gambling DR, Lucas M.R. and Micheal J. (1989)**: Randomized study of combined spinal- epidural analgesia versus intravenous meperidine during labor, Impact on CS delivery rate. *Anesthesiology*; 89:1336-44.
 - **Garton KJ, Yuen P. Meinwald J (1995)**: Stereoselective metabolism of enflurane by human liver cytochrome P450 2E1. *Drug Metab Dispos* 23: 1426.
 - **Gassano, G.B., Ghetti, B., Gliozzi, E. and Hanssion, E. (1967)**: Autoradiographic distribution study of "short-acting" and "long- acting" barbiturates. ³⁵S- thiopentone and ¹⁴C- phenobarbarbitone. *Brit. J. Anaesth.*, 39,11.
 - **Genom A, Leroy G and Mercier C. (1975)**: Anesthesia alfentanil utilising common agent of de narcose. *Cahier de Anesthesiologie*; 23:513.

- **Giles W.B., Trudinger B.J., & Baird J.P. (1985):** Fetal umbilical arterial flow velocity waveform and placental resistance: pathological correlation. *Br J obstet Gynaecol* 92,31-38.
- **Gin T, Gregory MA, and Oh T.E. (1991):** Maternal and fetal levels of propofol at caesarean section. *Anesth Intensive care*;18:180.
- **Goland RS, Wardlay SI, Blum M, Tropper PJ, stark RI. (1988):** Biologically active corticotropin- releasing hormone in maternal and fetal plasma during pregnancy. *Am J obstet Gynecol* 159:884.
- **Goodlin RC, Anderson J.C., Gallagher T.F. (1984):** Relationship between amniotic fluid volume and maternal plasma volume expansion *Am J obstet Gynecol*, 146-505.
- **Grawford, J.S. and Rubofsky, S. (1965).** Placental transmission and neonatal metabolism of promazine. *Brit. J. Anaesth.*, 37,303.
- **Gregory MA and Davidson D.G. (1991):** Plasma etomidate levels in mother and fetus. *Anesthesia*; 46:716,.
- **Gude NM, King RG and Brennecke S.P. (1998):** Autoacid interactions in the regulation of blood flow in human placenta. *Clin Exp pharmacol physiol*; 25:106.
- **Guyton, (2002):** Endocrinology and reproduction (626-628), *Human physiology and mechanisms of diseases*.
- Hamshaw- Thomas A, Rogerson N, Reynolds F. (1984):** Transfer of bupivacaine, lignocaine and pethidine across the rabbit placenta: Influence of maternal protein binding and fetal flow. *Placenta*; 5: 61-70.
- Harmer F.M. (1995):** Anesthesia in obstetrics, Dewhurst's text book of obstetrics and gynecology for postgraduate, Vol1,5th Ed Blackwell science Ltd. London.

- Hahre FW, Hook W and Hon E.H. (1969):** Continuous lumbar epidural anesthesia in obstetric. *Anesth Analg Curr Res*; 48:909.
- Helling HD, Gattereau D, Lefevre Y, Bolte E. (1970):** Steroid production from plasma cholesterol, I. Conversion of plasma cholesterol to placental progesterone in humans. *J Clin Endocrinol Metab* 30:624.
- Helliwell PJ and Hutom A.M. (1950):** Tichloro- ethylene anesthesia. *Anesthesia*; 5:4.
- Hertig A.T. (1962):** The placenta: some new knowledge about an old organ. *Obstet gynecal.* 20: 859.
- Heslop- Christensen, J., Andreassen, F. and Jansen, J. A (1981):** Pharmacokinetics of thiopentone in Caesarean section. *Acta anaesh. Scand.*, 25, 174.
- Hogan Q, Abram S. (1997):** Neural blockade for diagnosis and prognosis: A review. *Anesthesiology* 86: 215.
- Holaday DA, Smith F.R. (1981):** Clinical characteristics and biotransformation of sevoflurane in healthy human volunteers. *Anesthesiology* 54: 100.
- Hollmen Al, Jouppila R and Matta L. (1978):** Neurological activity of infants following anesthesia for CS. *Anesthesiology*, 33:123-45.
- Hunt CO, Naulty JS, Malinow A. (1989):** Epidural butorphanol-bupivacaine for an agesia during labor. *Anesth Analg*;68:323.
- Hunt CO, Naulty JS, Malinow A. (1989):** Preoperative analgesia with subarachnoid fentanyl- bupivacaine. *Anesthesiology*, 71:535.
- Husemeyer RP, Cumming AJ and Rosankiewicz J.R. (1982):** A study of pethidine kinetics and analgesia in woman in labor following IV, IM and peidural administration. *Br J clin pharm*;13:171.
- Iida H, Watanabe Y, Dohi S. (1997):** Direct effects of ropivacaine and bupivacaine on psinal pial vessels in canine. *Anesthesiology* 87:75.

- Irested T, Ekblom A, Olofsson C and Emandersson B.M. (1998):** Continuous epidural infusion with ropivacaine 2.5 mg/ml or bupivacaine 2.5 mg/ml for labor pain relief. *Acta Anesthesiol Scand*; 4 (8):890-6.
- Irving F.C. (1989):** Advantage and disadvantage of barbiturates in obstetrics. *Am J obstet Gynecol*; 160:1184.
- Jeffrey Met (2000):** Metabolism and toxicity of Inhaled Anesthetics, *Anaesthesia*.
- John J Savarese, (2002):** profiles in Anaesthetic practice, A current practice of Relaxation, clinical anaesthesiology.
- Johnson E.f., Herman N., Arney T.L., Gonzalez H, Johnson V., Dowling J.W. (1995):** Bupivacaine transfer across human term placenta: a study using the dual perfused placental model., *Anaesthesiology* 82:459-468.
- Johnson EF, Herman N, Arney TL, Gonzalez H, Johnson V, Downing JW. Bupivacaine transfer across the human term placenta (1995):** A study using the dual perfused human placental model *Anesthesiology*, 82: 459-468.
- Kakinohana M, Fujimine T, Kakinohana O. (1999):** Propofol anesthesia for an emergent caesarean section in a patient with asthma. *Masui*, Aug ; 48(8): 900-2.
- Kan GS, Tong WN, Chan AM and Lou J.W. (1998):** Recovery from mivacurium block with or without atropine following continuous infusion in obstetric patients. *Anesth Intensive Care*; 24(5):5-5-9.
- Kan, Kessin E, Rosen and Samuel (1998):** IV remifentanyl, placental transfer, maternal and neonatal effects. *Anesthesiology*; 88:1467-74.
- Kanto, J, Virtanen R, Lisalo E. (1981):** Placental transfer and pharmacokinetics of atropine after single maternal intravenous and intramuscular

ingestion. Acta. Anaesth. Scand.25,85.

Kaufman, (1993): The placenta and fetal membranes, Dathology of human plucenta, williams obstetrics.

- **Kelly RE, Dimer M, Turneer LS (1993):** Succinylcholine increases intraocular pressure in the human eye with the extraocular muscles detached. Anesthesiology 79:948.
- **Kerkamp H, Gielen M, Edstrom H. (1990):** Comparison of 0.75% ropivacaine with epinephrine and 0.75% bupivacaine with epinephrine in lumbar anesthesia. Reg Anesth 15: 204.
- **Kharasch ED, Armstrong AD, Gunn K. (1995):** Clinical sevoflurane metabolism and disposition. II. The role of cytochrome P 450 2 E1 in fluoride and hexafluoroisopropanol formation. Anesthesiology 82: 1379.
- **Kharasch ED, Karol MD, Lanni C. (1995):** Clinical sevoflurane metabolism and disposition. I. Sevoflurane and metabolite pharmacokinetics. Anesthesiology 82: 1369.
- **Kharasch ED, Thummel K.E. (1993):** Identification of cytochrome P 450 2E1 as the predominant enzyme catalyzing human liver microsomal defluorination of sevoflurane, isoflurane, and methoxyflurane. Anesthesiology 79: 797.
- **Khaw K, Ngankee WD and Critchely H.L. (2000):** Epidural meperidine does not cause hemodynamic changes in the term parturient. Can J Anesth ; 47:155-159.
- **Kivalo 1 and Soarikoski S. (1976):**Placental transfer of 14C- dimethyltubocurarine during caesarean section. Br J Anesth;48: 239.

- **Kivalo I and saarikoski, S. (1977):** Placental transfer of atropine at full term pregnancy. Brit. J Anasesthesia, 49:1017.
- **Knuppel RA and Goodlin R.C. (1994):** Maternal- placental fetal unit. In: current obstetric and gynecologic diagnosis and treatment, 7th edition. Appleton and lange.
- **Kobayashi Y, Ryoichi O, Takeda J. (1992):** Serum and urinary inorganic fluoride concentrations after prolonged inhalation of sevoflurane in humans. Anesth Analg 74: 753.
- **Koop D. (1992):** Oxidative and reductive metabolism by cytochrome P 450-2E1. FASEB J 6: 728.
- **Kopecky EA, Simone C, Knie B, Koren G. (1999):** Transfer of morphine across the human placenta and its interaction with naloxone. Life Sci ; 65 (22):2359-71.
- **Krishna R, Zakowski M and Mark S. (1997):** Sufentanil transfer in the human placenta during in vitro perfusion. Can J Anesth; 44(9): 996-1001.
- **Kriss J, Disk WF, Lwyser KH and Schronz D. (1994):** Thiopentone, thiopentone /ketamine and ketamine for induction of Anesthesia for caesarean section. Eur J Anesthesiol; 11:115.
- **Kristianson B, Mango K and Wickstrom L. (1990) :** Late effects on the infant of Euflurane anesthesia for caesarean section. Acta Anesthesiol Scand ; 24(3) :187-9.
- **Kudoh A and Matsuki A. (2000):** Halothane and sevoflurane decrease nor- epinephrine stimulated glucose transport in neonatal cardiomyocyte. Anesth Analgesia; Nov; 91(5):1151-9.

-
- **Kunert BR, Zupan KJ and Brown D.E. (1987):** Bupivacaine disposition in mother, fetus and neonate after spinal anesthesia for cesarean section. *Anesth Analg* ; 66:407.
 - **Kyegombe D, Franklin C and Turner P. (1973):** Drug metabolic enzymes in the human placenta, their induction and repression *lancet*; 1:405.
 - **Laegried L, Hagberg M, Lunberg A. (1992):** The effect of BZD on the fetus and newborn. *Neuropediatrics*; Feb (23):18-23.
 - **Lanier WL, Milde JH, Michenfelder J.D. (1986):** Cerebral stimulation following succinylcholine in dogs. *Anesthesiology* 64:551.
 - **Latto IP and Waldran B.A. (1977):** Anesthesia for caesarean section. *Br J Anesth* ; 49:371.
 - **Leach L and firth J.A. (1997):** Structure and permeability of human placenta. *Maicrosc Res Tech* ; 38: 137.
 - **Lebowitz PW, Ramsey FM, Savarese J.J. (1981):** Combination of pancuronium and metocurine: Neuromuscular and hemodynamic advantages over pancuronium alone. *Anesth Analg* 60: 12.
 - **LEE C, Kwan WF, Chen B.J and Tsai S.K. (1996):** Neuromuscular blocking effect and placental gradient of pipecurium bromide in elective caesarean section. *Proc Natt. Sci counc Reub China B*; 16(3):119-25.
 - **Levinson G, Shnider SM, Gildea JE and De Lorimer A.A. (1973):** Maternal and fetal cardiovascular and acid- base changes during ketamine anesthesia in pregnant eve. *Br J Anesth*; 45:1111.
 - **Line CA, Schmith VD, Belmont M.R. (1996):** Pharmacokinetic of cisatracurium in patients receiving nitrous oxide / opioid / barbiturate anaesthesia. *Anacsthiology* 84:300.

- **Ling H, Hiroshi S and Tsujimolo C. (2001):** Effects of uterine and umbilical blood flows on the transfer of propofol across the human placenta during in vitro perfusion. *Anesthesia and Analgesia*; 93:151-156.
- **Little B, Chang T, Dill W and Swift A.Y. (1972):** Study of ketamine as an obstetric anesthetic agent. *Am J Obst Gyn*;113:247.
- **Litwiller R.W. (1969):** Succinylcholine hydrolysis. *Anesthesiology* 31:356.
- **Loftus JR, Holholbrook R and Cohen S.E. (1991):** FHR after epidural lidocaine and bupivacaine for elective cesarean section. *Anesthesiology* ; 48 :350.
- **Loquent p, Broughton- pipkin F, Symonds EM, Rubin P.C. (1988):** Blood velocity wave form and placental vascular formation *lancet* 2:1257.
- **Lotto IP and Wainwrght A.C. (1972):** Anesthesia for caesarean section. *Br J Anesth* ; 44: 1050,1972.
- **Mac Donald. (1993):** Perinatal pharmacology, problems in obstetric anaesthesia 20-71.
- **Maddinein VR, Mirakhur RK, McCoy E.P. (1993):** Neuromuscular effect and intubating conditions following mivacurium: A comparison with suxamethonium. *Anaesthesia* 48:940.
- **Malan T.P. (1995):** Sevoflurane and renal function. *Anesth analg* 81: S39.
- **Mandelli, M., Morselli, P. L., Nordio, S., Pardi, J., principi, N., Sereni, F. and Tognoni, G.(1975):** Placental transfer diazepam and its disposition in the newborn. *Clin. Parmacol. Ther.*, 17,565.

- **Manullang RT, Chunk K, and Ega D.T. (2000):** The use of remifentanyl for caesarean section in parturient with recurrent aortic coarctation. *Can J Anesth*; 47:454-459.
- **Markow AG, Mc Guiness GA, Ereuberg A and Kennedy R. L. (1980):** The neonatal neurobehavioral effect of bupivacaine, mepivacaine and 2-chloroprocaine use for pudenda block. *Anesthesiology*; 52:309.
- **Marx GF, Eckstein KL and Halevy S. (1973):** Placental transmission and maternal fetal elimination of flurxene. *Anesth Analg Curr Kes*; 52:654.
- **Marx Gf, Hwang H.S and Chandra P. (1979):** Postpartum premeres with different doses of ketamine. *Anesthesiology*; 50L 163.
- **Marx Gf, Josh CW and Orkin L.R. (1970):** Placental transfer of nitrous oxide. *Anesthesiology*; 32:429.
- **Mather LE, Cousins M.J. (1979):** Local anesthetics and their current clinical use. *Drugs* 18: 185.
- **Matteo RS, Backus WW, McDaniel D.D. (1985):** Pharmacokinetics and pharmacodynamics of d- tubocurarine and metocurine in the elderly. *Anesth Analg* 64:23.
- **McBride, R. J., Dundee, J, W., Moore, J., Toner, W. and Howard, P.J.(1979):** A study of the plasma concentrations of lorazepam in mother and neonate. *Brit. J. Anaesth.*, 51,971.
- **McCowan L.M., Mullen B.M., Ritchie J.W.K. (1972):** Umbilical artery flow velocity wave forms and the placental vascular bed. *Am J obstet. Gynecol* 157,400-402.
- **Meretoja OA, Taivainen T, Wirtavuouri K. (1994):** Pharmacodynamics of mivacurium in infants. *Br J Anaesth* 73:490.

-
- **Mesitelman C, plaud B, Donati F. (1991):** Neuromuscular effects of succinylcholine on the vocal cords and adductor pollicis muscles. *Anesth Analg* 73:278.
 - **Miller RD, Matteo RS, Bental L.Z. (1977):** The pharmacokinetics of d-tubocurarine in man with and without renal failure. *J pharmacol Exp there* 202:1.
 - **Miller RD, Rupp SM, Fisher D.M. (1984):** Clinical pharmacology of vecuronium and atracurium. *Anesthesiology* 61:444.
 - **Miranda AF, Kyni W, Sivalingan N and Sebastianni (1992):** Propofol and Mehoexitone for elective caeserean, section: comparative study. *med J Malysia* ; Dec ;47 (4): 280-6.
 - **Moir DD and Carfy M.J. (1980):** Obstetric anesthesia and anagesia, 2nd Ed. Bailliere, indall, London.
 - **Moore J, McNoble IG and Glynn J.P. (1973) :** The placental transfer of pentazacine and pethidine. *Br J Anesth*; 45 :798.
 - **Moore K.L. (1982):** The developing human, 3rd edn. Philadelphia.
 - **Moore, J., McNabb, T. G. and Glynn, J.P.(1973):** The placental transfer of pentazocine and pethidine. *Brit. J. Anaesth.*, 45,798.
 - **Moreland, T. A., Brice, J. E.J.H., Walker, C.H.M. and parija, A.C. (1980):** Naloxone pkarmacokinetics in the newborn. *Brit. J. clin. Pharmacol.*, 9, 609.
 - **Morgan G.E, Mikhail Ms and Murray M.J. (2002):** obstetric anaeshtia, chinical anaesthiolgy 3rd Ed. Lange Medical, Books.
 - **Morgan GE, Mikiail MS and Murray M. J. (2002):** Obstetric anesthesia. *Clinical anesthesiology* 3rd Ed. Lange Medical books.
 - **Morishima H, Pedersen H, Finster M. (1985):** Bupivacaine toxicity in pregnant and nonpregnant ewes. *Anesthesiology* 63: 134.

- **Morishima HO, Pederson H, Firnter M. (1981):** Toxicity of lidocaine in adult, newborn and fetal sheep. *anesthesiology*; 55:61.
- **Murray JM, Trinick T.R. (1992):** Plasma fluoroide concentrations during and after prolonged anesthesia: A comparison of halothane and isoflurane. *Anesth Analg* 74: 236.
- **Mutsnzaki V, Liy Masuhivo K. (1992):** Trophoblast-derived tumom necrosis factor (induces release of human chorionic gonadotropin J clin Endocrinol Metab 24: 184.
- **Naguib M, Abdulatif M, Al- Ghamdi A. (1993):** Dose- response relationships for edrophonium and neositgmine antagonism of mivacurium- induced neuromuscular block. *Br J Anaesth* 71 : 709.
- **Naguib M. (1994):** Neuromuscular effects of rocuronium bromide and mivacurium chloride administered alone and in combination. *Anesthesiology* 81: 388.
- **Nandi RP, Morrison P.H. and Morgan B.M. (1990):** Effect of general anesthesia on fetus during caesarean section. *Anesth Review*;8:103.
- **Nelson PE, Erickson J.R, Abouleish E. (1996):** FHR changes after intrathecal sufentanil or epidural bupivacaine for labor analgesia incidence and clinical significance. *Anesth Analg*;83:742-6.
- **Nikkola EM, Ekbland KP, and Jukkar J.M. (1997):** Intravenous fentanyl patients controlled analgesia during labor. *Can J Anesth* (44);12:1248-1255.
- **Noelge G, Hinsken H, Buzello W. (1984):** Comparison between the continios infusion of vecuronium and the intermittent administration of pancuronium and vecuronium. *Br J Anaesth* 56:473.

- **Novarrow M. (2000):** Desflurane general anaesthesia for caesarean section compared with isoflurane and epidural anaesthesia, *Anesthesiology*, 35 (4): 232-6.
- **Nybeel- Lindahl G, Carisson C and Ingemarsson L. (1981) :** Maternal and fetal concentration of morphine after epidural administration during labor. *Am J Obs Gyn*;139:20.
- **O' Donoghue, S. E.F. (1971):** Distribution of pethidine and chlorpromazine in maternal, fetal and neonatal biological fluids. *Nature (Lond.)*229, 124.
- **Older, P.O. and Harris, J.M. (1968):** Placental transfer of tubocurarine. *Brit. J. Anaesth.*, 40,459.
- **Olive G and Rey E. (1993):** BZD and pregnancy, transplacental passage, labor and lactation. *Epilepsia*; 9 (4 suppl 2): 87B-96B.
- **Olufalabi AJ, Glass PS, Booth JV and Wakeling H.G. (2000):** A preliminary investigation of remifentanyl as a labor analgesic. *Anesth Analg*; 91:606-8.
- **Ostheimer G.W. (1992):** Principle of perinatal pharmacology. Manual of obstetric anesthesia. Churchill livingstone, New York.
- **Pacific G.M. and Nottoli R. (1999):** Placental transfer of drugs administered to the mother, clinical pharmacology for anaesthesia (28-235-269).
- **Palmer MC, Macuilla EJ, Cork C.R. (1999):** The incidence of FHR changes after intrathecal fentanyl labor analgesia. *Anesth Analg* ; 88: 577-81.
- **Patrick W. Soutter (2003):** Gynaecology, P26,
- **Perault MC, Farreliere S, Minet P, Rembliez C. (2000):** BZD and pregnancy. *Therapie*; Sep-Oct 55(5): 587-95.

- **Percival R. (1980):** Function of placenta, Manual of obstetrics, 14th Ed. Churchill LTD, p43.
- **Phillips TJ and Madodonald R.R. (1971):** Comparative effects of pethidine, trichloroethylene and entonox on fetal and neonatal acid- base and PO₂. Br Med J;3:558.
- **POPPERS PJ and Mostri A.R. (1970):** Maternal and fetal methaemoglobinaemia caused by prilocaine. Acat Anesthesiol Scand;37: 258.
- **Rao GS, Meridian DJ, Tong Y.S. (1979):** Biochemical toxicology of chronic nitrous oxide exposures. Pharmacologist 21: 216.
- **Rathmell JP, Viscomi CM and Ashburn M.A. (1997):** Management of no- obstetric pain during pregnancy and lactation Anesthesia Analgesia ;85:1074-87.
- **Rayburn W, Rathke A, Fuschen MP, Weidner W. (1989):** Fentanyl citrate analgesia during labour. Am J obstet Gynecol ;161:202-6.
- **Refstad, S.O. and Lindbaek, E.(1980):** Ventilatory depression of the newborn of women receiving pethidine or pentazocine. A double-blind comparative trial. Brit. J. Anaesh., 52, 265.
- **Reussel FG, Van der Am, Gribnau F.W. (1998):** Mechanism of drug transfer across human placenta, pharm world sci, Aug 20(4):139-148.
- **Reynolds F, Laishley R, Morgan B, Lee A. (1989):** Effect of time and adrenaline on the feto-maternal distribution of bupivacaine. Br J Anaesth; 62: 509-514.
- **Reynolds S.R.M. (1954):** Haemodynamic characteristics for the fetal circulation. Am J obstet Gynecol 68:69,1954.

-
- **Rice SA, Mazze RI, Baden J.M. (1985):** Effects of subchronic intermittent exposure to nitrous oxide in Swiss Webster mice. *J Environ Pathol Toxicol Oncol* 6: 271.
 - **Ritchie J. W. K. (1995):** The fetus, placental and amniotic fluid, Dewhurst's text book of obstetrics and Gynecology for post graduates 873:82.
 - **Robert A. Knuppel and Robert C. Goodlin (1991):** Maternal- Placental-fetal unit really neonatal physiology, 7-151-159.
 - **Robert W. Shaw (2003):** *Gynaecology*. 23:25..
 - **Rodman JH, Koehntop JE; Brundage DM and Hegland M.G. (1986):** Pharmacokinetics of fentanyl in neonates. *Anesth, analgesia*, 65:227-32.
 - **Rooth G. and Sjostadts (1997):** Placental transfer of gases and fixed acids. *Placenta* ; 18:527.
 - **Rosen A.M. (2002):** Nitrous oxide for relief of labor pain a systemic review. *Am J Obstet Gynecol* ; May 186 (5 supp nature): 51106.
 - **Rosenberg A.A. (2002):** The neonate, delivery room management of the newborn. *Obstetrics, normal and problem pregnancies* 4th ed. Churchill livingstone.
 - **Rudolf stienstra and Anton Burn (2003):** Placental transfer of local anaesthetics agents.
 - **Russel FG, Van der Am, Gribnau F.W. (1998):** Mechanism of drug transfer across the human placenta. *Pharm World Sci*; Aug 20(4) : 139-48.
 - **Schifferli, P-Y and Caldeyro- Barcia R. (1973):** Effect of atropine and beta- adrenergic drugs on the heart rate of human fetus. In : *Fetal pharmacology* PP. 259-278 Ed. By L. Boreus.

-
- **Schneider H and Mohlen K.M. (1979):** Transfer of glutamic acid across the human placenta perused in vitro. Br J Obst Gyn, 86:299.
 - **Schneider SM and Moya F. (1964):** Effects of meperidine in the newborn infant. Am J Obst Gyn; 89:100.
 - **Shearer ES. Hunter J.M. et al., (1991):** Transplacental distribution of atracurium, landonosine and monoquaternary alcohol during elective caesurean section. Br J. anesth; 66 (5): 5512-6.
 - **Shier, R. W., Sprague, A. D. and Dilts, P.V. (1973):** Placental transfer of meperidine HCl. Part II. Am.J. obstet. Gynecol., 115,556.
 - **Shulteus R.R. (1985):** Hemodynamic effects of ketamine and thiopentone durings anesthesia for caesarean section. Can J Anesth Soc, 32:592.
 - **Smiley RM, Ornstein E, Pantuck E.J. (1991):** Metabolism of desflurane and isoflurane to fluoride ion in surgical patients. Can J Anaesth 38: 965.
 - **Smith CE, Donati F, Bevan D.R. (1988):** Dose- response curves for succinylcholine: Single versus cumulative techniques. Anesthesiology 69:338.
 - **Snchez L, Alcavaz, Q and Laguorda. S. (1998):** Placental transfer and neonatal effects of propofol in CS. J Clin pharm there, Feb; 23(11): 19-23.
 - **Sohn YJ, Bencini AF, Scaf A.H.J (1986):** Comparative pharmacokinetics and dynamics of vecuronium and pancuronium in anesthetized patients. Anesth Analg 65:233.
 - **Stephani SJ, Hughes SC and Shnider S.M. (1982):** Neonatal neurobehavioral effects of inhalational analgesia for vaginal delivery. Anesthesiology;56:351.

-
- **Stephen, G. W. and Cooper, L. V. (1977):** The role of analgesic in respiratory depression: a rabbit model. *Anaesthesia*, 32,324.
 - **Stuart L. Stanton (2003):** Surgical anatomy of the uterus, *Gynaecology* 3rd (ed), 23:30.
 - **Sutton TS, Koblin DD, Gruenke L.D. (1991):** Flurode metabolites after prolonged exposure of volunteers and patients to desflurane. *Anesth Analg* 73: 180.
 - **Swartz J, Cumming M. and Biell D. (1987):** the effect of ketamine anesthesia on the acidotic fetal lamb. *Can J Anesth*;34:233.
 - **Thomas G.R. (1976):** Routine phenobarbitone for prevention of neonatal hyperbilirubinaemia. *Obst Gyn*; 47:304.
 - Thomas, J., Climie, C.R. and Mather, L.E. (1969):** The placental - transfer of alcuronium. *Brit. J. Anaesth.*, 41,297.
 - **Thomson AM, Billewicz W2, Hytten F.E. (1969):** The weight or the placenta in relation to birth weight. *Br J obstet gynecal* 76:865.
 - **Thummel K, Kharasch E. Podoll T. (1993):** Human liver microsomal enflurane defluorination catalyzed by cytochrome P 450 2 El. *Drug Metab Dispos* 21: 350.
 - **Tomson, G., Garle, R.I.M. Thalme, B., Nisell, H., Nylund, L. and Rane, A. (1982):** Maternal kinetics and transphacental passage of pethidine during labour. *Brit. J. clin, pharmacol.*, 13,653.
 - **Tucker GT, Boyes RN and Moore D.C. (1980):** Binding of amide type local anesthesia in human plasma. *Anesthesiology*, 33: 304.
 - **Tucker GT. (1986):** Pharmacokinetics of Local Anaesthetics. *Br J Anaesth*; 58: 717-731.

-
- **Van Dyke RA, Baker MT, Janson L. (1988):** Reductive metabolism of halothane by purified cytochrome P450. *Biochem Pharmacol* 37: 2357.
 - **Van Dyke R.A. (1983):** Enflurane, isoflurane, and methoxyflurane metabolism in rat hepatic microsomes from ethanol- treated animals. *Anesthesiology* 58: 22.
 - **Venola N, Passani N, Zambello A and Fonzo R. (2001):** Low concentration ropivacaine in labor epidural analgesia. A prospective study on obstetric and neonatal outcome. *Minerva Ginecol*, 53(6): 397-403.
 - **Vickers, Morgan M, Spencer; Ms Read (199):** Drugs in Anaesthesia and Intensive Care practice, General Anesthetics.
 - **Viscomi CM, Hood D.D. and Eisenach J.C. (1990):** FHR variability after epidural fentanyl during labor. *Anesth Analg* ;71:679.
 - **Wang C, Cheng Y, Leing J. (1999):** Diazepam, with pregnancy and lactation. *Human Yike Da Xue, Baca*: 24 (1):53-5.
 - **Waskell L. (1994):** Metabolism of the volatile anesthetics. In Rice SA. Fish KJ (eds): *Anesthetic Toxicity*. New York, Raven Press. P-49.
 - **Weiskopf RB, Eger EI, Ionescu U.P. (1992):** Desfluran does not produce hepatic or renal injury in human volunteers. *Anesth Analg* 74: 570.
 - **Whitfield R.C. (1995):** The fetus, placenta, Amniotic fluid. Dewhurst textbook of obstetric and Gynecology for postgraduate 5th Ed Blackwell science ltd London.
 - **Wislocvi G.B, Dempsey E.W. (1955):** Electron microscopy of the human placenta. *Anet Rec* 123:133.

- **Wojtezak A.J. (2000):** Haemodynamic effects of halothane and isoflurane in chick embryo. *Anesth Analg*;90:1331-5.
- **Wood M and Wood A. (2001):** Drugs and anesthesia (pharmacology for anesthesiologists.) William and wilkins, Baltimore-Londo.
- **Wood WA, Stanski DR, Curtis J and Rosen M. (1982):** The role of fetal liver in the distribution of thiopentone from mother to fetus, *Anesthesiology*, 57:A 390.
- **Writer C, Gatt G, stienstra E. (1998):** Neonatal outcome and mode of delivery after epidural analgesia for labor with ropivacaine and bupivacaine. *Br J Anesth*; 81:713-717.
- **Zakowski Mark, Angelito and Gilbert, (1994):** Transfer and uptake of alfentanil in human placenta during vitro perfusion. *Anesth Analgesia*; 79:1089-93.