

References

References

Adamson JP, Lewis L, and Stein JD (1959):

Application of abdominal pressure for artificial respiration.
JAMA ; 169 : 1613-1617 .

Alsous F, Amoateng-Adjepong Y, and Manthous C (1999):

Noninvasive ventilation : experience at a community teaching hospital.
Intensive Care Med; 25: 458-63.

Ambrosino N, Foglio K, Rubini F, Clini E, Nava S, and Vitacca M (1995):

Noninvasive mechanical ventilation in acute respiratory failure due to chronic obstructive pulmonary disease: correlates for success.
Thorax ; 50: 755-757.

Annane D, Quera-Salva M, Lofaso F, Vercken J, Lesieur O, Fromageot C, Clair B, Gajdos P, and Raphael J (1999):

Mechanisms underlying effects of nocturnal ventilation on daytime blood gases in neuromuscular diseases.
Eur Respir J ; 13: 157-62

Anthony M (2000):

Review of the role of noninvasive ventilation in the emergency department.
J Accid Emerg Med; 17:79-85.

Anton T, Tarrega J, Giner J, Guell R, and Sanchis J (2003):

Acute physiologic effects of nasal and full-face masks during noninvasive positive pressure ventilation in patients with acute exacerbation of chronic obstructive pulmonary disease.
Respir Care; 48 : 922-25.

Antonelli M, Conti G, Bui M, et al (2000) :

Noninvasive ventilation for treatment of acute respiratory failure in patients undergoing solid organ transplantation: a randomized trial.
JAMA ; 283: 235 – 41.

- Antonelli M, Conti G, Rocco M, Arcangeli A, Cavaliere F, Proietti R, and Meduri G (2002) :**
Noninvasive positive pressure ventilation versus conventional oxygen supplementation in hypoxemic patients undergoing diagnostic bronchoscopy.
Chest ; 121:1149-1154.
- Antonelli M, Conti G, Rocco M, Bui M, de Blasi R, Vivino G, Gasparetto A, and Meduri G (1998) :**
Comparison of noninvasive positive pressure ventilation and conventional mechanical ventilation in patients with acute respiratory failure.
N Engl J Med ; 339 :429-35.
- Antonelli M, Esquinas A, Conti G, Gonzales G, Conflonieri M, Pelaia P, Principi T, Gengoretti C, Beltrame F, Meduri G, Nencini C, Pirottina M, Pasqui M, Di Palma A, And Genovesi S (1999) :**
Risk factors for failure of non-invasive ventilation in acute hypoxemic respiratory failure : a multicenter study.
Intensive Care MED ; 25 : 56.
- Appendini L, Palessio A, Zanaboni S, Carone M, Gukov B, Donner C, and Rossi A (1994):**
Physiologic effects of positive end-expiratory pressure and mask pressure support during exacerbations of chronic obstructive pulmonary disease.
Am J Respir Crit Care Med ; 149: 1069-76
- Auriant I, Jallot A, Herv A, Cerrina J, Adurie F, Fournier J, Lescot B, and Parquin F (2000) :**
Noninvasive ventilation reduces mortality in acute respiratory failure following lung resection.
Am. J. Respir. Crit. Care Med.;164 (7): 1231-1235.
- Aslanian P, El Atrous S, Isabey D, Valente E , Corsi D, Harf A, Lemarie F, and Brochard L (1998):**
Effects of flow triggering on breathing effort during partial ventilatory support.

Am J Respir Crit Care Med ; 57: 135-143.

Bach J (2003):

Successful pregnancies for ventilator user.
Am J Phy Med Rehabil; 82:226-9.

Bach J and Alba A (1990):

Management of chronic alveolar hypoventilation by nasal ventilation.
Chest 1990; 97: 52-72 .

Bach J and Alba A (1990) :

Noninvasive options for ventilatory support of the traumatic high level quadriplegic.
Chest ; 98: 613-619.

Bach J and Alba A (1991):

Total ventilatory support by the intermittent abdominal pressure ventilator.
Chest ; 99: 630-636.

Bach J, Dominique R, Leger P, and Langevin B (1995):

Sleep fragmentation in kyphoscoliotic individuals with alveolar hypoventilation treated by NPPV.
Chest ; 107: 1552-1558

Baudouin S , Blumenthal S, Cooper B, Davidson C, Davison A, Elliot M, Kinnear W , Paton R, Sawicka E, and Turner L (2002):

Non-invasive ventilation in acute respiratory failure .
Thorax ; 57 :192-211.

Belman M, Soo-Hoo G, Kuei J, and Shadmehr R (1990):

Efficacy of positive versus negative pressure ventilation in unloading the respiratory muscles.
Chest ;99:850-856.

Benhamou D, Muir J, and Melen B (1998) :

Mechanical ventilation in elderly patients.
Monaldi Arch Chest Dis; 53:547-51.

Benumof J (1995):

Functions of the respiratory system.

In: Anaesthesia for thoracic surgery, 2nd ed., Philadelphia WB Saunders: 25-24.

Bott J, Carroll M, Conway J, Keilty S, Ward E, Brown A, Paul E, Elliott M, Godfrey R, Wedzicha J, and Moxham J (1993):

Randomized controlled trial of nasal ventilation in acute ventilatory

failure due to chronic obstructive airways disease.

Lancet ; 341: 1555-57.

Bradley D, Holloway R, McLaughlin P, Ross B, Walters J, and Liu P (1992):

Cardiac output response to continuous positive airway pressure in congestive heart failure.

Am Rev Respir Dis ; 145: 377-82.

Brochard L, Mancebo J, and Elliot M (2002):

Noninvasive ventilation for acute respiratory failure.

Respir Care ; 19: 712-21.

Brochard L, Mancebo J, Wysocki M, Lofaso F, Conti G, Rauss A, Simmoneau G, Benito S, Gasparetto A, and Lemaire F (1995):

Noninvasive ventilation for acute exacerbations of chronic obstructive pulmonary disease.

N Engl J Med ; 333: 817-822

Brochard L, Rauss A, Benito S, Conti G, Mancebo J, Rekik N, Gasparetto A, Lemaire F (1994):

Comparison of three methods of gradual withdrawal from ventilatory support during weaning from mechanical ventilation.

Am J Respir Crit Care Med ; 150: 896-903.

Calderini E, Confalonieri M, Puccio P, Francavilla N, Stella L, and Gregoret C (1999):

Patient-ventilator asynchrony during noninvasive ventilation: the role of expiratory trigger.

Intensive Care Med ; 25: 662-67.

Confalonieri M, Potena A, Carbone G, et al (1999):

Acute respiratory failure in patients with severe community-acquired

pneumonia. A prospective randomized evaluation of noninvasive ventilation.

Am J Respir Crit Care Med ;160:1585-91.

Campbell R, Johannigman J, Branson R, Austin P, Mataica G, and Banks G (2002):

Battery duration of portable ventilators : effects of control variable,

peak end-expiratory pressure, and inspired oxygen concentration.

Respir Care; 47:1173-83.

Carrey Z, Gottfried S, and Levy R (1990):

Ventilatory muscle support in respiratory failure with nasal positive pressure ventilation.

Chest ; 97: 150-8.

Celikel T, Sungur M, Ceyhan B, and Karakurt S (1998):

Comparison of noninvasive positive pressure ventilation with standard medical therapy in hypercapnic acute respiratory failure.

Chest ; 114: 1636-42

Chalmers RM, Howard RS, Wiles CM, and Spencer GT (1994):

Use of the rocking bed in treatment of neurogenic respiratory insufficiency.

Q J Med ; 87: 423-9.

Claman D, Piper A, Sanders M, Stiller RA, and Votteri B (1996):

Nocturnal noninvasive positive pressure ventilatory assistance.

Chest ; 110: 1581-1588.

Clarke D, Vaughn L, and Raffin T (1994):

Noninvasive positive pressure ventilation for patients with terminal respiratory failure: the ethical and economic cost of delaying the inevitable are too great.

Am J Crit Care ; 3: 4-5

Clini E (1997):

Patient ventilator interfaces: practical aspect in chronic situation.

Monaldi Arch chest dis.;52: 76-9.

Colice G.L. (1994):

Historical perspective on the development of mechanical ventilation.

In: Tobin M.J. Principles and practice of mechanical ventilation. 1 - 35.

Corrado A, Gorini M, Ginanni R, Pelagatti C, Villella G, Buoncristiano U, Guidi F, Pagni E, Peris A, and DePaolo E(1998):

Negative pressure ventilation versus conventional mechanical ventilation

in the treatment of acute respiratory failure in COPD patients.

Eur Respir J, 12:519-525.

Corrado A, Gorini M, Villella G, and DePaola E(1996):

Negative pressure ventilation in the treatment of acute respiratory failure: an old noninvasive technique reconsidered.

Eur Respir J; 9: 1531-1544.

Corrado A, Confalonieri M, Marchese S, Mollica C, Villella G, Massimo Gorini, and Della Porta R (2002):

Iron Lung versus Mask Ventilation in the treatment of acute on chronic Respiratory Failure in COPD Patients. A multicenter study.

Chest, 121;189-195.

Criner G, Brennan K, Travaline J, and Kreimer D (1999):

Efficacy and compliance with noninvasive positive pressure ventilation

in patients with chronic respiratory failure.

Chest; 116: 667-75.

Dafne C, Falvia F, and Edurado J (2003):

Noninvasive ventilation in pediatrics.

J Pediatr (Rio J); 79:161-8.

Dalziel J (1838):

On sleep and apparatus for promoting artificial respiration.

Br Assoc Adv Sci; 1 : 127 .

Delclaux C, L'Her E, Alberti C, et al (2000) :

Treatment of acute hypoxemic nonhypercapnic respiratory insufficiency

with continuous positive airway pressure delivered by a face mask: a randomized controlled trial.

JAMA ; 284 : 2352-60.

DeLucas P, Tarancon C, Puente L, et al (1993):

Nasal continuous positive airway pressure in patients with COPD in acute respiratory failure. A study of the immediate effects.
Chest ;104:1694-7.

Diaz O, Iglesia R, and Ferrer M (1997) :

Effects of noninvasive ventilation on pulmonary gas exchange and hemodynamics during acute hypercapnic exacerbation of chronic obstructive pulmonary diseases.

Am J Respir Crit Care Med ; 156:1840-45.

Disdier C, Masa J, and Julian G (2002):

Noninvasive ventilation.

Madrid medical group; 1: 231-9.

Drinker P and Shore L (1929):

An apparatus for the prolonged administration of artificial respiration.
J Clin Invest ; 7 :229.

Easa D, Mundie TG, Finn K, Hashiro G, and Balaraman V (1994):

Continuous negative extrathoracic pressure versus positive end-expiratory

pressure in piglets after saline lung lavage.

Pediatr Pulmonol ;17:161-168.

Edward A, Panacek, and Kirk J (2002) :

Role of noninvasive ventilation in the management of acutely decompensated heart failure.

Rev Cardiovasc Med. ;3:35-40 .

Eisenmenger R (1939):

Therapeutic application of supra-abdominal suction and compressed air in

relation to respiration and circulation.

Wien Med Wochenschr ; 89 : 1032-1049 .

Elliott M (2002):

Acute noninvasive ventilation : what are the current indication?
J R Coll Physicians Edinb ; 32:266-274.

Elliott M (2003):

Improving the care for patients with acute sever respiratory disease.
Thorax; 58: 285-88.

Elliott M and Simonds A (1995) :

Nocturnal assisted ventilation using bilevel airway pressure:
 the effect of expiratory airway pressure.
Eur Respir J ;8:436-40

Estop R, Villasante C, De Lucas P et al (2001):

Mechanical ventilation at home .
Arch Bronconeumol ; 37: 142 – 9.

Evans T, Elliott M, Ranieri M et al (2002):

Pulmonary medicine and critical care medicine in Europe.
Eur Respir J; 19: 1202-6.

Ferguson T and Gilmartin M (1995):

CO₂ rebreathing during BiPAP ventilatory assistance.
Am J Crit Care Med ;151:1126-35.

Gabrielli A, Caruso L, Layon A, and Antonelli M (2003):

Yet another look at noninvasive ventilation.
Chest; 124:428-31.

Ganong J (2001):

Respiration.
 In: Review of medical physiology, 18th ed., Appleton and Lange, section VII: 591-625.

Gappa M, Costeloe K, Southall D, Rabbette P, and Stocks J(1994):

Effect of continuous negative extrathoracic pressure on respiratory mechanics and timing in infants recovering from neonatal respiratory distress syndrome.

Pediatr Res ;36:364-372.

Girault C, Briel A, Hellot MF, Tamion F, Woinet D, Leroy J, and Bonmarchand G (2003):

Noninvasive mechanical ventilation in clinical practice: A

2-year experience in a medical intensive care unit.

Crit Care Med ;31(2) :552-9.

Girault C, Richard JC, Chevron V, et al (1997):

Comparative physiologic effects of noninvasive assist-control and pressure support ventilation in acute hypercapnic respiratory failure.

Chest ;111:1639-48

Giuliani R, Mascia L, Recchia F, Carracciolo A, Flora T, and Ranieri V (1995):

Patient-ventilator interaction during synchronized intermittent mandatory

ventilation. Effects of flow triggering.

Am J Respir Crit Care Med ; 151:1-9.

Goldberg A (2002):

Noninvasive Mechanical Ventilation at home.

Chest ; 121:321-324.

Goldberg P, Reissmann H, Maltais F, et al (1995):

Efficacy of noninvasive CPAP in COPD with acute respiratory failure.

Eur Respir J ;8:1894-900.

Goldstein R, Molotiu N, Skrastins R, Long S, Rosie J, Contreras M, Popkin J, Rutherford R, and Phillipson A (1987) :

Reversal of sleep-induced hypoventilation and chronic respiratory failure by nocturnal negative pressure ventilation in patients with restrictive ventilatory impairment.

Am Rev Respir Dis ; 135: 1049-1055.

Granton J, Shapiro C, and Kesten S (2002):

Noninvasive nocturnal ventilatory support in advanced lung disease from
cystic fibrosis.
Respir Care ;47(6):675-681.

Grasso S and Ranieri V (2001):
Proportional assisted ventilation.
Respir Care Clin N Am ;7 :465-73.

Griffiths M and Evans t (2002):
The pulmonary physician in critical care: towards comprehensive
critical
care.
Thorax;57:77-8.

Guerin C, Girard R, Chemorin C, de Varax R, and Fournier G (1997):
Facial mask noninvasive mechanical ventilation reduce the incidence
of
nosocomial pneumonia: a prospective epidemiological survey from a
single ICU.
Intensive Care Med; 23:1024-32.

Guyton A and Hall J (1996) :
Respiration: Pulmonary ventilation .
In: textbook of medical physiology, 9th ed., unit VII; Ch 37: 447-89.

Harding FM, Davies RJO, Stradling JR (1995):
Effects of short term high frequency negative pressure ventilation on
gas
exchange using the Hayek oscillator in normal subjects.
Thorax; 50 : 44-49.

Hart N, Hunt A, Polkay M, Fouroux B, Lofaso F, and Simonds A (2002):
Comparison of proportional assisted ventilation and pressure support
ventilation in chronic respiratory failure due to neuromuscular and
chest
wall deformity.

Thorax; 57: 979-81.

Hartmann H, Jawad M, Noyes J, Samuels M, Southall D (1994):

Negative extrathoracic pressure ventilation in central hypoventilation syndrome.

Arch Dis Child ;70 :418-423.

Haworth C, Dodd M, Woodcock A, et al (2000):

Pneumothorax in adults with cystic fibrosis dependent on nasal intermittent positive pressure ventilation (NIPPV): a management dilemma.

Thorax ;55:620-2.

Hayek Z and Sohar E (1993) :

External high frequency oscillation—concept and practice.

Intensive Care World :10.

Hilbert G, Gruson D, Vargas F, Valentino R, Gbikpi-enissan G, Dupon M, Reiffers j, Jean P, and Cardinaud (2001):

Noninvasive Ventilation in Immunosuppressed Patients with pulmonary

Infiltrates, Fever, and Acute Respiratory Failure.

N Eng. J M , 344;7:481-487.

Hilbert G, Gruson D, Vargas F, et al (2000):

Noninvasive ventilation for acute respiratory failure.

Quite low time consumption for nurses.

Eur Respir J ;16:710-16.

Highcock M, Morrish E, Jamieson S, Shneerson J, and Smith I (2002):

An overnight comparison of two ventilators used in treatment of chronic

respiratory failure.

Eur Respir J ; 20:942-5.

Hilberg R and Jhonson D (1997):

Noninvasive ventilation .

N Engl J Med ; 337:1746-51.

Hill N (1986):

Clinical applications of body ventilators.
Chest ; 90 : 897-905.

Hill N (1994):

Use of the Rocking bed, pneumobelt, and other noninvasive aids to ventilation.

In: Tobin MJ, editor. Principles and practices of mechanical ventilation. New York: McGraw Hill; 1994. p. 413-425.

Hill N (1997):

Complications of noninvasive positive pressure ventilation.
Respir Care ; 42: 432-442 .

Hill N (2002):

Initiation of noninvasive positive pressure ventilation.
Solunum; 4:269-74.

Hill N (2003):

Noninvasive strategies in the acute care setting . What to use for acute pulmonary edema: CPAP or NPPV or neither.
Abstracts of 32nd congress of society of critical care medicine ; San Antino, texas.

Hill N, Carlisle C, and Kramer N (1997):

Does the exhalation valve really matter during bilevel nasal ventilation?
Am J Respir Crit Care Med ; 155: A408 .

Hodes H (1955):

Treatment of respiratory difficulty in poliomyelitis.
In: Poliomyelitis: papers and discussions presented at the Third International Poliomyelitis Conference. Philadelphia: Lippincott; p. 91-113.

Honig E and Ingram R (1998):

Chronic bronchitis, emphysema and airway obstruction.
Harrison's principles of internal medicine. 14th ed.;1451-60.

Iber C, Davies S , and Mahowald M (1989):

Nocturnal rocking bed therapy: improvement in sleep fragmentation in patients with respiratory muscle weakness.

Sleep; 12: 405-412.

Jasmer R, Luce J, and Matthay J (1997):

Noninvasive positive pressure ventilation for acute respiratory failure.

Chest ; 111: 1672-78.

Jones S, Packham S, Hebden M, Smith AP (1998):

Domiciliary nocturnal intermittent positive pressure ventilation in patients with respiratory failure due to severe COPD: long- term follow up and effect on survival.

Thorax ; 53: 495-498

Joris J, Sottiaux T, Chiche J, Desai C, and Lamy M (1997):

Effect of bi-level positive airway pressure (BiPAP) nasal ventilation

on the postoperative pulmonary restrictive syndrome in obese patients undergoing gastroplasty.

Chest; 111: 665-670.

Jounieaux V, Aubert G, Dury M, Delguste P, and Rodenstein D (1995):

Effects of nasal positive-pressure hyperventilation on the glottis in normal sleeping subjects.

J Appl Physiol ; 79: 186-83

Jubran A, Van G, and Tobin M (1995):

Variability of patient-ventilator interaction with pressure support ventilation in patients with chronic obstructive pulmonary disease.

Am J Respir Crit Care Med ; 152:129-36.

Kacmarek R (1999):

NPPV: patient-ventilator synchrony, the difference between success and failure?.

Intensive Care Med ; 25: 645-47

Karner N, Hill N, and Millman R (1996):

Assessment and treatment of sleep disordered breathing in neuromuscular

and chest wall diseases.

Top Pulm Med; 3:336-42.

Keenan S, Powers C, McMormack D, and Block G (2002):

Noninvasive positive pressure ventilation for postextubation respiratory

distress : randomized controlled trial.

JAMA ; 287: 3238-44.

Keenan S, Gregor J, Sibbald W, Cook D, and Gafni A (2000):

Noninvasive positive pressure ventilation in acute setting of sever , acute exacerbation of chronic obstructive pulmonary disease : more effective and less expensive.

Crit Care Med; 28: 2094-102.

Kelly A (2001):

Continuous positive airway pressure: a new modality for the pre-hospital

treatment of pulmonary edema?.

Pre-hospital Immediate Care ;5:12-15.

Kelly A, Geogakas C, Bau S et al (1997):

Experience with the use of continuous positive airway pressure therapy in

emergency management of acute sever pulmonary edema.

Aust N J Med; 27:317-22.

Kosowsky J, Storrw A, and Carleton S (2000):

Continuous and bilevel positive airway pressure in the treatment of cardiogenic pulmonary edema.

Am J Emerg Med ; 18:91-95.

Kramer N, Meyer TJ, Meharg J, et al. (1995):

Randomized, prospective trial of noninvasive positive pressure ventilation in acute respiratory failure.

Am J Respir Crit Care Med ;151:1799-806.

Kwok H, McMormack J, Cece R, Houtchens J, and Hill N (2003):

- Controlled trial of oronasal versus nasal mask ventilation in the treatment of acute respiratory failure.
Crit Care Med; 31: 468-73.
- Leger S and Leger P (1999):**
The art of interface. Tools for administering noninvasive ventilation.
Med Klin; 94:35-9.
- Levy R, Bradley T, Newman S, Macklem P, and Martin J (1989):**
Negative pressure ventilation. Effects on ventilation during sleep in normal subjects.
Chest ;95:95-99.
- Lightowler J, Wedzicha J, Elliott M, and Ram F (2003):**
Non-invasive positive pressure ventilation to treat respiratory failure resulting from exacerbations of chronic obstructive pulmonary disease:
Cochrane systematic review and meta- analysis.
BMJ ; 326 :185.
- Lisboa C, Diaz O, and Fadic R (2003):**
Noninvasive mechanical ventilation in patients with neuromuscular diseases and in patients with chest restriction.
Arch Bronconeumol ; 39: 314 – 320.
- Lobato D and Alises M (2003):**
Noninvasive ventilation.
Arch Bronconeumol ; 39: 566 –79.
- Lockhat D, Langleben D, and Zidulka A (1992) :**
Hemodynamic differences between continual positive and two types of negative pressure ventilation.
Am Rev Respir Dis ;146:677-680.
- Lofaso F, Brochard L, Thierry H, Hubert L, Harf A, and Isabey D (1996):**
Home versus intensive care pressure support devices:
Experimental and clinical comparison.
Am J Respir Crit Care Med ; 153: 1591-1599.

- Lofaso F, Fodil R, Lorino H, Leroux K, Quintel A, Leroy A, and Harf A (2000):**
Inaccuracy of tidal volume delivered by home mechanical ventilators.
Eur Respir J ; 15: 338-341.
- Make B, Hill N, Goldberg A, Bach J, Criner G, Dunne P, Gilmartin M, Heffner J, Kacmarek R, Keens T, McInturff S, O'Donohue W, Oppenheimer E, and Robert D (1998):**
Mechanical ventilation beyond the intensive care unit.
Chest ; 113: 289-344.
- Martin T, Hovis G, Constantion G, Bierman M, Donahoe M, Rogers R, Kreit G, Sciurba F, Stiller R, and Sanders M (2000):**
A randomized prospective evaluation of noninvasive ventilation for acute
respiratory failure.
Am J Respir Crit Care Med; 161:807-13.
- Masa F, Celli B, Riesco J, Cos J, Disdier C, and Sojo A (1997) :**
Noninvasive positive pressure ventilation and not oxygen may prevent
overt ventilatory failure in patients with chest wall diseases.
Chest ; 112: 207-213.
- Masa F, Gonzalez S, Laverty A, Wallis C, and Lane R (2002):**
The use of nasal continuous positive airway pressure to treat
obstructive
sleep apnoea.
Archives of Disease in Childhood ;87:438-443.
- Masip J, Betbese AJ, Paez J, *et al* (2000):**
Non-invasive pressure support ventilation versus conventional oxygen
therapy in acute cardiogenic pulmonary oedema: a randomized trial.
Lancet ;356:2126-32.
- Matthias L (2003):**
Indications of non-invasive ventilation in intensive care: a practical
approach. *Respiratory Support in Critical Care* issue 12 .
- McCormick J, Mehta S, Hipona R, Cece R, and Hill N (1997):**

Face mask vs nasal mask for administration of noninvasive positive pressure ventilation in acute respiratory failure.
Am J Respir Crit Care Med ; 155: 407.

McMinn R (1994):

Lungs.

In: Last's anatomy regional and applied, 9th ed., Churchill Living stone: Ch. 4: 282-6.

Metha S and Hill NS (2001):

Noninvasive ventilation.

Am. J. Respir. Crit. Care Med.;163: 540-577.

Mehta S, Jay G, Woolard R, Hipona R, Connelly E, Cimini D, Drinkwine J, and Hill N (1997):

Randomized prospective trial of bilevel versus continuous positive airway

pressure in acute pulmonary edema.

Crit Care Med ; 25: 620-28.

Meyer T and Hill N (1994):

Noninvasive positive pressure ventilation to treat respiratory failure.

Ann Intern Med ;120:760-70.

Meyer T, Pressman M, Benditt J, McCool F, Millman R, Natarajan R, and Hill N (1997):

Air leaking through the mouth during nocturnal nasal ventilation:

Effect on sleep quality. *Sleep* ; 20: 561-69.

Meduri G (1996):

Noninvasive positive-pressure ventilation in patients with acute respiratory failure.

Clin Chest Med ; 17: 513-53

Miguel J , Diaz P, Ramos D, Pérez Parra J, Buendia J Garcia M, Marcos R, and Moro G (2003):

Analysis of withdrawal from noninvasive mechanical ventilation in patients with obesity-hypoventilation syndrome .

Arch Bronconeumol ; 39 : 292 – 297.

Mohr CH and Hill NS(1990):

Long-term follow-up of nocturnal ventilatory assistance in patients with respiratory failure due to Duchenne-type muscular dystrophy.
Chest; 97: 91-96.

Motley L, Lang L, and Gordon B (1948):

Use of intermittent positive pressure breathing combined with nebulization in pulmonary disease.

Am J Med ; 5: 853-856 .

Monks P, Broomhead C, Dilkes M, and McKelvie P (1995):

The use of the Hayek oscillator during microlaryngeal surgery.

Anaesthesia ;50:865-869

Montserrat J , Martos JA, Alarcon A, Celis R, Plaza V, and Picado C (1991):

Effect of negative pressure ventilation on arterial blood gas pressures and

inspiratory muscle strength during an exacerbation of chronic obstructive

lung disease.

Thorax ; 46: 6-8.

Mundie T , Finn K, Balaraman V, Sood S, and Easa D (1995):

Continuous negative extrathoracic pressure and positive end-expiratory

pressure. A comparative study in Escherichia coli endotoxin-treated neonatal piglets.

Chest ;107:249-255.

Muir, JF, Cuvelier, A, Tengang, B, et al (1997) :
L

Long term nasal intermittent positive pressure ventilation + oxygen therapy (LTOT) versus LTOT alone in severe hypercapnic COPD: preliminary results of a European multicenter trial .

Respir Crit Care Med 155,408.

Nauffal D, Domenech R, Martinez M, Compte L, Macian V, and Perpaz M (2002):

Noninvasive positive pressure home ventilation in restrictive disorders :outcome and impact health related quality of life.

Resp Med; 96:777-83.

Nava S, Ambrosino N, Clini E, Prato M, Orlando G, Vitacca M, Brigada P, Fraccia C, and Rubinni F (1998):

Non-invasive mechanical ventilation in the weaning of patients with respiratory failure due to chronic obstructive pulmonary disease: a randomized study.

Ann Intern Med ; 128: 721-728.

Nava S, Ambrosino N, Zocchi L, and Rampulla C (1990) :

Diaphragmatic rest during negative pressure ventilation by pneumowrap. *Chest* ; 98 : 857-865.

Nava S, Evangelisti I, Rampulla C, Campagnoni ML, Fracchia C, and Rubini F (1997):

Human and financial costs of noninvasive mechanical ventilation in patients affected by COPD and acute respiratory failure. *Chest* ; 111: 1631-1638.

Navalesi P, Fanfulla F, Frigeiro P, Gregoretti C, and Nava S (2000)):

Physiologic evaluation of noninvasive mechanical ventilation delivered with three types of masks in patients with chronic hypercapnic respiratory failure.

Chest ; 28: 1785-1790

Navalesi P and Pollini A (2000):

Acute respiratory failure in patients with severe community acquired pneumonia: A prospective randomized evaluation of noninvasive ventilation.

Am. J. Respir. Crit. Care Med.; 162(2) : 761-762.

Nunn J(1993):

Diffusion.

In: Nunn's applied respiratory physiology; 4th ed., Oxford : Butter Worth-Heinman, Ch 8 :184-207.

Palmer K , Spencer S, Wickramasinghe Y, Wright T, Southall D, and Rolfe P (1995):

Effects of positive and negative pressure ventilation on cerebral blood

volume of newborn infants.

Acta Paediatr ;84:132-139.

Pang D, Keenan SP, Cook DJ, et al (1998):

The effect of positive pressure airway support on mortality and the need

for intubation in cardiogenic pulmonary edema: a systematic review.

Chest ;114:1185-92.

Pankow W, Hijjeh N, Schuttler F, Penzel T, Becker H, Peter J, and Wichert P(1997):

Influence of noninvasive positive pressure ventilation on inspiratory muscle activity in obese subjects.

Eur Respir J ; 10: 2847-2852.

Patel R and Petrini M (1998):

Respiratory muscle performance, pulmonary mechanics, and gas exchange between the BiPAP S/T-D system and Servo Ventilator 900C

with bilevel positive airway pressure ventilation following gradual pressure support weaning.

Chest ; 114: 1390-1396.

Parreira V, Delguste P, Jounieaux V, Aubert G, Dury M, and Rodenstein D(1997) :

Effectiveness of controlled and spontaneous modes in nasal two-level positive pressure ventilation in awake and asleep normal subjects.

Chest ; 112: 1267-77.

Patrick L, Hill N, and Criner G (1999):

Clinical indication for noninvasive positive pressure ventilation in chronic respiratory failure.

Chest;116 :521-34.

Patrick W and Webster K (1996) :

Noninvasive positive-pressure ventilation in acute respiratory distress without prior chronic respiratory failure.

Am J Respir Crit Care Med 1996;153:1005-11.

Penny D, Hayek Z, Rawl E, Rigby M, Reddington A (1992):

Ventilation with external high frequency oscillation around a negative baseline increases pulmonary blood flow after the Fontan operation.

Cardiol Young ; 2 :277-280.

Petros A, Fernando S, Shenoy V, and Al-Saady M (1995):

The Hayek oscillator. Nomograms for tidal volume and minute ventilation using external high frequency oscillation.

Anaesthesia ; 50 : 601-606.

Peltier L (1993):

Obstructive apnoea in artificially hyperventilated subjects during sleep.

J Appl Physiol;5:614-618.

Piper A and Sullivan C (1996):

Effects of long-term nocturnal nasal ventilation on spontaneous breathing during sleep in neuromuscular and chest wall disorders.

Eur Respir J ; 9: 1515-22.

Plant P, Owen J, and Elliott M (2000):

Early use of non-invasive ventilation for acute exacerbations of chronic

obstructive pulmonary disease on general respiratory wards: a multicentre randomised controlled trial.

Lancet ; 355 :1931-35.

Plant P, Owen J, and Elliott M (2000):

One-year prevalence study of acidosis in patients admitted to hospital with an exacerbation of COPD. Implications for noninvasive ventilation.

Thorax; 53: 863-66.

Plant P , Owen J , Parrott S , and Elliott M (2003):

Cost effectiveness of ward based non-invasive ventilation for acute exacerbations of chronic obstructive pulmonary disease: economic analysis of randomised controlled trial.

BMJ ;326:956-959 .

Plum F and Whedon G (1951) :

The rapid-rocking bed: its effect on the ventilation of poliomyelitis patients with respiratory paralysis.

N Engl J Med; 245: 235-240.

Rabatin J and Gay P(1999):

Noninvasive ventilation.

Mayo Clin Proc ;74 :817-20.

Raine J, Cowan F, Samuels M, Wertheim D, and Southall D (1994):

Continuous negative extrathoracic pressure and cerebral blood flow velocity: a pilot study.

Acta Paediatr ;83:438-439.

Rennote M, Baele P, Aubert G, et al (1995) :

Nasal continuous positive airway pressure in the perioperative management of patient with obstructive sleep apnea submitted to surgery.

Chest; 107 :367-74.

Rehder K, Selsler A, and Rodartre J (1997):

Regional intrapulmonary gas distribution in anaesthetized man.

J Appl Physiol; 23:215-26.

Richards G, Cistulli P, Ungar G, Berthon-Jones M, Sullivan C (1996):

Mouth leak with nasal continuous positive airway pressure increases nasal airway resistance.

Am J Respir Crit Care Med ; 154: 182-186.

Robert M (2003):

Noninvasive positive pressure ventilation: The little thing do make the difference.

Respir Care; 48:10.

Romanes G (1997):

The cavity of the thorax: Lungs.

In: Cunningham's manual of practical anatomy, 15th ed., Oxford medical publications, Vol. 2: 30-36.

Russell-Jones D, Treacher D, Lenicker H, Tashanov M, and Spencer T (1989):

Central hypoventilation in a seven year old child following pertussis treated with negative pressure ventilation.

Postgrad Med J ; **65** :768-770.

Samuels M and Southall D (1989) :

Negative extrathoracic pressure in the treatment of respiratory failure in

infants and young children.

BMJ ; **299** : 1253-1257.

Samuels M and Southall D (1998) :

The role of negative pressure ventilation.

Arch Dis Child ; **79** :94.

Schettino G, Chamongkolchart S, HessD, and Kacmarek R (2003) :

Position of exhalation port and mask design affect CO₂ breathing during noninvasive positive pressure breathing.

Crit Care Med; **31**:2178-82.

Schmidt G (2003) :

Noninvasive strategies in the acute care setting . Indication for noninvasive ventilation acute care program.

Abstracts of 32nd congress of society of critical care medicine ; San Antino, texas.

Shneerson J (1991):

Assisted ventilation.5. Non-invasive and domiciliary ventilation: negative pressure techniques.

Thorax; **46** :131-135.

Shneerson J and Simonds K (2002):

Noninvasive ventilation for chest wall and neuromuscular disorders.

Eur Respir J ;**20**:480-7.

Schonhofer B, Geibel M, Sonneborn M, Haidl P, and Kohler D (1997):

Daytime mechanical ventilation in chronic respiratory insufficiency.

Eur Respir J ; **10**: 2840-46.

Schonhofer B, Sonneborn M, Haidl P, et al (1997):

Comparison of two different modes of noninvasive mechanical ventilation in chronic respiratory failure: volume versus pressure

controlled devices.

Eur Respir J ;10:184-91

Simon P, Skatrud J, Badr M, Griffin D, Iber C, and Dempsey J (1991):

Role of airway mechanoreceptors in the inhibition of inspiration during mechanical ventilation in humans.

Am Rev Respir Dis ; 144 :1033-1041.

Skaburskis M, Helal R, and Zidulka A (1987):

Hemodynamic effects of external continuous negative pressure ventilation compared with those of continuous positive pressure ventilation in dogs with acute lung injury.

Am Rev Respir Dis ; 136 :886-891.

Smith I and Shneerson J (1997):

Secondary failure of nasal intermittent positive pressure ventilation using

the Monnal, D.: effects of changing ventilator.

Thorax ; 52: 89-91

Soo Hoo G, Santiago S, and Williams J (1994):

Nasal mechanical ventilation for hypercapnic respiratory failure in chronic obstructive pulmonary disease: determinants of success and failure.

Crit Care Med ; 27: 417-34.

Spalding K and Opie L (1958):

Artificial respiration with the Tunncliffe breathing-jacket.

Lancet ; 1 : 613-615.

Spitzer SA, Fink G, Mittelman M (1993):

External high-frequency ventilation in severe chronic obstructive pulmonary disease.

Chest ; 104: 1698-1701.

Sternburg L and Sternburg D (1986):

View from the seesaw.

New York: Dodd, Mead and Co. ; 1986.

Steven M and Sinha S (2003):

Invasive and Noninvasive Neonatal Mechanical ventilation.
Resp Care ; 48 : 4.

Sullivan C, Issa F, Berthon-Jones M, and Eves L (1981):

Reversal of obstructive sleep apnea by continuous positive airway pressure applied through the nares.
Lancet ; 1: 862-865.

Takami Y and Hiroshi I (2003) :

Beneficial effects of bilevel positive airway pressure after Surgery under
cardiopulmonary bypass.
Interactive cardiovascular and thoracic surgery; 2 (2): 156-9.

Thomson A (1997):

The role of negative pressure ventilation.
Arch Dis Child ; 77 : 454-58.

Tobin M, Jubran A, and Laghi (2001):

Patient ventilator interaction.
Am. J. Respir. Crit. Care Med.;163: 1059-63.

Tsuboi T , Ohi M, Kita H et al (1999):

The efficacy of custom fabricated nasal mask on gas exchange during nasal intermittent positive pressure ventilation.
Eur Resp J;13:152-6.

Turkington P and Elliot M (2000) :

Rationale for use of noninvasive ventilation in chronic ventilatory failure .
Thorax; 54: 417-23.

Tzeng A and Bach J (2000):

Prevention of pulmonary morbidity for patients with neuromuscular diseases.
Chest; 118: 1390-6.

Valetova V, Trembach A, and Vasil'ev V (2002):

Noninvasive assisted lung ventilation during ambulatory laparoscopic cholecystectomy.

Anesteziol Reanimatol ;4 : 60-3.

Viroslav J, Rosenblatt R, Morris-Tomazevic S (1996):

Respiratory management, survival, and quality of life for high-level traumatic tetraplegics.

Respir Care Clin N Am ; 2: 313-322.

Vittica M, Clini E, Pagani N, Bianchi L, Rossi A, and Ambrosino N (2000):

Physiologic effects of early administered mask proportional assisted ventilation in patients with chronic obstructive pulmonary disease and acute respiratory failure.

Crit Care Med; 28: 1791-97.

Wagener S and Headley R (2003):

Cystic Fibrosis: Current Trends in Respiratory Care.

Respir Care ;48(3): 234 – 245.

Wallis C (2000):

Management. Non-invasive home ventilation.

Paediatr Respir Rev ;1(2):165-71.

Weavind L and Wecknen O (2003):

Newer modes of ventilation : an overview.

The internet J of anaesthesiology; 4:4.

Wood K, Lewis L, VonHarz B, and Kollef M (1998):

The use of noninvasive positive pressure ventilation in the emergency department: result of randomized clinical trial.

Chest; 113:1339-46.

Wood K, Flaten A, and Backes (2000):

Inspissated secretion: a life-threatening complication of noninvasive prolonged ventilation.

Respir Care;45:491-93.

William P, Kratochvil J, Ritz R, Hess D, and Kacmarek R (2000):

Pressure support and pressure assist/control; are there difference?. An evaluation of newest intensive care unit ventilators.
Respir Care; 45:1169-81.

Woollam C (1976):

The development of apparatus for intermittent negative pressure respiration.
Anaesthesia ; 3: 666-685 .

Wright J (1947):

The respired rocking bed in poliomyelitis.
Am J Nurs ; 47: 454-455 .

Wysocki M (1999):

Noninvasive ventilation in acute cardiogenic pulmonary edema: better than continuous positive airway pressure?.
Intensive Care Med;25:1-2.

Yamamoto F, Kato R, Sato J, Nishino T (2003):

Anesthesia for awake craniotomy with non-invasive positive pressure ventilation.
Br J Anaesth ; 90: 382-5.

Yang G, Alba A, Lee M, and Khan A (1989):

Pneumobelt for sleep in the ventilator user: clinical experience.
Arch Phys Med Rehabit 70: 707-11.