




REFERENCES

REFERENCES

Aantaa, R.; Kallio, A. and Virtanen, R. (1993):

Dexmedetomidine, a novel α_2 -adrenergic agonist. A review of its pharmacodynamic characteristics.

Drugs of the Future, 18(1): 49-56.

Abdel Hadi, N.; Abdel Raouf, M.; Wedad, M. and Mansou , L. (2003):

The effect of isosorbide dinitrate aerosol in controlling the pressor responses to endotracheal intubation in hypertensive patients.

Egyptian Journal of Anaesthesia, 19: 33-8.

Abernethy, D. and Schwartz, J. (1999):

Calcium antagonist drugs.

New England Journal of Medicine, 341(19): 1447-53.

Abrams, J. (1996):

Therapy of angina pectoris with long acting nitrates which agent and when?

Canadian Journal of Cardiology, 12(Suppl C): 9C-16C.

Adachi, Y.; Aramaki, Y.; Watanabe, K.; Higuchi, H. and Satoh, T. (2001):

Co-administration of nitrous oxide reduces the pressor response against oro-tracheal intubation during induction of anesthesia with propofol infusion at a low rate.

Masui, 50(4): 405-9.

Adachi, Y.U.; Satomoto, M.; Higuchi, H.; and Watanabe, K. (2002):

Fentanyl attenuates the hemodynamic response to endotracheal intubation more than the response to laryngoscopy.

Anesthesia and Analgesia, 95: 1.

Ainsworth, Q. P. and Howells, T. H. (1989):

Transilluminated tracheal intubation.

British Journal of Anaesthesia, 62: 494-7.

Albertin, A.; Casati, A.; Deni, F.; Danelli, G.; Comotti, I.; Grifoni, F. and Fanelli, G. (2000):

Clinical comparison of either small doses of fentanyl or remifentanyl for blunting cardiovascular changes induced by tracheal intubation.

Minerva Anaesthesiol., 66(10): 691-6.

Alexander, R.; Hill, R.; El-Moalem, H.A.; and Gan, T.J. (2001):

Remifentanyl prevents the hemodynamic response to orotracheal intubation.

The Journal of Applied Research, 1: 2.

Al-Shaikh, B. (2002):

A new cuff design prevents N₂O diffusion.

Anaesthesiology, 97: 3.

Angaran, D. M.; Schultz, N. J. and Tschida, V. H. (1986):

Esmolol hydrochloride: an ultrashort-acting, beta-adrenergic blocking agent.

Clin. Pharm., 5: 288-303.

Arndts, D.; Doevendans, J.; Kirsten, R. and Heintz B. (1983):

New aspects of the pharmacokinetics and pharmacodynamics of clonidine in man.

European Journal of Clinical Pharmacology, 24(1): 21-23.

Asai, T. and Morris, S. (1994):

The laryngeal mask airway: its features, effects and role

Canadian Journal of Anaesthesia, 41: 930-60.

Bachmann M. B.; Biscoping, J.; Schurg, R. and Henkelmann, G. (1991):

Pharmacokinetics and pharmacodynamics of local anesthetics

Anaesthesiol. Reanim., 16(6): 359-73.

Bahk, J.H. and Lim, Y.J. (2001):

Use of intracuff lidocaine during general anesthesia.

Anesthesia and Analgesia, 92: 4.

Barak, M.; Ziser, A.; Greenberg, A.; Lischinsky, S. and Rosenberg, B. (2003):

Hemodynamic and catecholamine response to tracheal intubation: direct laryngoscopy compared with fiberoptic intubation. Journal of Clinical Anaesthesia, 15: 2.

Beebe, D.S. (2001):

Complications of tracheal intubation.

Seminars in Anaesthesia, Perioperative Medicine, and Pain, 20: 3.

Bensky, K.P.; Donahue-Spencer, L.; Hertz, G.E.; Anderson M.T. and James R. (2000):

The dose-related effects of bolus esmolol on heart rate and blood pressure following laryngoscopy and intubation.

AANA Journal, 68(5): 437-42.

Benumof, J. L. (1996):

Laryngeal mask airway and the ASA difficult airway algorithm. Anesthesiology, 84: 686-99.

Berkovitz, B.K.B. and Moxham, B.J. (2002):

Head and neck anatomy: A clinical reference, 1st ed., 6: 192-200, 11: 398-410, 12: 416-431.

Published by Martin Dunitz Ltd, a member of the Taylor&Francis group.

Bishop, M. J.; Weymuller, Jr. E. A. and Fink, B. R. (1984):

Laryngeal effects of prolonged intubation.

Anesthesia and Analgesia, 63: 355.

Bonet, S.; Agusti, A. and Arnau, J. M. (2000):

β -Adrenergic blocking agents in heart failure: Benefits of vasodilating and non vasodilating agents according to patients' characteristics.

Arch. Intern. Med., 160: 621.

Boufflers, E.; Maslowski, D. and Menu, H. (1998):

The clinical use of the cuffed oropharyngeal airway (CCPA).

Ann. Fr. Anesth. Reanim., 17: 206-9

Boussofara, M.; Bracco, D. and Ravussin, P. (2001):

Comparison of the effects of clonidine and hydroxyzine on haemodynamic and catecholamine reactions to microlaryngoscopy.

European Journal of Anaesthesiology, 18(2): 75-8.

Brichet, A.; Verkindre, C.; Dupont, J.; Carlier, M. L.; Durras, J. and Wurtz, A. (1999):

Multidisciplinary approach to management of postintubation tracheal stenoses.

European Respiratory Journal, 13(4): 888-93.

Brown, D. L. (1999):

Atlas of Regional Anesthesia, 2nd ed., P: 195-216.

Copyright © W. B. Saunders Company.

Burgoyne, L. and Cyna, A. (2001):

Laryngeal mask versus intubating laryngeal mask: insertion and ventilation by inexperienced resuscitators.

Anaesthesia and Intensive Care, 29(6): 604-8.

Butler, K.H. and Clyne, B. (2003):

Management of the difficult airway: alternative airway techniques and adjuncts.

Emergency Medicine Clinics of North America, 21: 2.

Butterworth, J. F. and Stichartz, G.R. (1990):

Molecular mechanisms of local anesthetics.

Anesthesiology, 72: 711.

Caplan, R. A.; Benumof, J. L. and Berry, F. A. (1993):

Practice guidelines for management of the difficult air way: a report by the American Society of Anesthesiology Task Force on Management of the Difficult Airway.

Anesthesiology, 78: 597-602.

Carpenter, R. L.; Eger, E. and Johnson, B. H. (1986):

Pharmacokinetics of inhaled anesthetics in humans.

Anesthesia and Analgesia, 65: 575-82.

Carrat, X.; Verhulst, J.; Duroux, S.; Pescio, P.; De ars, F. and Traissac, L. (2000):

Postintubation interarytenoid adhesion.

Annals of Otology, Rhinology and Laryngology, 109: 7. 6-40.

Casati, A.; Cappelleri, G.; Fanelli, G.; Magistris, L.; Beccaria, P.; Albertin, A. and Torri, G. (1999):

The pressor response after laryngeal mask or cuffed oropharyngeal airway insertion.

Acta Anaesthesiologica Scandinavica, 43(10): 1053-6.

Casati, A.; Fanelli, G.; Albertin, A.; Deni, F.; Danelli, C.; Grifoni, F. and Torri, G. (2001):

Small doses of remifentanyl or sufentanyl for blunting cardiovascular changes induced by tracheal intubation: a double-blind comparison.

European Journal of Anaesthesiology, 18(2): 108-12.

Cavo, J. W. J. (1985):

True vocal cord paralysis following intubation.

Laryngoscope, 95: 1352-9.

Chang, D.P.; Huang, M.H.; Lei, C.H.; Pang, W.W. and Tung, C.C. (1999):

The effects of tramadol versus fentanyl in attenuating haemodynamic response following tracheal intubation.

Acta Anaesthesiologica Sinica Journal, 37(4): 191.

Combes, X.; Schauvliege, F.; Peyrouset, O.; Motamed, C.; Kirov, K.; Dhonneur, G. and Duvaldestin, P. (2001):

Intracuff pressure and tracheal morbidity. Influence of filling cuff with saline during nitrous oxide anesthesia.

Anaesthesiology, 95: 5.

Crosby, E. T. and Lui, A. (1990):

The adult cervical spine: implications for airway management.

Canadian Journal of Anaesthesia, 37(1): 77-93.

da Silva, O. and Stevens, D. (1999):

Complications of airway management in very-low birth-weight infants.

Biol. Neonate, 40-5.

Dyck, J. B. and Shafer, S. L. (1993):

Dexmedetomidine pharmacokinetics and pharmacodynamics.

Anaesth. Pharm. Review, 1: 238-45.

Ebегue, A.F. (2001):

Second dose thiopentone attenuates the haemodynamic response to laryngoscopy and intubation.

Middle East Journal of Anaesthesiology, 16(1): 81-90.

Ebert, T. J. (1999):

Is gaining control of the autonomic nervous system important to our speciality?

Anesthesiology, 90: 651.

Ebert, T. J.; Hall, J. E.; Barney, J. A.; Uhrich, T. D. and Colvin, M. D. (2000):

The effects of increasing plasma concentrations of dexmedetomidine in humans.

Anesthesiology, 93: 382-94.

Egan, T. D.; Minto, C. F.; Herman, D. J.; Barr, J.; Mui, K. T. and Shafer, S. L. (1996):

Remifentanyl versus alfentanil: Comparative pharmacokinetics and pharmacodynamics in healthy adult male volunteers.

Anesthesiology, 84: 821-33.

Eltzschig, H.K.; Darsow, R.; Schroeder, T.H.; Hettseimer, H. and Guggenberger, H. (2001):

Effect of tracheal intubation or laryngeal mask airway™ insertion on intraocular pressure using balanced anesthesia with sevoflurane and remifentanyl.

Journal of Clinical Anaesthesia, 13: 4.

Engel, T.P.; Applegate II, R.L.; Chung, D.M. and Sanchez, A. (2001):

Management of the difficult airway, 2nd ed., 1(1): 23-40

Copyright © by Cook Incorporated and its licensors.

Estebe, J.P.; Dollo, G.; Corre, P.L.; Naoures, A.L. Cleevanne, F.; Verge, R.L. and Ecoffey, C. (2002):

Alkalinization of intracuff lidocaine improves endotracheal tube-induced emergence phenomena.

Anesthesia and Analgesia, 94: 1.

Ezello, G. C. (2002):

Text book of physiology, 1st ed., 41: 466-71.

Oxford University press.

Farré, R.; Rotger, M.; Ferre, M.; Torres, A. and Navajas, D (2002):

Automatic regulation of the cuff pressure in endotracheally-intubated patients.

European Respiratory Journal, 20(4): 1010-3.

Felten, M.L.; Schmautz, E.; Cerceau, S.D.; Orliaguet, G.A and Pierre A. Carli, P.A. (2003):

Endotracheal tube cuff pressure is unpredictable in children.

Anesthesia and Analgesia, 97: 6.

Figueredo, E. and Garcia-Fuentes, E.M. (2001):

Assessment of the efficacy of esmolol on the haemodynamic changes induced by laryngoscopy and tracheal intubation: a meta-analysis.

Acta Anaesthesiologica Scandinavica, 45(8): 1011-22.

Friederich, J. A. and Butterworth, J. F. (1995):

Sodium nitroprusside: Twenty years and continuing.

Anesthesia and Analgesia, 81: 152-62.

Frink, E. J. and Pattison, B. D. (1989):

Posterior arytenoids dislocation following uneventful endotracheal intubation and anesthesia.

Anesthesiology, 70: 358.

Fujimura, A.; Ebihara, A.; Ohashi, K.; Shiga, T.; Iimagai, Y.; Nakashima, H. and Kotegawa, T. (1994):

Comparison of the pharmacokinetics, pharmacodynamics, and safety of oral (Catapres) and transdermal clonidine in healthy subjects

Journal of Clinical Pharmacology, 34: 260-5.

Gabbott, D.A. (2001):

The airway in resuscitation.

The Middle East Journal of Emergency Medicine, 1: 1.

Georgiou, M.; Parlapani, A.; Argiriadou, H.; Papagianopoulos, P.; Katsikis, G. and Kaprini, E. (2002):

Sufentanil or clonidine for blunting the increase in intraocular pressure during rapid-sequence induction.

European Journal of Anaesthesiology, 19(11): 819-22.

Ghai, B.; Sharma, A. and Akhtar, S. (2001):

Comparative evaluation of intraocular pressure changes subsequent to insertion of laryngeal mask airway and endotracheal tube.

Journal of Postgraduate Medicine (JPGM), 47(3): 181- .

Giles, R. W.; Berger, J. H. and Barash, P. G. (1982):

Continuous monitoring of left ventricular performance with the computerized nuclear probe during laryngoscopy and intubation before coronary artery bypass surgery.

American Journal of Cardiology, 50: 735.

Glass, P. S. A. (1995):

Pharmacology of remifentanyl.

European Journal of Anaesthesiology, 12(Suppl. 10): 73-4.

Goldberg, J. S.; Rawle, P. R.; Zehnder, J. L. and Sladen, J. N. (1990):

Colorimetric end-tidal carbon dioxide monitoring for tracheal intubation.

Anesthesia and Analgesia, 70(2): 191-4.

Gorczyński, R. J. (1985):

Basic pharmacology of esmolol.

American Journal of Cardiology, 56: 3F-13F.

Graf, B. M.; Martin, E. and Basnjak, Z. J. (1997):

Stereospecific effects of bupivacaine isomers on atrioventricular conduction in the isolated perfused guinea pig heart.

Anesthesiology, 86: 410.

Gray, H. (2000):

Anatomy of the human body, 20th ed., Fig. 956, 1a-tl : larynx, 1b- the trachea and bronchi, thoroughly rev. and re-edited by Warren H. Lewis, On-line ed., Published by Bartleby.com.

Greenberg, R. S.; Brimacombe, J. and Berry, A. (1998):

A randomized controlled trial comparing the cuffed oropharyngeal airway and the laryngeal mask airway in spontaneously breathing anesthetized adults.

Anesthesiology, 88: 970-7

Gupta, R.; Ghei, A. and Chhabra, B. (2003):

A study to compare efficacy of cuffed oropharyngeal airway with LMA in anaesthetised spontaneously breathing patients.

Indian Journal of Anaesthesia, 47(4): 289-92.

Habib, A.S.; Parker, J.L.; Maguire, A.M.; Rowbotham, D.J. and Thompson, J.P. (2002):

Effects of remifentanyl and alfentanil on the cardiovascular responses to induction of anaesthesia and tracheal intubation in the elderly.

British Journal of Anaesthesia, 88(3): 430-3.

Hall, V. A. and Guest, J. M. (1992):

Sodium nitroprusside-induced cyanide intoxication and prevention with sodium thiosulfate prophylaxis.

American Journal of Critical Care, 2: 19-27.

Hariri, M. A. and Duncan, P. W. (1989):

Infective complications of brief nasotracheal intubation.

Journal of Laryngology and Otology, 103: 1217.

Harris, M. A. and Kumar, M. (1997):

A rare complication of endotracheal intubation.

Lancet, 350: 1820-1.

Hasuo, H.; Tomiyasu, S.; Hojo, M.; Fujigaki, T.; Fukunaki, M. and Sumikawa, K. (1998):

Effect of ONO-1101, a novel short-acting β -blocker, on hemodynamic responses to isoflurane inhalation and TI.

Journal of Anaesthesia, 12: 115-8.

Heidegger, T.; Gerig, H. J. and Ulrich, B. (2001):

Validation of a simple algorithm for tracheal intubation: daily practice is the key to success in emergencies. An analysis of 13,248 intubations.

Anesthesia and Analgesia, 92: 517-22.

Helmerts, J. H.; van Leeuwen, L. and Zuurmond, W. W. (1994):

Sufentanil pharmacokinetics in young adult and elderly surgical patients.

European Journal of Anaesthesiology, 11: 181-5.

Heringlake, M.; Doerges, V. and Ocker, H. (1999):

A comparison of the cuffed oropharyngeal airway (COPA) with the laryngeal mask airway (LMA) during manually controlled positive pressure ventilation.

Journal of Clinical Anaesthesia, 11: 590-5.

Hirshman, C. A.; Edelstein, G. and Peetz, S. (1982):

Mechanism of action of inhalational anesthesia on airways.

Anesthesiology, 56: 107-11.

Hudson, R. J.; Stanski, D. R. and Burch, P. G. (1983):

Pharmacokinetics of methohexital and thiopental in surgical patients.

Anesthesiology, 59: 215-9.

Hung, O. R.; Pytko, S. and Morris, I. (1995):

Clinical trial of a new lightwand device (Trachlight) to intubate the trachea.

Anesthesiology, 83: 509-14.

Hurford, W. E. (1999):

Orotracheal intubation outside the operating room: Anatomic considerations and techniques.

Respiratory care, 44(6): 615-26.

Irish, J. C. and Gullane, P. J. (1995):

Complications of intubation and other airway management procedures.

Anaesthesiology Clinics of North America, 13: 709-23.

Kamibayashi, T.; Harasawa, K. and Maze, T. (1997):

Alpha-2 adrenergic agonists.

Canadian Journal of Anaesthesia, 44: R13.

Kamra, S.; Wig, J. and Sapru, R.P. (1986):

Topical nitroglycerin. A safeguard against pressor responses to tracheal intubation.

Anaesthesia, 41(11): 1087-91.

Kanaide, M.; Fukusaki, M.; Tamura, S.; Takada, M. Miyako, M. and Sumikawa, K. (2003):

Hemodynamic and catecholamine responses during tracheal intubation using a lightwand device (Trachlight) in elderly patients with hypertension.

Journal of Anaesthesia, 17(3): 161-5.

Kaplan, J. D. and Schuster, D. P. (1991):

Physiologic consequences of tracheal intubation.

Clinics in Chest Medicine, 12(3): 425-32.

Karasawa, F.; Matsuoka, N.; Kodama, M.; Okuda, T.; Mori, T. and Kawatani, Y. (2002):

Repeated deflation of a gas-barrier cuff to stabilize cuff pressure during nitrous oxide anesthesia.

Anesthesia and Analgesia, 95: 1.

Karasawa, F.; Mori, T.; Kawatani, Y.; Ohshima, T. and Satoh, T. (2001):

Deflationary phenomenon of the nitrous oxide-filled endotracheal tube cuff after cessation of nitrous oxide administration.

Anesthesia and Analgesia, 92: 1.

Karasawa, F.; Mori, T.; Okuda, T. and Satoh, T. (2001):

Profile soft-seal cuff, a new endotracheal tube, effectively inhibits an increase in the cuff pressure through high compliance rather than low diffusion of nitrous oxide.

Anesthesia and Analgesia, 92: 1.

Karasawa, F.; Takita, A.; Takamatsu, I.; Mori, T.; Ohshima, T. and Kawatani, Y. (2002):

Rapid deflation of the bronchial cuff of the double-lumen tube after decreasing the concentration of inspired nitrous oxide.

Anesthesia and Analgesia, 95: 1.

Khan, F.A. and Kamal, R.S. (1989):

Effect of buprenorphine on the cardiovascular response to tracheal intubation.

Anaesthesia, 44(5): 394-7.

Kihara, S.; Brimacombe, J.; Yaguchi, Y.; Watanabe, S. Taguchi, N. and Komatsuzaki, T. (2003):

Hemodynamic responses among three tracheal intubation devices in normotensive and hypertensive patients.

Anesthesia and Analgesia, 96(3): 890-5.

Kil, H. K.; Bishop, M. J. and Bedford, R. F. (1995):

Physiological and pathophysiological responses to intubation.

Anaesthesiology Clinics of North America, 13: 361-75.

Kopman, D. F. (2003):

Acute complications of artificial airways.

Clinics in Chest Medicine, 24: 3.

Kumar, N.; Batra, Y.K.; Bala, I.; and Gopalan, S. (1993):

Nifedipine attenuates the hypertensive response to tracheal intubation in pregnancy-induced hypertension.

Canadian Journal of Anaesthesia, 40(4): 329-33.

Kumar, S.; Mishra, M.N.; Mishra, L.S.; and Bathla, S. (2003):

Comparative study of the efficacy of i.v. esmolol, tiltiazem and magnesium sulphate in attenuating haemodynamic response to laryngoscopy and tracheal intubation.

Indian Journal of Anaesthesia, 47(1): 41-44.

Lee, M.A. and Lee, C.H. (1984):

The effect of sodium nitroprusside on response of the cardiovascular system during endotracheal intubation.

Korean Journal of Anesthesiology; 17(4): 281-7.

Louis, W. J.; McNeil, J. J.; Anavekar, S. N.; Conway, E. J.; Workman, B., Howes, L. G.; Drummer, O. H. and Jarrott, B. (1987):

Comparison of pharmacokinetics and pharmacodynamics of adrenoceptor agonists and antagonists as antihypertensive agents.

Journal of Cardiovascular Pharmacology, 10(Suppl. 12) S100-3.

Luscher, T. F. and Cosentino, F. (1998):

The classification of calcium antagonists and their selection in the treatment of hypertension.

British Journal of Anaesthesia, 74: 497-9.

Maguire, A.M.; Kumar, N.; Parker, J.L.; Rowbotham, D.J. and Thompson, J.P. (2001):

Comparison of effects of remifentanyl and fentanyl on cardiovascular response to tracheal intubation in hypertensive

patients.

British Journal of Anaesthesia, 86(1): 90-3.

Maguire, A.; Thompson, J.P.; Guest, C.; Sadler, P.J.; Srupish, J.W. and West, K.J. (2001):

Comparison of the effects of IV alfentanil and esmolol on the cardiovascular response to double-lumen endobronchial intubation.

Anaesthesia, 56(4): 319-25.

Mahajan, R.P.; Grover, V.K.; Sharma, S.L.; and Singh, H (1988):

Intranasal nitroglycerin and intraocular pressure during general anesthesia.

Anesthesia and Analgesia, 67(7): 631-6.

Marty-Ane, C. H.; Picard, E.; Jonquet, O. and Mary, H. (1995):

Membranous tracheal rupture after endotracheal intubation.

Annals of Thoracic Surgery, 60: 1367-71.

Massard, G.; Rouge, C.; Dabbagh, A.; Kessler, R.; Herz, J. G. and Roeslin, N. (1996):

Tracheobronchial lacerations after intubation and tracheostomy.

Annals of Thoracic Surgery, 61(5): 1483-7.

Mayer, O. (1995):

Pharmacokinetics and pharmacodynamics of calcium channel blockers

Vnitr. Lek., 41(3):164-6.

McCulloch, T. M. and Bishop, M. J. (1991):

Complications of translaryngeal intubation.

Clinics in Chest Medicine, 12(3): 507-21.

McIntyre, J. W. R. (1989):

Laryngoscope design and the difficult adult tracheal intubation.

Canadian Journal of Anaesthesia, 36: 94.

Mikawa, K.; Hasegawa, M.; Suzuki, T.; Maekawa, N.; Kaetsu, H.; Goto, R.; Yaku, H. and Obara, H. (1992):

Attenuation of hypertensive response to TI with nitroglycerin.
Journal of Clinical Anaesthesia, 4(5): 367-71.

Mikawa, K.; Nishina, K.; Maekawa, N.; and Obara, H. (1996):

Comparison of nicardipine, diltiazem and verapamil for controlling the cardiovascular responses to tracheal intubation.
British Journal of Anaesthesia, 76(2): 221-6.

Miller, R. (2000):

Anesthesia, 5th ed., P: 1429, 1444-5, 2183.
Copyright © Churchill Livingstone.

Morgan, D. J.; Blackman, G. L. and Paull, J. D. (1981):

Pharmacokinetics and plasma binding of thiopental: Studies in surgical patients.
Anesthesiology, 54: 468-73.

Morgan, G. E.; Mikhail, M. S.; Murray, M. J. and Larrison, M. P. (2002):

Clinical anesthesiology, 3rd ed., Vol. 1, I(5): 60-1, 70-1, II(13): 225.
Copyright © by McGraw-Hill Companies.

Nishikawa, K.; Kawamata, M.; and Namiki, A. (2001):

Lightwand intubation is associated with less hemodynamic changes than fiberoptic intubation in normotensive, but not in hypertensive patients over the age of 60.
Canadian Journal of Anaesthesia, 48: 1148-54.

Nishikawa, K.; Kawana, S. and Namiki, A. (2001):

Comparison of the lightwand technique with direct laryngoscopy for awake endotracheal intubation in emergency cases.
Journal of Clinical Anaesthesia, 13: 4.

Nishikawa, K.; Omote, K.; Kawana, S.; and Namiki, A. (2010):

A comparison of hemodynamic changes after endotracheal intubation by using the lightwand device and the laryngoscope in normotensive and hypertensive patients.

Anesthesia and Analgesia, 90(5): 1203-7.

O'Callaghan-Enright, S. and Finucane, B. T. (1995):

Anesthetizing the airway.

Anesthesiology Clinics of North America, 13(2): 325-36.

Olsson, G. L. (1987):

Bronchospasm during anaesthesia. A computer-aided incidence study of 136929 patients.

Acta Anaesthesiologica Scandinavica, 31: 244-52.

Ovassapian, A. (2001):

Flexible bronchoscopy update: The flexible bronchoscope: A Tool for Anesthesiologists.

Clinics in Chest Medicine, 22: 2.

Panchal, I. D.; Dalal, A. R. and Vasa, C. (2003):

A comparative study between the cuffed oropharyngeal airway and the LMA in spontaneously breathing anaesthetized adults.

Indian Journal of Anaesthesia, 47(3): 194-7.

Parker, J. D. (1996):

Potential problems with intermittent nitrate therapy.

Canadian Journal of Cardiology, 12 (suppl. C): 22-4.

Parker, J. D. and Parker, J. O. (1998):

Nitrate therapy for stable angina pectoris.

New England Journal of Medicine, 338: 520.

Peppard, S. B. and Dickens, J. H. (1983):

Laryngeal injury following short-term intubation.

Annals of Otolaryngology, Rhinology and Laryngology, 92: 327-30.

Perez, P.J.M.; Olmedilla, A.L.; Jimeno, F.C.; and Navia, R. I. (1991):

Effect of an intravenous nitroglycerin bolus on the hemodynamic impact of laryngoscopy and intubation.

Rev. Esp. Anaesthesiol. Reanim., 38(4): 234-7.

Petcu, L. G. and Sasaki, C. T. (1991):

Laryngeal anatomy and physiology.

Clinics in Chest Medicine, 12(3): 415-23.

Polansky, M. (1997):

Airway management: The basics of endotracheal intubation.

The Internet Journal of Academic Physician Assistants, 1: 1.

Pollack, C. V. (2001):

The laryngeal mask airway: a comprehensive review for the emergency physician.

Journal of Emergency Medicine, 20(1): 53-66.

Prys-Roberts, C.; Greene, L. T. and Meloche, R. (1971):

Studies of anaesthesia in relation to hypertension. Hemodynamic consequences of induction and endotracheal intubation.

British Journal of Anaesthesia, 43: 531.

Rathore, A.; Gupta, H.K.; Tanwar, G.L. and Rehman, H. (2002):

Attenuation of the pressure response to laryngoscopy and endotracheal intubation with different doses of esmolol.

Indian Journal of Anaesthesia, 46(6): 449-52.

Richards, A. and McConachie, I. (1995):

The pharmacology of local anesthetic drugs.

CACC, 6(1): 41-7.

Rinde-Hoffman, D.; Glasser, S. P. and Arnett, D. K. (1991):

Update on nitrate therapy.

Journal of Clinical Pharmacology, 31: 697.

Robin, E. D. and McCauley, R. (1992):

Nitroprusside-related cyanide poisoning: Time for urgent, effective intervention.

Chest, 102: 1842-45.

Rouby, J. J.; Andreev, A. and Leger, P. (1991):

Peripheral vascular effects of thiopental and propofol in humans with artificial hearts.

Anesthesiology, 75: 32-42.

Salihoglu, Z. (2002):

Comparison of effects of remifentanyl, alfentanil and fentanyl on cardiovascular responses to tracheal intubation in morbidly obese patients.

European Journal of Anaesthesiology, 19(2): 125-8.

Sasada, M. and Smith, S. (2003):

Drugs in anaesthesia & intensive care, 3rd ed., P: 10-394

Oxford University press.

Scheinin, B.; Lindgren, L.; Randell, T.; Scheinin, H. and Scheinin, M. (1992):

Dexmedetomidine attenuates sympathoadrenal responses to tracheal intubation and reduces the need for thiopentone and perioperative fentanyl.

British Journal of Anaesthesia, 68(2): 126-31.

Schug, S. A.; Zech, D. and Grond, S. (1992):

Adverse effects of systemic opioid analgesics.

Drug safety, 7: 200-13.

Scott, L. J. and Perry, C. M. (2000):

Tramadol: A review of its use in perioperative pain.

Drugs, 60(1): 139-76.

Seegobin, R. D. and van Hasselt, G. L. (1984):

Endotracheal cuff pressure and tracheal mucosal blood flow: endoscopic study of effects of four large volume cuffs.

British Medical Journal (Clin. Res. Ed.), 288(6422): 965-8.

Shipton, E. A. (2000):

Tramadol: Present and future.

Anaesthesia and Intensive care, 28: 363-74.

Singh, M. (2003):

Stress response and anaesthesia altering the peri- and post-operative management.

Indian Journal of Anaesthesia, 47 (6): 427-34.

Singh, S. and Smith, J.E. (2003):

Cardiovascular changes after the three stages of nasotracheal intubation.

British Journal of Anaesthesia, 91(5): 667-71.

Singleton, M. A.; Rosen, J. I. and Fisher, D. M. (1988):

Pharmacokinetics of fentanyl in the elderly.

British Journal of Anaesthesia, 60: 619-22.

Sklar, B.Z.; Lurie, S.; Ezri, T.; Krichelli, D.; Savir, I. and Soroker, D. (1992):

Lidocaine inhalation attenuates the circulatory response to laryngoscopy and endotracheal intubation.

Journal of Clinical Anaesthesia, 4(5): 382-5.

Smith, I.; White, P. F.; Nathanson, M. and Gouldson, R. (1994):

Propofol: An update on its use.

Anesthesiology, 81: 1005.

Soliz, J. M.; Sinha, A. C. and Thakar, D. R. (2002):

Airway Management: A Review and Update.

The Internet Journal of Anesthesiology, 6: 1.

Soltani, H.A. and Aghadavoudi, O. (2002):

The effect of different lidocaine application methods on postoperative cough and sore throat.

Journal of Clinical Anaesthesia, 14: 1.

Somri, M.; Fradis, M.; Malatskey, S.; Vaida, S. and Gaitini L. (2002):

Simple on-line endotracheal cuff pressure relief valve.

Annals of Otolaryngology, Rhinology and Laryngology, 111(2): 90-2.

Sosis, M. and Lazar, S. (1987):

Jaw dislocation during general anesthesia.

Canadian Journal of Anaesthesia, 34: 407-8.

Sosis, M. B. (2003):

The pinch test: a simple technique for determining the pressure of an endotracheal tube cuff.

Journal of Clinical Anaesthesia, 15: 2.

Stanski, D. R. and Maitre, P. O. (1990):

Population pharmacokinetics and pharmacodynamics of thiopental.

Anesthesiology, 72: 412-22.

Stauffer, J. L. (1999):

Complications of endotracheal intubation and tracheostomy.

Respiratory care, 44: 828.

Stein, C. (1993):

Peripheral mechanisms of opioid analgesia.

Anesthesia and Analgesia, 76: 182.

Streitz, J. M. and Shapshay, S. M. (1991):

Airway injury after tracheotomy and endotracheal intubation.

Surg. Clin. North America, 71(6): 1211-30.

Sum, C. Y.; Yacobi, A. and Kartzinel, R. (1983):

Kinetics of esmolol, an ultra-short-acting beta blocker, and of its major metabolite.

Clinical Pharmacology and Therapeutics, 34: 427-34.

Swerdlow, B. N. and Holley, F. O. (1987):

Intravenous anaesthetic agents: pharmacokinetic-pharmacodynamic relationships.

Clin. Pharmacokinet., 12: 79-110.

Takahashi, S.; Mizutani, T.; Miyabe, M.; and Toyooka, H. (2002):

Hemodynamic responses to tracheal intubation with aryngoscope versus lightwand intubating device (Trachlight®) in adults with normal airway.

Anesthesia and Analgesia, 95: 2.

Takita, K.; Morimoto, Y. and Kemmotsu, O. (2001):

Tracheal lidocaine attenuates the cardiovascular response to endotracheal intubation.

Canadian Journal of Anaesthesia, 48(8): 732-6.

Tramer, M. R.; Moore, R. A. and McQuay, H. J. (1997):

Propofol and bradycardia.

British Journal of Anaesthesia, 78: 642-51.

van Vlymen, J. M.; Fu, W. and White, P. F. (1999):

Use of the cuffed oropharyngeal airway as an alternative to the laryngeal mask airway with positive-pressure ventilation

Anesthesiology, 90: 1306-10.

Verdile, V. P.; Chiang, J. L. and Bedger, R. (1990):

Nasotracheal intubation using a flexible lighted stylet.

Annals of Emergency Medicine, 506-10.

Victory, R.A.; Gajraj, N.M.; Pace, N.A.; Ostman, L.P. and White, P.F. (1995):

Nebulized bupivacaine attenuates the heart rate response following tracheal intubation.

Journal of Clinical Anaesthesia, 7(1): 9-13.

Wang, L.; Luo, A. and Wu, X. (1999):

Bolus administration of esmolol for preventing the haemodynamic response to tracheal intubation: a multicentre clinical study.

Zhonghua Yi Xue Za Zhi., 79(11): 828-31.

Wang, Y.M.; Chung, K.C.; Lu, H.F.; Huang, Y.W.; Lin, K.C. Yang, L.C. and Lin, C.R. (2003):

Lidocaine: the optimal timing of intravenous administration in attenuation of increase of intraocular pressure during tracheal intubation.

Acta Anaesthesiologica Sinica Journal, 41(2): 71-5.

Warner, M. E.; Benenfeld, S. M.; Warner, M. A.; Schroeder, L. R. and Maxson, P. M. (1999):

Perianesthetic dental injuries: frequency, outcomes, and risk factors. Anesthesiology, 90: 1302-5.

Weber, S. (2002):

Traumatic complications of airway management.

Anesthesiology Clinics of North America, 20: 3.

Weis, F. R. (1992):

Lightwand intubation for cervical spine injuries.

Anesthesia and Analgesia, 74: 622.

Willens, J. S. and Myslinski, N. R. (1993):

Pharmacodynamics, pharmacokinetics and clinical uses of fentanyl, sufentanil and alfentanil.

Heart lung, 22: 239-51.

Yacobi, A.; Kartzinel, R.; Lai, C. M. and Sum, C. Y. (1983):

Esmolol: a pharmacokinetic profile of a new cardioselective beta-blocking agent.

J. Pharm. Sci., 72: 710-1.

Yang, K. L. (1995):

Tracheal stenosis after a brief intubation.
Anesthesia and Analgesia, 80(3): 625-7.

Yao Fun-sun F. (2003):

Yao & Artusio's Anesthesiology: Problem-oriented patient management, 5th ed., vol. 1, II(14): 352-3.
Copyright © by Lippincott-Raven Publishers.

Yoo, K.Y.; Jeong, S.T.; Ha, I.H.; and Lee, J. (2003):

Nitrous oxide attenuates pressor but augments norepinephrine response to laryngoscopy and endotracheal intubation.
Anesthesia and Analgesia, 96(5): 1516-21.

Yoo, K.Y.; Lee, J.U.; Kim, H.S. and Im, W.M. (2001):

Hemodynamic and catecholamine responses to laryngoscopy and tracheal intubation in patients with complete spinal cord injuries.
Anesthesiology, 95: 3.

Zalunardo, M.P.; Zollinger, A.; Szelloe, P.; Spahn, D.R.; Seifert, B. and Pasch, T. (2001):

Cardiovascular stress protection following anesthesia induction.
Comparison of clonidine and esmolol.
Anaesthesist., 50(1): 21-5.

Zargar, J.A.; Naqash, I.A.; Gurcoo, S.A.; and Din, M.U. (2002):

Comparative evaluation of the effect of metoprolol and esmolol on rate pressure product and ECG changes during laryngoscopy and endotracheal intubation in controlled hypertensive patients.
Indian Journal of Anaesthesia, 46(5): 365-8.