

CONTENTS

	<u>Page</u>
INTRODUCTION	1
REVIEW OF LITERATURE	4
MATERIAL AND METHODS	12
RESULTS	34
DISCUSSION	98
SUMMARY	110
REFERENCES	125
ARABIC SUMMARY	--

LIST OF TABLES

	<u>Page</u>
Table (1): Plasma hormonal levels in immature male mice following intraperitoneal administration of 10 µg prolactin for 30 minutes, 24 hr and 72 hr.	38
Table (2): Plasma hormonal levels in immature female mice following intraperitoneal administration of 10 µg prolactin for 30 minutes, 24 hr. and 72 hr.	39
Table (3): Plasma hormonal levels in virgin mice following intraperitoneal administration of 10 µg of prolactin for 30 minutes, 24 hr and 72 hr.	40
Table (4): Plasma hormonal levels in immature male mice following intraperitoneal administration of 59 µg of prolactin for 30 minutes, 24 hr and 72 hr.	45
Table (5): Plasma hormonal levels in immature female mice following intraperitoneal administration of 59 µg of prolactin for 30 minutes, 24 hr and 72 hr.	46
Table (6): Plasma hormonal levels in virgin mice following intraperitoneal administration of 59 µg of prolactin for 30 minutes, 24 hr and 72 hr.	47

	<u>Page</u>
Table (7): Plasma hormonal levels in immature male mice following intraperitoneal administration of 144 µg of prolactin for 30 minutes, 24 hr and 72 hr.	53
Table (8): Plasma hormonal level in immature female mice following intraperitoneal administration of 144 µg of prolactin for 30 minutes, 24 hr and 72 hr.	54
Table (9): Plasma hormonal levels in virgin mice following intraperitoneal administration of 144 µg of prolactin for 30 min. 24 hr and 72 hr.	55
Table (10): Plasma hormonal levels in immature male mice following intraperitoneal administration of 350 µg of prolactin for 30 minutes, 24 hr and 72 hr.	61
Table (11): Plasma hormonal levels in immature female mice following intraperitoneal administration of 350 µg of prolactin for 30 minutes, 24 hr and 72 hr.	62

- III -

	<u>Page</u>
Table (12): Plasma hormonal levels in virgin mice following intraperitoneal administration of 350 µg of prolactin for 30 minutes, 24 hr and 72 hr.	63
Table (13): Liver uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in immature male mice 24 hr following the intraperitoneal administration of different doses of prolactin.	68
Table (14): Liver uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in immature female mice 24 hr following the intraperitoneal administration of different doses of prolactin.	69
Table (15): Liver uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in virgin mice 24 hr following the intraperitoneal administration of different doses of prolactin.	70
Table (16): Kidney uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in immature male mice 24 hr following the intraperitoneal administration of different doses of prolactin.	74

	<u>Page</u>
Table (17): Kidney uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in immature female mice 24 hr following the intraperitoneal administration of different doses of prolactin.	75
Table (18): Kidney uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in virgin mice 24 hr following the intraperitoneal administration of different doses of prolactin.	76
Table (19): Muscle uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in immature male mice 24 hr following the intraperitoneal administration of different doses of prolactin.	81
Table (20): Muscle uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, C ¹⁴ -Arginine (mg/gm tissue) in immature female mice 24 hr. following the intraperitoneal administration of different doses of prolactin.	82
Table (21): Muscle uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, and C ¹⁴ -Arginine (mg/gm tissue) in virgin mice 24 hr following the intraperitoneal administration of different doses of prolactin.	83

	<u>Page</u>
Table (22): Gonads uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine and C ¹⁴ -Arginine (mg/gm tissue) male mice 24 hr following the intraperitoneal administration of different doses of prolactin.	88
Table (23): Gonads uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine and C ¹⁴ -Arginine mg/gm tissue) in immature female mice 24 hr following the intraperitoneal administration of different doses of prolactin.	89
Table (24): Gonads uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine and C ¹⁴ -Arginine (mg/gm tissue) in virgin mice 24 hr following the intraperitoneal administration of different doses of prolactin.	90
Table (25): Brain uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, and C ¹⁴ -Arginine (mg/gm tissue) in immature male mice 24 hr following the intraperitoneal administration of different doses of prolactin.	94
Table (26): Brain uptake of C ¹⁴ -Galactose, C ¹⁴ -Valine, and C ¹⁴ -Arginine(mg/gm tissue) in immature female mice 24 hr following the intraperitoneal administration of different doses of prolactin.	95

	<u>Page</u>
Table (27): Brain uptake of C^{14} -Galactose, C^{14} -Valine, and C^{14} -Arginine (mg/gm tissue) in virgin mice 24 hr following the intraperitoneal administration of different doses of prolactin.	96