

REFERENCES

REFERENCES

- Abdel-Azim (1992)*: Physiological genetic studies on some Maize inbreds and their hybrids. M.Sc. Genetic Dept. Ain Shams Univ.
- Abd El-Raheem, A.A. and M.A. Rashed (1991)*: Genetic analysis for same bread wheat characters (*Triticum aestivum* L.) and their correlation with some isozyme markers under salt water stress. Zagazig. J. Agric. Res. 18: 445.
- Abdel-Tawab, F.M. and M.A. Rashed (1985)*: Esterase, peroxidase and catalase isozyme polymorphism in zea, Teosinte and sorghum from different origins. Egypt. Genet. Cytol., 14: 275-281.
- Abdel-Tawab, F.M.; A.A. Tayel, S.H.; Hussanien and Samia, D. Antoun (1989)*: Cultivar identification of *Vicia Faba* L. By biochemical genetic markers. 1. Seed protein electrophoresis and isozyme polymorphism. Egypt J. Genet. Cytol. 18: 83-95.
- Abdel-Tawab, F.M.; M.A. Rahed, Eman, M. Fahmy and F.M. El-Domyati (1993)*: Soybean cultivar identification by biochemical genetic markers. Ann. Agric. Sci. 4th Conf. Agric. Dev. Res. Ain Shams Univ. Cairo (Abst).
- Afify, M.R. and M.A. Shousha (1988)*: Effect of low-dose irradiation on soybean protein solubility, trypsin inhibitor activity and protein patterns separated by PAGE. J. Agric. Food Chem. 36: 810.
- Albert, R. and M. Popp (1977)*: Chemical composition of halophytes from the Neusiedler Lake region in Austria. Oekologia 27: 157-170.
- Allkin, R.; D.J. Goyder, F. Biosby, R.J.; White (1986)*: Names and synonyms of species and subspecies in the Viciae: Issue 3. Viciae Database Project, Publication No. 5.

- Anonymous, (1984):* The salt-tolerant rice and oats from tissue culture. The Tissue Culture for Crops Project (TCCP). Newsletter, 3.
- Ashraf, M. and T. McNeilly (1988):* Variability in salt tolerance of nine spring wheat cultivars. J. Agron. Crop. Sci. 160: 14-21.
- Ayers, A.D.; J.W. Brown and C.H. Wadleigh (1952):* Salt tolerance of barely and wheat in soil plots receiving several salinization regimes. Agron. J. 44: 307-3310.
- Bahieldin, (1990):* Genetic analysis of the physiological basis of tolerance to environmental stresses in wheat by means of aneuploidy. Ph. D. Genetic Department, Ain Shams Univ.
- Ball, P.W. (1968):* Flora Europea 2. pp. 128-1336. eds. T.G. Tutin et al. London: Cambridge Univ., Press (C.F. Cubero, 1984).
- Bar-Num, N. and A. Poljakoff-Mayber (1979):* intervarietal differences in the amino acid composition of pea roots as related to their response to salinity. Ann. Bot., 44: 309-314.
- Barrent, N.M. and A.W. Naylor (1966):* Amino acid and protein metabolism in Bermuda grass during water stress. plant Physiol., 41: 1222.
- Barratt, D.H.P. (1979):* Studies on the field bean *Vicia faba* L. report., Long Ashton res. Stn. Publ: Bristol, U.K., 1980, pp. 61-62.
- Bartok, T. and F. Sagi (1990):* A new endosperm-supported callus induction method for wheat (*Triticum aestivum* L.). Plant cell., Tissue and Organ Culture, 22: 37-41.
- Bassiri, A. and I. Rouhani (1977);* Identification of broad bean cultivars based on isozyme patterns. Euphytica, 26: 279-286.
- Bates, L.S.; R.P. Waldren and I.D. Teare (1973):* Rapid determination of free proline for wter-stress studies. Plant and Sol, 39: 205-207.
- Beckmann, J.S. and M. Soller (1983):* Restriction fragment length polymorphism in genetic improvement: methodologies, mapping and cost. Theor. Appl. Genet., 67: 33-43.

- Beckmann, J.S. and M. Soller (1986):* Restriction fragment length polymorphism in plant genetic improvement. In: B.J. Mifflin (ed.). Oxford survey of plant molecular biology, Vol. 3. Oxford Press, Oxford, pp. 196-250.
- Beckmann, J.S. and M. Soller (1990):* Toward a unified approach to genetic mapping of eukaryotes based on sequence tagged microsatellite sires. *Biotechnology*, 8: 930-932.
- Bejaoui, M. (1985):* Interaction between chloride and some phytohormones on soybean (*Glycine max*) growth. *J. Plant Physiol.* 120 (2): 25-110 (In Frwita Engl. Summ).
- Bernstein, L. (1975):* Effect of salinity and sodicity on plant growth. *Ann. Rev. Phytopathol.*, 12: 295.
- Bonierbale, M.W.; R.L. Paisted and S.D. Tanksley (1988):* RFLP maps based on a common set of clones reveal modes of chromosomal evolution in potato and tomato. *Genetics*, 120: 1095-1103.
- Boond, D.A. (1976):* Field bean *Vicia faba* (Leguminosae-Papilionatae). In evolution of crop plants. Ed. by N.W. Simmonds. Longman, London, pp. 179-182.
- Boulter, D.; D.A. Thurman and E. Derbyshire (1967):* A disc electrophoretic study of globulin proteins of legume seeds with reference to their systematics. *New Phytologist*. 66: 27-36.
- Busse, G. (1986):* In vitro cultivation of *vicia faba* and induction of morphogenesis. *Vegetables*. 105 (1/2) 97-104.
- Cardy, B.J. and W.D. Beversdrof (1984):* Identification of soybean cultivars using isoenzymes electrophoresis. *Seed Sci. And Technol.*, 12: 943.
- Chakrabarti, T. and G. Subrahmanyam (1985):* Analysis of DNA from related and diverse species of barley. *Plant Sci.*, 42: 183-190.

- Change, Yin-Fu; W.C. Wang; C.Y. Warfield; H.T. Nguyen and J.R. Wong (1991):** Plant regeneration from protoplasts isolate from long-term cell cultures of wheat (*Triticum aestivum* L.) Plant Cell Reports 9: 611-614.
- Champman, S.R. and L. Hart (1977):** Developing salt tolerant crop varieties. P. 10-11. In: Warren (ed.) Now, Vol. 13, Montana State University.
- Chauhan, R.P.S.; S.P.C. Chauhan and D. Kumar (1980):** Free proline accumulation in cereals in relation to salt tolerance. Plant and Soil 57: 167.
- Chen, S.; Y. Guo and Y. Liu (1990):** The NaCl-tolerance of proline analog resistant variant in rice. (Abst). 7th International Congress in Plant Tissue and cell Culture. p. 150 Amsterdam.
- Chen, D.X. (1984):** Induction of callus and plantlet regeneration from *Vicia faba* embryos (Preliminary report). Acta Agric. Univ. Pekinesis 10 (3): 331-332.
- Chen and Tabaeizadeh (1991):** Alteration of gene expressis on in tomato plants (*Lycopersicon esculentum*) by drought and salt stress. Biological Science, Univ. of quebec, Montreal, Canda.
- Collander, R. (1941):** Selecying absorption of cations by higher plants plant physiol. 16e: 691-720.
- Conger, B.V. (1986):** Cloning agricultural plants via in Vitro techniques. CRC press. Inc. Florida.
- Cooke, R.J. (1990):** The use of electrophoresis for the distinctness testingt of varieties of autogamous species. Plant Breed. Abst. 60: 2902.
- Cubero, J.I. (1984):** Problems and perspectives in breeding for protein content in *Vicia faba*. FABIS news letter No. 9, 1-9.
- Danielsson, C.E. (1949):** Seed globulins of the Gramineae and Leguminosae Biochem. J., 44: 3387-400.

- Dix, P.J.; U.A. Mclysaght and A. Plunkkett (1986):** Salt stress: Resistance mechanisms and in vitro selectin procedures. In: Plant Tissue Culture and Its Agricultural Application. eds. Withers, L.A. and P.G. Alderson. . 469-478. London, Boston, Durban, singpaora.
- Dregne, H.L. (1964):** Plant response to fertilizers on a saline soil. N. Mex. Agric. Exp. Sta, Res. Rpt. 94 (after Agric. res. Her. 49 (2): 25).
- Durrant, M.J.; A.P. Draycott and P.A. Payne (1974):** Some effects of sodium chloride on germination and seedling growth of sugar beet. Ann. Bot. 38: 1045-1051.
- Dweikat, I.; S. Mackkenzie, M. Levy and H. Ohm (1993):** Pedigree assessment using RAPPD-DGGE in sereal crop species. Theor. Appl. Genet., 85: 497-505.
- Epstein, E. (1972):** Physiological genetics of plant nutrition. in "Mineral nutrition of plants principles and perspectives" Epstein, E. (ed), John Wiley and Sons, New York, pp. 325-344.
- Epstein, E. (1977):** Genetic potential for solving problems of soil mineral stress: Adaptation of crops to salinity. In: M.J. Wright (ed.) Proc. of workshop, Cornell Univerity Exp. Stn. Special Bull., New York.
- Epstein, E.; J.D. Norlyn, D.W. Rush; R.W. Kingsbury; D.B. Kelley; G.A. Cunningham and A.F. Wrona (1980):** Saline culture of crops: a genetic approach. Science 210: 399-404.
- Ericson, M.C. and S.H. Alfinito (1984):** Proteins pproduced during salt stress in tobacco cell cultures. Plant Physiol. 74: 506-509.
- Fahmy, Eman, M. and K.A. Okasha (1992):** Phylogenetic relationships and biochemical genetic markers as fingerprints in two fragaria species. J. Agric. Sci. Mansoura Univ. 17: 335.
- Fedchenko, B.A. (1948):** Vicia. In flora of the URSS, ed. V.L. Kormrov, Moscow. (C.F. cubero 1984).

- Forde, M.B. and S.E. Gardiner (1990):* Electrophoretic seed protein banding patterns as genetic resource discreptors. Interant. Workshop on tissue culture for the conservation of aidodiversity and Genet. Resources. May, 1990 Kuala-Lumpur, Malaysia.
- Forster, B.P.; J. Goorham and T.E. Miller (1987)K:* Salt tolerance of an amphiploid betwee *Triticum aestivum* and *Aggropyron junceum*. Plant breeding 98: 1-8.
- Fox, D.J.; D.A. Thurman, and D. Boulter (1964):* Studies on the proteins of seeds of the Leguminoseae-1. Albumins. Phytochemistry, 3, 417-419.
- Francois, L.E.; E.V. Mass; T.J. Donovan and V.L. Youngs. (1986):* Effect of salinity on grain yield and quality, vegetative growth and germination of semi-dwarf and durum wheat. Agron. J. 78: 1053-1058.
- Gates, P. and D. Boulter (1979):* The sue of seed isoenzymes as an aid to the breeding of field beans (*Vica faba* L.). The New phytologist, 83 (3): 783-792.
- Gawel, N.J. and R.L. Jarett (1991):* A modified CTAB DNA-Rep., 91: 262-266.
- Goodrich, W.J.; R.J. Cooke; A.G. Morgan (1985):* The application of electrophoresis to the characterization of cultivars of vlcia fab L. vegetables, FAbis No. 13, 8-11.
- Gorman, M.B. and Y.T. Kiang (1977):* Variety-secific electrophoretic variants of flour soybean enzymes. Crop Sci., 17: 963.
- Hassa, N.S. and D.A. Wilkins (1988):* Rapid electrophoresis of Oat (*Avena sativa* L.) prolamins from single seeds for cultivar identification. Cereal Chem. 65: 153-154.
- He, S.; H. Ohm and S. Mackenzie (1992):* Detection of DNA sequence polymophisms among wheat varieties. Theor. Appl. Genet., 84: 573-578.

- Heikal, M.M. and Shaddad, M.A. (1981):* Allevation of osmotic stress on seed germination and seedling growth of cotton pea and wheat by proline. *Phyton* 22: 275-287.
- Heikkal, M.M.; M.A. Shaddad and A.M. Ahmed (1982):* Effect of water stress and gibberillic acid on germination of flux, sesone and onion sees. *Biologia Plantarum*, 24: 124-129.
- Henkel, P.A. (1964):* Physiology of plants under drought. *Ann. Rev. Plant Physiol.*, 15: 363.
- Heyser, J. W.; M.W. Nabors; C. Mackinnon; T.A. Dykes; K.J. Demott; D.C. Kautzman and A. Mujeeb-Kazi (1985):* Long-term, high frequency plant regeneration and the induction of somatic embryogenesis in callus cultures of wheat (*Triticum aestivum* L.). *Z. Pflanzenphysio.* 94: 218-233.
- Hsiao, T.G.; E. Acerado; E. Fereres and D.W. Henderson (1976):* Stress metabolism. *Philos. Trans. R. Soc. Lond. B.* 273-479-500.
- Hyghes, S.A. and P.A. Murphy (1983):* Varietal influence on the quantity of glycinin in soybean. *J. Agric. Food Chem.* 31: 376.
- Hurkman and Tamaka (1987):* *Plant physiol.* 83: 517-524.
- Hussein, A.; H. Ramirz; W. Busflux and W. Roca (1986):* Feild beans (*Phaseolus vulgaris* L.) Cultivar identification by electrophorograms of cotyledon stroage prooteins. *Euphytica*.
- Hurkman, W.J.; Fornari, C.B. and Tanaka, C.K. (1989):* Acomparision of the effect of salt on polypeptides and granslatable M RNAS in roots of a salt-tolerant and a salt-sensitive cultiver of b arley. *Am. Soc. of plant Physiol.*, 90. (4), 1444-1456.
- Hutton, E.M. (1971):* Variation in salt response between tropical pasture legumes. *Sabro Newsletter*, 3; 75-81.
- Iglesias, L. (1984a):* peroxidase isoenzymes in roots of soybean (*G. max* L.) cultivars. *Plant Breed. Abst.*, 54: 9352.

- Innis, M.A.; D.H. Gelfand; J.J. Sninsky and T.J. White (1990):** PCR protocols, Ed. 1. Academic Press, San Diego, p. 482.
- Jacoby, B. and A. Ratner (1974):** Mechanism of sodium exclusion in bean and corn plants. A re-evaluation. In X. Wehrmann (ed.) J. Plant Analysis and Fertilizer problems. German Society of Plant Nutrition, Hannover, P. 175.
- Johnson, C.M.; P.R. Stoudi, R.C.; Broysere and A.B. Cariton (1957):** Comparative chlorine requirement of different plants species. Plant And Soil., 337.
- Kahler, A.L.; S. Hecht-Pagliuso and R.W. Allard (1981):** Genetics of isozymes variants in barley. 11. 6-Phosphogluconate Dehydrogenase, Glutamate Oxallate Transaminase and Acid Phosphatase. Crop Sci. 21: 536.
- Karadimova, M.V. (1990):** Selection for NaCl tolerance in callus cultures of wheat (*Triticum aestivum* and *T. durum*). (Abst.) 7th International Congress on plant Tissue and Cell culture. P. 157. Amsterdam.
- Kessmann, H.; D.A. Choudhary and R.A. Dixon. (1990):** Stress responses in alfalfa (*Medicago sativa* L.) 111. Induction of medicarpin and cytochrome P450 enzyme activities in elicitor-treated cell suspension cultures and protoplasts. Plant cell reports 9: 38-41.
- Khan, S.S. and K.H. Sheikh (1977):** Effect of different levels of salinity on seed germination and growth of *Capasicum annum* Var. Longum. *biologia (lohore)*, 22 (1): 15-26.
- Kiang, Y.T.; Y.C. Chaing and C.J. Bult. (1987):** Genetic study of glutamate oxaloacetic transaminase in soybean. *Genome*, 29: 370.
- Kingsbury, R.W. and E. Epstein (1984):** Selection for salt resistance spring wheat. *crop Sci.* 24: 301.
- Kingsbury, R.W. and E. Epstein (1986):** Salt sensitivity in wheat. *Plant Physiol.* 80: 651-654.

- Kishk, E.T. and A.a. Shalaby (1984a)*: Some experimental tests of Ethephon "Ethrol" as an agent for improving salt tolerance in some field crops. Desert Inst. Bull., A.R.E., 34. No. (1,2): 355-368.
- Kishor, P.B.K. and G.M. Reddy (1992)*: Salt stress in callus cultures of rice (*Oryza sativa* L.) Advances in gene technology: Feeding the world in the 21 st century. proceedings of the 1992. Miami Bio/Technology Winter Symposium.
- Klyshev, L.K. and N.M. Rakova (1964)*: Effect of salinization on the protein content in root of pea seedlings. Trudy Instituta botaniki AN KazSSR, 20: 156-165. (In Russian).
- Kochert, G.; T. Halward; W.D. Branch and C.E. Simpson (1991)*: RFLP variability in peanut (*Arabidopsis hypogaea* L.) cultivars and wild species. Theor. Appl. Genet., 81: 565-570.
- Krawetz, S.a. (1989)*: The polymerase chain reaction: opportunities for agriculture. Ag. Biotech. News. Info., 1:897-902.
- Kursanov, A.L. (1967)*: Plant physiology in the golden age of experimental biology. Izvestiya ANSSSR, Seriya Biologicheskaya, 5:695-701.
- Ladizinsky, G. (1975)*: Seed protein electrophoresis of the wild and cultivated species of section *Faba* of *Vicia*. Euphytica 24 (3): 785-688.
- (1983): Study of evolutionary problems by means of seed protein electrophoresis. In seed protein, Biochemistry, Genetics, nutritive value. Ed by W. Gottschalk, and H.P. Muller. Puple: the Hague, Netherlands; Martinus Nijhoff, Dr W. Junk. pp. 481-498.
- and T. Hymowitz (1979): Seed protein electrophoresis in taxonomic and evolutionary studies. Theoretical and applied Genetics 54, 145-151.
- Laemmli, U.K. (1970)*: Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature 227: 680-685.

- Landry, B.S.; R.V. Kesseli; B. Farrara and R.W. Michelmore (1987):** Genetics 116: 331-337.
- Lee, T.E.G. and O.J. Crocomo (1980):** Protein pattern changes in germinating bean seed. Turrialba, 30: 87.
- Lerner, H.R. (1985):** Adaptation to salinity at the plant cell levels. Plant and Soil 89: 3-14.
- Mahmoud, S.H.; J.A. Gatenhous, D. Boulter (1984):** inheritance and mapping of isoenzyme in pean (*Pisum sativus* L.). Thor. and Appl Genet., 68: 559.
- Malki, E. and Y. Waisel (1987):** Effect of pressure on germination of seeds of wheat (*Triticum aestivum* Cv. Barqai) in saline and in non-saline media. *Physiol. Plantarum* 70: 733-77.
- Marcshner, H. and H. ossenberg-Neubaus (1976):** Longstreckent ransport von natrium in bohnepflanzen.Z. Pflanzenernakr. dung Bodenked. 193-129-142.
- Mass, E.V. and G.J. Hoffman (1977):** Crop salt tolerance current assessment. ASCE. J. Irrig. and Drainage Div. 103: 115-134.
- Mass, E.V. and R.H. Nieman (1978):** Physiology of plant tolerance to salinity. p. 277-299. In: G.A. Jung (ed.) Crop tolerance to subotimal land conditions. Am Soc. Agron. Spec. Publ. No. 32. Madison, WI.
- Maas, E.V. (1986):** Salt tolerance of plants. Appl. Agric. Res. 1: 12-26.
- Maxted, N.; M.A. Ccallimassia and M.D. Bennett (1991a):** Cytotaxonomic studies of Eastern Mediterranean Vicia species (Leguminosae). *Plant Syst. Evol.*, 177; 221-234.
- Maxted, N.; A.M.A. Khattab and F.A. Bisby (1991b):** The newly discovered relatives of *Vicia faba* L. do litle to resolve the engima of its origin. *Bot. Chron.*, 10: 435-465.
- McCoy, T.J. (1987):** Tissue culture evaluation of NaCl tolerannce in medicago species: cellular versus whole plant response. *Plant Cell Reports* 6: 31.

- Mehta, B.V. and R.S. Lasai (1958):** Effect of various salinity regions on growth leaf expansion and transpiration rate of been plants. Soil. Sci., 109, 26.
- Metry (1993):** Physiological genetic studies for salt tolerance on some tissues cultures wheat varieties. M.Sc. Genetics, Ain Shams Univ.
- Michelmore, I. Paran and R.V. Kesseli (1991):** Identification of markers linked to disease-resiatnace genes by bulked segregant analysis: A rapid method to detect markers in specific genomic regions by using segregation populations. Genetics, Vol 88, pp. 9828-9832.
- Misra, A.N.; M. Misra, and N. Das. (1990):** Sodium chloride salt stress effect on pearl millet callus: 1. Growth water status. Protein and proline content. (Abst 7th international. Congress on plant tissue and Cell Culture, P. 22 9 Amesterdam.
- Mohamed, A.S. (1991):** Somaclonal variation in some agronomic characters in wheat. International Symposium On Plant Biotechnology and its Contribution to Plant Development, Multiplication and Invironment. Eds. J.M. Mascherpa, and C. Moncousin. P. 247-249. Geneva, Switaerland.
- Mohmand, A.S. (1991):** Somaclonal variation in some agronomic characters in wheat. International Symposium on plant Biotechnology and its Contribution to plant Development, Multiplication and Invironment. eds. J.M. Mascherpa, and C. Moncousin. P. 247-249. Geneva, Switzerland.
- Moorginski, L.A.; K.K. Kartha (1984):** Tissue culture of Legumes for crop improvement. Plant breeding Reiews. 2, 215-264.
- Mullet, J.E.; R. Creelman; H.M. Son; E. Bell; J. Jones; D. Dewalld and M. Whitsitt (1991):** Regulation of genes by abscisic acid and jasmonic acid during plant water deficiet. Molecular Biology of Plant growth and development. Third international Congress Society for Plant Molecular Biology. Abst. 143.

- Naboors, M.W.; A. Daiels; L. Nadolny and C. Brown (1975):*
Sodium chloride tolerant lines of tobacco cells. Plant Science Letters 4: 155-159.
- Narendra, K.; Singh Avtar, K.; Handa; M. Paul Hasegawa and Ray A. Bressan (1985):* Proteins associated with adaption of cultured tobacco cells to NaCl. Plant Physiol. 79: 126-137.
- Nettevich, E.; N.S. Berkutova and L.CG. Pogorelova (1983):*
Electrophoresis method in studying interavarietal variation in wheat grain quality. Seleksiya Semenovodstvo, USSR 11:8.
- Nieman, R.H. and L.L. Poulsen (1971):* Plant growth suppression on saline media: Interactions with light. Bot. Gaz. 132: 14-19.
- Nieman, R.H. and M.C. Shannon (1977):* Screening plants for salinity tolerance. P. 359-367. In: M.J. Wright. Plant adaptation to mineral stress in problem soils. Beltsville, Maryland.
- Norlyn, J.D. (1980):* Breeding salt tolerant plants P. 58-80. In: Genetic engineering of osmoregulation. Impact on plant productivity for food chemicals and energy. Plenum Press. New York.
- Novikov, U.S. (1942):* investigation into the salt tolerance of cotton. Seiyaxi, 5 Taskeut (C.F. Stroganov, 1962).
- Orf, J.H.; N. Kaizuma and T. Hymowitz (1980):* Six soybean seed protein electrophoretic variants. seed Sci. and Tech., 8: 401.
- Passera, C. and A. Albuzio (1978):* Effect of salinity on photosynthesis and photorespiration of two wheat species (*Triticum durum* cv. PEPE 2122 and *Triticum aestivum* CV. Marzotto). Can. J. Bot. 56: 121-126.
- Pasternak, D. (1987):* Salt tolerance and crop production. A comprehensive approach. Ann. Rev. Phytopathol. 25: 271-291. Pavlova, M.K. (1983). The use of in vitro culture methods in studies on gametophytes of higher plants. Plant breeding 40 (2) 65-69, 126.

- Pavolva, M.K.* The use of in vitro culture methods in studies on gametophytes of higher plants. *Ukrains'kii Botanichnii Zhurnal* (1983): 40 (2) 65-69, 126 (Uk, en, 31 ref.) institute Botaniki im. N.G. kholodnaga, Kiev, Ukrainian, SSR.
- Payne, R.C. and T.J. Koszykowski. (1978):* Esterase isozyme difference in seed extracts among soyebean varieties. *Crop Sci.* 18: 557.
- Perrino, P.; J. Maellaro, and A. Blanco (1977):* Electrophoretic patterns in species of the genus *Vicia*. *Genetica Agraria* 31 (1/2) 121-130.
- Pogna, N.E.; F. Mellini and DAL B.A. Peruffo (1985):* The role of polyacrylamide gel electrophoresis in varietal identification and in developing new varieties of durum wheat with good spaghetti-making quality. *Monografie di Genetica Agraria* 7: 199-212.
- Przybylska, J.; S. Blixt; H. Parzysch and Z.Z. Przybylski (1982):* Isoenzyme variation in genus *Pisum*. I. Electrophoretic Patterns of several enzyme systems. *Genetica Polonica*, 23: 103.
- Rahman, M.M. and K. Kaul (1989):* Differentiation of sodium chloride tolerant cell lines of tomato (*Lycopersicon esculentum* Mill). Cv. Jet Star. *J. Plant Physiol.* 133: 710-712.
- Ramagopal, S. (1987):* Molecular biology of salinity stress: preliminary studies and perspectives. P. 111-119. In: G. Bruening, J. Harada, T. Kosuge and A. Hollaender. Tailoring genes for crop improvement: An agricultural perspective. *basic life Science* Vol. 41, Plenum, New York.
- Rahsed, M.A. (1989):* Significance of protoplast in predicting salt tolerant wheat varieties. *Egypt J. Genet. Cytol.* 18: 285-290.
- Rahed, M.A.; A. Bahieldin and A.A. Goda. (1991):* The use of protein electrophoretic patterns isozymes for characterization of diverse male sterile cytoplasm of wheat. *Annals of Agric. Sci* 36: 307.

- Rosen, A. and M. Tal (1981):** Salt tolerance in the wild relatives of the cultivated tomato: Responses of naked protoplasts isolated from leaves of *Lycopersicon esculentum* L. and *Peruvianum* to NaCl and Proline. *Z. Pflanzenphysiol.* 102: 91-94.
- Roupakias, D. G. (1985):** Callus formation and regeneration from explants of *Vicia faba* L. and *Vicia narbonesis* L. *FABIS Newsletter* No. 11, 9-11.
- Roy and Ben-Hayyim (1985):** Selection and breeding for salinity tolerance in vitro. *Plant and Soil* 89, 243-252.
- Sabrah, N.S.; A.Y. El-Metiamy (1985):** Genetic distances between local and exotic cultivars of *Vicia faba* L. based on esterase isozymes variation. *Egyptian J. Genet. and Cytol.* 14 (2): 301-307.
- SAS. (1988):** Release 6.03 Ed. Cary, NC: SAS Institute Inc. NC. USA.
- Salam (1993):** Physiological genetic studies on some Egyptian wheat varieties. M.Sc. Genetics, Ain Shams Univ.
- Salih, F.A.; M.S. Omar and H.A. Saif (1990):** Effects of sodium chloride on some cellular contents in wheat callus cultures. (Abst) 7th International Congress on Plant Tissue and Cell Culture. . 230. Amsterdam.
- Sarin, N.B.; A. Gupta; S. Prakash and B. Biswas (1991):** Isolation and characterization of a salt-tolerant line of *Arachis hypogaea* using in vitro cultures. In: International Symposium on Plant Biotechnology and its Contribution to plant development, Multiplication and improvement Eds. J.M. and c. Moncousin. P. 210-222. Geneva, Switzerland.
- Sathish, P. and M.W. Nabors (1992):** NaCl-tolerant rice plant from NaCl tolerant rice cell lines obtained through anther culture. *Advances in Gene Technology: Feeding the world in the 21st Century. Proceedings of the 1992 Miami Bio/Technology Symposium.*

- Sayed, H.I.; A.S. Mashhady and M.H. Abddul-Aziz (1983):* Screening for salinity tolerance in wheat. p. 663-668. In: Proc 6th Int. Wheat Genet. Symp. Kyoto, Japan.
- Schaeffer, G.W.; P.S. Baenziger and J. Worley (1979):* Haploid plant development from anthers and in vitro embryo culture of wheat. Crop Sciences 19: 697-702.
- Schullze, S.; J. Grunewaldt; H. Schamidt (1985):* In vitro regeneration of *Vicia faba* L. Zeitschrife fur Pflanzenzuchtung 94 (3): 255-250.
- Shannon, L.M. (1968):* Plant isoenzymes. Ann. Rev. of plant physiol. Vol. 19: 187-210.
- Shannon, M.C. (1982):* Genetics of salt tolerance. New challenges. p. 271-2281. In: A. San Pietro (ed.) Biosaline Research. A look to the future (Environmental Science Research, Vol. 23). Plenum, New York.
- Shannon, M.C. (1984):* Breeding selection and the genetics of salt tolerance In; R.C. Staples and G.A. Toenniessen (ed.) Salinity tolerance in plants. Strategies for crop improvement. Wiley, New York.
- Shannon, M.C. (1985):* Principles and strategies in breeding for higher salt tolerance. P. 227-241. In: D. Pasternak and A. San Pietro (ed.): Biosalinity in action: Bioproduction with saline water. Dordrecht: Nijhoff.
- Sheffield, V.C.; D.R. Cox; L.S. Lerman and R.M. Myers (1989):* Attachment of a 40-base-pair G+C-rich sequence (SC clump) to genomic DNA fragments by the polymerase chain reaction results in improved detection of single base changes Proc. Natl. Acad. Sci. (USA), 86: 232-236.
- Sheooran, L.S. and O.P. Gorg (1978):* Effect of salinity on the activities of RNase, DNase and protease during germination and early seedling growth of mung bean. Physiol. Plant 44 (3): 171-174.

- Singh, T.N.; D. Aspinall and L.G. Paleg (1972):* Proline accumulation and varietal adaptability to drought in barley. - A. potential metabolic measure of drought resistance. *Nature Biol.* 236, 188.
- Singh, T.N.; D. Aspinall, L.G. Paleg and S.F. Boggess (1973b):* Stress metabolism. II. Changes in proline concentration in excised plant tissues. *Aust. J. Biol. Sci.*, 26, 57.
- Singh, A. (1982):* Effect of water stress salinity and alkalinity on seed germination of *Sesbania aegyptiaca*. *Geobios (Jodhpur)*, 9 (2): 26-65.
- Singh, K.N. and R.S. Rana (1985):* Genetic variability and character association in wheat varieties grown on sodic soil. *Indian J. Agric. Sci.* 55: 723-726.
- Singh, Handa, Hasegawa and Bressan (1985):* Protein associated with adaptation of tobacco cells to NaCl. *Plant Physiol.* Vol. 79; 126-137.
- Snell, F.D. and C.T. Snell (1954):* Colourimetric method of analysis Vol. N. New York. Van Nostrand.
- Srivastava, J.P. and S. Jana (1984):* Screening wheat and barley germ plasm for salt tolerance. P. 273-283. In: R.C. Staples and G.A. Toenniessen (ed.): *Salinity tolerance in plants: Strategies for crop improvement*. Wiley, New York.
- Stankevich, A.K. (1982):* Systematic position of some sections of the genus *Vicia*. *L. Trudy po Prikladnoi Botanike, Genetike i Selekcii* 72 (1): 21-27. (C.F. Faba Bean Abstr. 5 (4): 344, 1985).
- Staples, R.C. and C.A. Tenniessen, ed. (1984):* Salinity tolerance in plants: Strategies for crop improvement. 443 P. Wiley, New York.
- Stavarek, G.J. and D.W. Rains (1984):* Cell culture technique for selection and physiological studies of salt tolerance. P. 321-334. In *Salinity tolerance in plants: Strategies for crop improvement*. John Wiley and Sons, New York.

- Stegemann, H.; A.M.R. Afify and K.R.F. Hussein (1985):* Cultivar identification of dates (*Phoenix dactylifera*) by protein patterns. 2nd International Symposium of Biochemical Approaches to identification of cultivars. Braunschweig, West Germany, p. 44.
- Steward, F.C.; R.F. Lyndon and J.T. Barber (1965):* Acrylamide gel electrophoresis of soluble plant proteins A study on pea seedlings in relation to development. *Am. J. bot* 52: 155-164.
- Strogonov, B.P.; V.V. Kabanov, N.I. Shevjakova, L.P. Lapina, E.I. Komizerko, B.A. Popov; R.K. Dostanova and L.S. Prykhodko (1973):* Structure and function of plant cells in saline habitats. New trends in the study of salt tolerance. Israel Program for Scientific Translation, Jerusalem. Wiley, New York.
- Studier, F.W. (1973):* Analysis of bacteriophage T2 early RNAs and protein of slab gels. *J. Mol. Biol.*, 79: 237.
- Tal, M.; h. Heikin and K. Dehan (1978):* Salt tolerance in the wild relatives of the cultivated tomato: Responses of callus tissue of *Lycopersicon esculentum*, *L. peruvianum* and *Solanum pennellii* to high Salinity. *Z. Pflanzphysiol.* 86; 231-240.
- Tal, M. and J.W. Watts (1979):* Plant growth conditions and yield of viable protoplasts isolated from leaves of *Lycopersicon esculentum* and *L. Peruvianum*. *Z. pflanzphysiol.* 92: 207-214.
- Tanksley, S.D.; R. Bernatsky; N.L. Lapitan and J.P. Prince (1988):* Conservation of gene repetitive but not gene order in pepper and tomato. *Proc. Natl. Acad. Sci. (USA)*, 85: 6419-6423.
- Tanksley, S.d.; N.D. Young; A.H. Paterson and M.W. Bonnierbale (1989):* RFLP mapping in plant breeding: new tools for an old science. *Biotechnology*, 7: 257-264.
- Thompson, J.F.; C.R. Stewart and C.J. Morris (1966):* Changes in amino acid content of excised leaves during incubation. I. Effect of water content of leaves and atmospheric oxygen level. *Plant Physiol.*, 41, 1578.

- Utsumi, S.; Z. Yekouama and T. Mori (1980):* Comparative studies of subunit composition of legumins from various cultivars of *Vicia faba* L. seeds. *Agric. Biol. Chem.* 44: 595.
- Vajrabhaya, M.; T. thanapaisal and T. Vajrabhaya (1989):* Development of salt tolerant lines of KDML and LPT rice cultivars through tissue culture. *Plant Cell Reports* 8: 411-414.
- Van de Ven; W.T.G.; R. Waugh; N. Duncan; G. Ramsay; N. Dow and W. Powell (1991):* Development of a genetic linkage map in *Vicia faba* using molecular and biochemical techniques. *Aspects Appl. Biol.*, 27: 49-54.
- Van de Ven; W.T.G.; N. Duncan; G. Ramsay; M. Phillipps; W. Powell and R. Wough (1993):* Taxonomic relationships between *V. faba* and its relatives based on nuclear and mitochondrial RFLPs and PCR analysis. *Theor. Appl. Genet* 86: 71-80.
- Waisel (1972):* Biology of Halophytes. P. 178-184. Academic Press, New York. ISBN 0-12-730950-4.
- Webster, P.L. (1980):* Stress protein synthesis in pea root meristem cells. *Plant Sci. Lett.* 20: 141-145.
- Welsh, J. and M. McClelland (1990):* Fingerprinting genomes using PCR with arbitrary primers. *Nucleic acids Res.*, 18: 7213-7218.
- Wiinikov, L. (1990):* Gene expression in salt tolerant alfalfa cell cultures and the salt tolerant plant regenerated from tissue cultures. (Abst.) 7th international Congress on Plant Tissue and cell Culture. P. 144. Amsterdam.
- Williams, J.G.K.; A.R.; Kubelick; K.J. Livak; J.A. Rafalski and S.V. Tingey (1990):* DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.*, 18: 6531-6535.
- Williams, J.G.K.; J.A. Rafalski and S.V. Tingey (1992):* Genetic analysis using RAPD markers. *Methods Enzymol.* (in press).

- Wilson, J.R.; K.P. Hoydock and M.T. Robins (1970):* Response to salinity in glycine 5-changes in the chemical composition of three Australian species of *G. weightii* (*G. javanica*) over a range of salinity stresses. *Aust. J. Exp. Agric. Anim. Husbandry* 10: 158-165.
- Winter, E. (1982):* Salt tolerance of *Trifolium alexandrinum* L. III: Effect of salt on ultra-structure of phloem and xylem transfer cells in petioles and leaves. *Aust. J. Plant Physiol.* 9: 230-250.
- Winter, E. and J. Preston (1982):* Salt tolerance of *Trifolium alexandrinum* L. IV: Ion measurements by X-ray microanalysis in unfixed, frozen hydrated leaf cells at various stages of salt treatment. *Aust. J. Plant Physiol.* 9: 251-259.
- Wolff, G. (1980):* Investigation on the relation within the family papilionaceae on the basis of electrophoretic banding patterns. *Theoretical and Applied Genetics* 57: 225-232.
- Yamamoto, K.; O. Moritani and A. Ando (1982):* Karyotypic and isozymatic polymorphism in species of section *Faba* (Genus *Vicia*). *tech. Bull. agric. Kagawa Univ.*, 34:1.
- Yang, X. and C.F. Quiros (1993):* Identification and classification of celery cultivars with RAPD markers. *Theor. Appl. Genet.*, 86: 205-212.