

CHAPTER 11

REFERENCES

REFERENCES

Adams, T. and P. Dalzell, 1994. Artisanal fishing . Paper presented at the workshop on Marine / Coastal Biodiversity in the tropical Island Pacific Region, 2-9 November 1994, East West Centre, Honolulu, Hawaii. *South Pacific Commission, Noumea, New Caledonia*. 19p.

Alagaraja, K., 1984. Simple methods for estimation of parameters for assessing exploited fish stocks. *Indian J. Fish.*, 31 : 177 - 208.

Albaret, J. J., 1976. La reproduction de l'albacore (*Thunnus albacares* Bonnaterre, 1788) dans le Golfe de Guinee. These de 3^e cycle. *Universite de Paris*.

Aldonov, K. V. and A. D. Druzhinin, 1979. Some data on scavenger (family Lethrinidae) from the Gulf of Aden region. *Voprosy Ikhtologii*, 8 (4) : 527 - 535.

Al - Hussaini, A. H., 1947a. The anatomy and histology of the alimentary tract of the plankton - feeder, *Atherina forskali*. *Rupp. J. Morph.*, 80 : 251 - 286.

_____, 1947b. The feeding habits and the morphology of the alimentary tract of some teleosts living in the neighbourhood of the marine biological station, Ghardaqa, Red Sea. *Pub. Mar. Biol. Sta., Ghardaqa, Red Sea*, 5 : 1 - 61.

_____, 1949. On the functional morphology of the alimentary tract of some fish in relation to differences in their feeding habits. Anatomy and histology. *Quart. J. Micros. Sci.*, 90 : 109 - 139.

Allen, K. R., 1935. The food and migration of the perch, *Perca fluviatilis* in Windermere. *J. Anim. Ecol.*, 4 : 264 - 273.

_____, 1940. Studies on the biology of the early stages of salmon (*Salmo salar*). II . Feeding habits. *J. of Anim. Ecol.*, 9 : 47 - 76.

_____, 1942. Comparison of bottom faunas as sources of available fish food. *Trans. Am. Fish. Soc.*, 71: 275 - 283.

_____, 1951. The Horokiwi stream : a study of a trout population. *Fish. Bull. N.Z.*, 10 : 1 -238.

_____, 1966. A method of fitting growth curves of the von Bertalanffy type to observed data. *J. Fish. Res. Bd. Can.*, 23 : 18 - 23.

Arawomo, G. A. O., 1976. Food and feeding of three *Citharinus* species in lake Kainji, Nigeria. *J. Fish. Biol.*, 9 : 3 - 10.

Baddar, M. K., 1987. A preliminary study of the population dynamics of sheiry, the starry pigface bream *Lethrinus nebulosus*. *Kuwait Bull. Mar. Sci.*, 9 : 215 - 220.

Bagenal, T. B., 1971. The inter - relation of the size of fish eggs, the data of spawning and the production cycle. *J. Fish. Biol.*, 3 : 20 - 219.

Ball, J. N., 1961. On the food of the brown trout of Llyn Tegid. *Proc. Zool. Soc. Lond.*, 137 : 599 - 622.

Bautil, B. and C. R. Samboo, 1988. Preliminary stock assessment for the mahsena emperor (*Lethrinus mahsena*) on the Nazareth bank of Mauritius; workshop on the assessment of the fishery resources in the

south west Indian Ocean; Albion (Mauritius). *FAO Fish. Proj., RAF*
/ 79 / 065 : 260 - 267.

Beckman, W. C., 1948. The length - weight relationship, factor for conversions between standard and total length and coefficient of condition for seven Michigan fishes. *Trans. Am. Fish. Soc.*, 13 : 312 - 340.

Berhaut, J. A., 1973. Biologie des stades juveniles de teleosteens Mugilidae, *Mugil auratus* (Risso, 1810), *Mugil capito* (Cuv., 1829) et *Mugil saliens* (Risso, 1810). *Aquac.*, 2 : 251 - 266.

Bertalanffy, L. von, 1934. Untersuchungen über die Gesetzmäßigkeiten des wachstums. 1. Allgemeine Grundlagen der theorie. *Roux' Arch. Entwicklungsmech. Org.*, 131 : 613 - 653.

Bertalanffy, L. von, 1938. A quantitative theory of organic growth (Inquiries on growth Laws. 2). *Hum. Biol.*, 10 : 181 - 213.

Bertrand, J., 1986. Donneés concernant la reproduction de *Lethrinus mahsena* (Forsskal, 1775) sur les bancs de Saya de Malha (Ocean Indien). *Cybium*, 10 (1) : 15 - 29.

_____, 1988. Selectivity of hooks in the handline fishery of the Saya de Malha banks (Indian Ocean). *Fish. Res.*, 6 (3) : 249 - 255.

Beverton, R. J. H. and S. J. Holt, 1957. On the dynamics of exploited fish population. *U. K. Min. Agr. Fish. Food, Ish. Invest*, 19 : 533 p.

_____, 1959. A review of the life spans and mortality rates of fish in nature and their relation to growth and other physiological characteristics. In : Ciba Foundation colloquia on ageing (ed. G. E. W. Wolstenholme and M. O'Connor). Vol. 5 : 142 - 180.

Bhattacharya, C. G., 1967. A simple method of resolution of a distribution into Gaussian components. *Biometrics*, 23 : 115 - 135.

Broadway, J. E. and P. B. Moyle, 1978. Aspects of the ecology of the prickly sculpin *Cottus asper*, a persistent native species in Clear Lake Calif. *Envir. Biol. Fishes*, 3 : 337 - 343.

Brody, S., 1927. Growth rates. *Univ. Missouri Agric. Exp. Sta. Bull.*, 97.

_____, 1945. Bioenergetics and growth. *Reinhold, New York*: 1023p.

Bulkley, R. V.; V. L. Spykermann and L. E. Inman, 1976. Food of the pelagic young of walleyes and five species in Clear Lake Iowa. *Trans. Am. Fish. Soc.*, 105 : 77 - 83.

Caddy, J. F., 1980. Surplus production models. In : Selected lectures from the CIDA / FAO / CEECAF seminar on fishery resource evaluation. Casablanca, Morocco, 6 - 24 March, 1978 : *Rome, FAO Canada Funds - in - Trust, FAO / -TF / MJ 180 (c) Suppl.* : 29 - 55.

Carpenter, K. E., 1996. Lethrinidae. Emperor snappers. In : FAO identification Guide for Fishery Purposes (eds. K. E. Carpenter and V. Niem). *The Western Central Pacific*. 1 - 55.

Carpenter, K. E. and G. R. Allen, 1989. FAO species catalogue. Vol. 9. Emperor fishes and large - eye breams of the world (family Lethrinidae). *FAO Fish. Syn. No. 125, Vol. 9* : 118 p.

Cassie, R. M., 1954. Some uses of probability paper in the analysis of size frequency distributions. *Aust. J. Mar. Freshw. Res.*, 5 : 513 - 522.

Chapman, D. G., 1960. Statistical problems in dynamics of exploited fisheries populations. Proc. 4th. *Berkeley Symp. Math. Statist. Probl.*, 4: 153 - 168.

Chapman, D. G. and D. S. Robson, 1960. The analysis of a catch curve. *Biometrics*, 16 (3) : 354 - 368.

Charbonnier, D. and S. Garcia (eds.), 1983. Third technical consultation on stock assessment in the Balearic and Gulf of Lions statistical divisions. *FAO Fish. Rep.*, 305 : 110p.

Christy, F. T. Jr. and A. Scott, 1965. The common wealth in ocean fisheries. Problems of growth and economic allocation. *Baltimore, Johns Hopkins Press for accuources for the future Inc.* : 281 p.

Craig, J. F., 1978. A study of the food and feeding of perch (*Perca fluviatilis*) in Windermere. *Freshw. Biol.*, 8 : 59 - 68.

Crisp, D. T. ; R. H. Mann and J. C. Mc Cormack, 1978. The effects of impoundment and regulation upon the stomach contents of fish at Cow Green, Upper Teesdale. *J. Fish. Biol.*, 12 : 287 - 301.

Dalzell, P. ; S. Sharma and G. Nath, 1992. Estimation of exploitation rates in a multispecies emperor (pisces : Lethrinidae) fishery in Fiji, based on length frequency data. In : papers on fisheries science from the Pacific Islands (ed. P. Dalzel). Vol. 1. Inshore Fish. Res. Proj. Tech. Doc. No.1. *South Pacific Commission Noumea, New Caledonia.* 43 - 50.

Dineen, C. F., 1951. A comparative study of the food habits of *Cottus bairdii* and associated species of Salmonidae. *Am. Midl. Nat.*, 46 : 640 - 645.

Dipper, F. A., C. A. Bridges and A. Menz, 1977. Age, growth and feeding in the ballan wrasse *Labrus bergylta* Ascanius 1767. *J. Fish. Biol.*, 11 : 105 - 120.

Djama, Th., 1989. Maximum sustainable yield : the necessary evil. *Fishbyte*, 7 (2) : 2p.

Dunn, D. R., 1954. The feeding habits of some of the fishes and some members of the bottom fauna of Llyn Tegid (Bala Lake), Merionethshire. *J. Anim. Ecol.*, 23 : 224 - 233.

Eapen, P. K. and K. W. Krimly, 1982. Fisheries of Saudi Arabia. Kingdom of Saudi Arabia : Ministry of Agriculture and Water, *Agriculture Research Department report* : 93p.

Edwards, R. R., 1983. The Taiwanese pair trawler fishery in tropical Australian waters. *Fish. Res.*, 2 : 47 - 60.

Edwards, R. R. ; C. A. Bakhader and S. Shaher, 1985. Growth, mortality, age composition and fisheries yields of fish from the Gulf of Aden. *J. Fish. Biol.*, 27 : 13 - 21.

Edwards, R. R. ; A. Gaddaf and S. Shaher, 1987. The demersal fish stocks and the biometrics of fish on the PDR Yemen shelf of the Gulf of Aden. *Final report to UNESCO (Marine Science Division)*. 128 p.

Edwards, R. R. and S. Shaher, 1991. The biometrics of marine fishes from the Gulf of Aden. *Fishbyte*, 9 (2) : 27 - 29.

Efford, I. E. and K. Tsumura, 1973. A comparison of the food of salamanders and fish in Marion Lake, British Columbia. *Trans. Am. Fish. Soc.*, 102 : 33 - 47.

El - Dossary, N., 1987. Biological studies on some fishes of family Lethrinidae. *M. Sc. Thesis, Fac. Sci. High Presidency for Girls Education, El - Dammam, Saudi Arabia.* 431p.

El - Gammal, F. I., 1988. On the growth and demographic structure of *Lethrinus varigatus* Cuv. & Val. (1830) from the Red Sea. *Bull. Inst. Oceanog. & Fish., ARE*, 14 : 1-12.

Elliott, J. M. 1967. The food of trout (*Salmo trutta*) in a Dartmoor stream. *J. appl. Ecol.*, 4 : 59 - 71.

Fabens, A. J., 1965. Properties and fitting of the von Bertalanffy growth curve. *Growth*, 29 : 265 - 289.

FAO , 1981a. Demersal resources of the Gulf of Oman. Regional fishery survey and development project. *Rome, FAO, ID : DP/RAB/71/278/10* : 122p.

____, **1981b.** Pelagic resources of the Gulf and Gulf of Oman. Regional fishery survey and development project. *Rome, FAO, ID : DP/RAB/71/278/11* : 144p.

____, **1985.** Statistical basebook on African countries affected by emergencies. *Rome, FAO, Stat. Div., W/R 4527* : 100p.

____, **1989a.** Report of the sixth session of the working party on assessment of marine fishery resources. St. George's, Grenada, 15 - 19 May, 1989. *FAO Fish. Rep.*, 431 : *FP/L/R* 431p.

FAO, 1989b. Report of the seventh session of the CECAF sub - committee on management of resources within the limits of National Jurisdiction. *FAO Fish. Rep.*, 406 : 83p.

_____, 1992. Review of the state of world fisheries resources. Part 1. The marine resources. *FAO, Fish. Circ.*, C710 (Rev.8) : 325 - 335.

_____, 1993a. FAO year book. Fishery statistics : catches and landings, Vol. 72 (1991) Food and Agricultural Organization of the United Nations. Rome, Italy. *FAO Fish. Series*, 40 : 654p.

_____, 1993b. Fishery fleet statistics. FAO fishing information, Data and Statistics Service, 1970, 1975, 1980 - 1989. *FAO, Rome. Bull. Fish. Stat.*, 30 : 344p.

_____, 1994 . Review of the state of world marine fishery resources. *FAO Fish. Tech. Pap.*, 335 : 136p.

_____, 1995 . World fishery production 1950 - 1993. *FAO, Suppl. Rome.* 44p.

Fishelson, L., 1980. Marine reserves along the Sinai Peninsula (northern Red Sea). *Helgol. Wiss. Meeresunter*, 33 : 624 - 640.

Ford, E., 1933. An account of the herring investigations conducted at Plymouth during the years from 1924 to 1933. *J. Mar. Biol. Assoc. U. K.*, 19 : 305 - 384.

Forsskal, P., 1775. Descriptiones animalium avium, amphibiorum, piscium, insectorum, vermium, quae in itinere orientali observavit. Post mortem auctoris edidit Carsten Niebuhr. *Molleri, Hauniae.* 20 + xxx iv + 164p., 43 pls.

Fouda, M. M. and G. V. Hermosa, Jr, 1993. A checklist of Oman fishes. *Sultan Qaboos University Press, Sultanate of Oman*. 42p.

Fox, W. W. Jr., 1970. An exponential surplus - yield model for optimizing exploited fish populations. *Trans. Am. Fish. Soc.*, 99 : 80 - 88.

Frost, W. E., 1939. The food consumed by the brown trout (*Salmo trutta* L.) in acid and alkaline waters. *Proc. Roy. Irish Acad.*, 45 : 193 - 206.

_____, 1943. The natural history of the minnow *Phoxinus phoxinus*. *J. Anim. Ecol.*, 12 : 139 - 162.

_____, 1946. On the food relationships of the fish in windermere. *Biol. Jaarb.*, 13 : 216 - 231.

_____, 1954. The food of the pike, *Esox lucius*, L., in windermere. *J. Anim. Ecol.*, 23 : 339 - 360.

Fry, F. E. J., 1949. Statistics of a lake trout fishery . *Biometrics*, 5 : 27 - 67.

Fulton, T. W., 1911. The sovereignty of the sea . *Edinburgh and London*.

_____, 1920. Report on the marking experiments of plaice, made by S/S Goldseeker, in the years 1910 - 1913. *Sci. Invest. Fish. Scotl.* (1919),1.

Garcia, S. and L. Le Reste, 1981. Life cycles, dynamics, exploitation and management of coastal penaeid shrimp stocks. *FAO Fish. Tech. Pap.*, 203 : 215 p.

Gibbons, J. R. H. and J. H. Gee, 1972. Ecological segregation between longnose and blacknose dace (*Rhinichthys*) in the Mink River, Manitoba. *J. Fish. Res. Bd. Can.*, 29 : 1245 - 1252.

Glenn, C. L. and E. J. Ward, 1968. Wet weight as a method of measuring stomach contents of walleyes *Stizostedion vitreum vitreum*. *J. Fish. Res. Bd. Can.*, 25 : 1505 - 1507.

Gompertz, B., 1825. On the nature of the function expressive of the law of human mortality. *Phil. Trans. R. Soc.*, 36 : 513 - 585.

Gregor, J., 1969. An algorithm for the decomposition of a distribution into Gaussian components. *Biometrics*, 25 (1) : 79 - 93.

Gueno, Y. and J. C. Guen, 1968. Determination de l'age et de la croissance de *Sardinella eba* (Val.) dans la region de Pointe - Noire. *Cah. ORSTOM (Ser. Oceanogr.)*, 6 (2) : 69 - 82.

Gulland, J. A., 1965. Estimation of mortality rates. Annex to Arctic fisheries working group report *ICED C. M. Doc.*, 3 (mimeo).

_____, 1969. Manual of methods for fish stock assessment. Part 1. Fish population analysis. *FAO Man. Fish. Sci.*, 4 : 154p.

_____, 1974. Guidelines for fishery management. Rome, FAO, Indian Ocean Programme, *IOFC/DEV/74/36* : 84p.

Gulland, J. A., 1977. Goals and objectives of fishery management. *FAO Fish. Tech. Pap.*, 166 : 14p.

_____, 1978. Fishery management : new strategies for new conditions. *Trans. Am. Fish. Soc.*, 107 : 1 - 11.

_____, 1983. Fish stock assessment : a manual of basic methods. Chichester, U. K., Wiley Interscience, *FAO/ Wiley series on food and agriculture*, Vol. 1 : 223p.

Gulland, J. A. and S. L. Holt, 1959. Estimation of growth parameters for data at unequal time intervals. *J. Cons. Perm. Int. Explor. Mer*, 25 (1) : 47 - 49.

Haram, O. J. and J. W. Jones, 1971. Some observations on the food of the gwyniad *Coregonus clupeoides pennantii* Valenciennes of Llyn Tegid (Lake Bala) North Wales. *J. Fish. Biol.*, 3 : 287 - 295.

Harding, J. P., 1949. The use of probability paper for the graphical analysis of polymodal frequency distributions. *J. Mar. Biol. Ass. U. K.*, 28 : 141 - 153.

Hashim, M. T. and A. A. Shakour, 1981. Age determination and growth studies of *Lethrinus mahsena* and *L. xanthochilus*. Jeddah, *J. Mar. Res.*, 1 : 11 - 20.

Hasselblad, V., 1966. Estimation of parameters for a mixture of normal distributions. *Technometrics*, 8 : 431 - 444.

Healey, M. C., 1986. Multiattribute analysis and the concept of optimum yield. *Can. J. Fish. Aquat. Sci.*, 41 : 1393 - 1406.

Hickling, C. F. and E. Rutenberg, 1936. The ovary as an indicator of the spawning period in fisheries. *J. Mar. Biol. Assoc.*, 21 : 311 - 317.

Higham, J. R. A. and W. R. Nicholson, 1964. Sexual maturation and spawning of Atlantic Menhaden. *Fish. Bull.*, 63 (2) : 255 -271.

Hile, R., 1936. Age and growth of the cisco *Leucichthys artedi* (Le Sueur) in the lakes of the north eastern highlands, Wisconsin. *Bull. U. S. Bur. Fish.*, 48 : 211 - 317.

Hjort, J., 1910. Report on herring investigation . *Publ. Circons. Cons. Perm. Explor. Mer.* 53p.

_____, 1914. Fluctuations in the great fisheries of northern Europe, viewed in the light of biological research. *Rapp. P.-V. Reun. Cons. Perm. Int. Explor. Mer.*, 20 : 1 - 228.

Hoffbauer, C., 1898. Die Altersbestimmung des karpfens an seiner Schuppe. *Allgemeine Fischereizeitung*, 23 (19).

Holt, S. J., 1960. A preliminary comparative study of the growth, maturity and mortality of sardines. In : proceedings of the world scientific meeting on the biology of sardines and related species (eds. H. Rosa, Jr. and G. Murphy). *Rome, FAO, Vol.*, 2 : 553 - 561.

_____, 1962. The application of comparative population studies to fishery biology - an exploration. In : the exploitation of natural animal populations, British Ecological Society Symposium No. 2 (eds. E. D. Le Cren and M. W. Holdgate). *Oxford , Blackwell Scientific publications* : 51 - 69.

Holt, S. J. , 1965. A note on the relationship between mortality rate and the duration of life in an exploited fish population. *ICNAF Res. Bull.*, 2 : 73 - 75.

Holt, S. J. and L. M. Talbot, 1978. New principles for the conservation of wild living resources. *Wildl. Monogr.*, 59 : 33p.

Hunt, B. P. and N. F. Carbine, 1951. Food of young pike and associated fishes in Peterson's Ditches, Houghton Lake, Michigan. *Trans. Am. Fish. Soc.*, 80 : 67 - 73.

Hunt, P. C. and J. W. Jones, 1972. The food of brown trout in Llyn Alaw, Anglesey, North Wales. *J. Fish. Biol.*, 4 : 333 - 352.

Huang, T. S. and J. L. Yen, 1985. Embryonic and larval development of the scavenger, *Lethrinus nebulosus*. *Bull. Taiwan Fish. Res. Inst.*, 38 : 147 - 155.

Hynes, H. B. N., 1950. The food of freshwater sticklebacks (*Gasterosteus aculeatus*, and *Pygosteus pungitius*) with a review of methods used in studies of the food of fishes. *J. Anim. Ecol.*, 19 : 36 - 58.

Hyslop, E. J., 1980. Stomach contents analysis : a review of methods and their application. *J. Fish. Biol.*, 17 : 411 - 429.

Ibrahim, M. A.; K. A. El-Bary and J. A. Al-Khayat, 1988. Stock assessment and management of *Lethrinus nebulosus* (family Lethrinidae) of Qatar waters, Arabian Gulf. *Bull. Nat. Inst. Oceanog. & Fish., AREE*, 14 (2) : 99 - 111.

Ikusemiju, K. and C. I. O. Olaniyan, 1977. The food and feeding habits of the cat fishes, *Chrysichthys walkeri* (Gunther), *Chrysichthys filamentosus* (Boulenger) and *Chrysichthys nigrodigitatus* (Lacepede) in the Lekki Lagoon, Nigeria. *J. Fish. Biol.*, 10 : 105 - 112.

Jones, A., 1973. The ecology of young turbot *Scophthalmus maximus* (L.) at Borth, Cardiganshire, Wales. *J. Fish. Biol.*, 5 : 367 - 383.

Jones, R., 1976. Mesh regulation in the demersal of the south China Sea area. Manila, south China Sea Fisheries Development and coordinating Programme, SCB/76/WP/34 : 75p.

_____, 1984. Assessing the effects of changes in exploitation pattern using length composition data (with notes on VPA and cohort analysis) *FAO Fish. Tech. Pap.*, (256) : 118p.

Jones, R. and N. P. Van Zalinge, 1981. Estimates of mortality rate and population size for shrimp in Kuwait waters. *Kuwait Bull. Mar. Sci.*, 2 : 273 - 288.

Jude, D. J., 1971. Food and feeding habits of gizzard shad in pool 19, Mississippi River. *Trans. Am. Fish. Soc.*, 102 : 378 - 383.

Katzir, G. and E. Schechtman, 1986. Interactions during feeding among certain coral reef fishes in Elat. *Mar. Biol.*, 91 : 441 - 447.

Kedidi, S. M., 1984. Stock assessment for the red spot emperor *Lethrinus lentjan* from areas adjacent to Suakin and Muhammad Qol (Sudan) survey conducted during 1982 - 1984. *FAO, Rome, Italy. RAB/83/023/07* : 27p.

Kedidi, S. M., 1984. The Red Sea reef associated fishery of the Sudan. Catches, efforts and catches per fishing effort survey conducted during 1982 - 1984. *FAO, RAB / 83 / 023 / 06* : 18p.

Kedidi, S. M. ; T. Abu Shusha and K. Allam, 1984. Biology and stock assessment of the red spot emperor *Lethrinus lentjan*, from waters adjacent to Tuwwal, Saudi Arabia. Proj. Dev. Fish. Red Sea and Gulf of Aden. *RAB/81/002/15, FAO* : 21p.

Kedidi, S. M. and M. Bouhlef, 1985. Stock assessment for the long face emperor *Lethrinus miniatus* from the Djiboutian waters, Red Sea - Gulf of Aden. *FAO, Rome. RAB/83/023/12* : 25p.

Kennedy, M. and P. Fitzmaurice, 1972. Some aspects of the biology of gudgeon *Gobio gobio* (L.) in Irish waters. *J. Fish. Biol.*, 4 : 425 - 440.

Kesteven, G. L., 1960. Manual of field methods in fisheries biology. *FAO, Man. Fish. Sci. No. 1* : 152p.

Komrakov, O. Ye. and V. P. Meshcheryakov, 1972. Some features of the biology of *Lethrinus nebulosus* from the area of the north - west coast of Australia. In : *Issledovaniya po biologii ryb i promyslovoy okeanogr.* (studies of the biology of fish and commercial oceanography) No. 7. Published by the Pacific Institute for sea Fisheries and Oceanography.

Kruger, F., 1973. Zur mathematik des tierischen wachstums. II. Vergleich einiger wachstums funktionen. *Helgolander wiss. Meeresunters.*, 25 : 509 - 550.

Kulbicki, M. ; G. Mou Tham ; P. Thollot and L. Wantiez, 1993. Length - weight relationships of fish from the lagoon of New - Caledonia. *NADA, The ICLARM J.* 16 (2 - 3) : 26 - 29.

Kuo, C. L., 1988. The study of fishery biology on porgies *Lethrinus nebulosus*, Forsskal in waters of Australia. Sino - Jap. Colloq. *Assess. Manag. Mar. Fish. Res.* : 125 - 131.

Kuo, C. L. and S. S. Lee, 1986. Age and growth of common porgy, *Lethrinus nebulosus* (Forsskal) in shelf waters off north west Australia. *J. Fish. Soc. Taiwan*, 3 : 39 - 57.

Lagler, K. F., 1956. Freshwater fishery biology. Wm. C. Brown Co., Dubuque, Iowa : 421p.

Lagler, K. F. ; C. E. Bardach ; R. R. Miller and D. R. Passion, 1977. Ichthyology. John Wiley and Sons. New York.

Larimore, W. R., 1957. Ecological life history of the warmouth (Centrarchidae). *Bull. Ill. St. Nat. Hist. Surv.*, 27 : 81 - 82.

Larkin, P. A., 1977. An epitaph for the concept of maximum sustainable yield. *Trans. Am. Fish. Soc.*, 106 (1) : 1 - 11.

Lea, E., 1910. On the methods used in the herring investigations. *Publ. Circumst., Cons. Int. Explor. Mer.*, 53 : 7 - 174.

Lebeau, A. and J. C. Cueff, 1975. Biologie et peche du capitaine *Lethrinus enigmaticus* (Smith, 1959) du band de Saya de Malha (Ocean Indien). *Rev. Trav. Inst. Peches marit.*, 39 (4) : 415 - 442.

Lebeau, A and J. C. Cueff, 1976. Biologie et peche des lethrinide's sur les haut fonds de Saya de Malha. *Trav.et documents ORSTOM*,47: 333 - 348.

Le Cren, E. D., 1951. The length - weight relationship and seasonal cycle in gonad weight and condition in the perch. *J. Anim. Ecol.*,20 : 201 - 219.

Le Drew, B. R. and J. M. Green, 1975. Biology of the radiated shanny *Ulvaria subbifureata* storer in Newfoundland (pisces : Stichaeidae). *J. Fish. Biol.*,7 : 485 - 495.

Lee, J. U., 1990. Effects of changes in fishing effort on yield of Kuwait's commercial fish stocks. *Bull. Korean Fish. Soc.*, Vol. 23 (6) : 475 - 483.

Lee, J. U. and F. Al-Baz, 1989. Assessment of fish stocks exploited by fish traps in the Arabian Gulf area. *Asian Fish. Sci.*, 2 : 213 - 231.

Lee, R. M., 1920. A review of the methods of age and growth determination in fishes by means of scales. *Fish. Invest. Min. Agr. Fish.*, Ser. 2, 4 (2) : 1 - 32.

Lee, S. C., 1986. Fishes of the genus *Lethrinus* (percoidi : Lethrinidae) of Taiwan. *Bull. Inst. Zool., Acad. Sinica*, 25 (1) : 79 - 89.

Loubens, G., 1978. Biologie de quelques especes de poissons du lagon neo - Caledonien I. Determination de l'age (otolithometrie). *Cah. ORSTOM* , Ser. Oceanogr., 16 (3-4) : 263 - 283.

Loubens, G., 1980a. Biologie de quelques especes de poissons du lagon neo - Caledonien. II. Sexualite et reproduction. *Cah. Indo-Pac.*, 2 (1) : 41 - 72.

_____, 1980b. Biologie de quelques especes de poissons du lagon neo-Caledonien. III. Croissance. *Cah. Indo-Pac.*, 2 (2) : 101 - 153.

Man, H. S. H. and I. J. Hodgkiss, 1977. Studies on the ichthyo-fauna in Plover Cove Reservoir, Hong Kong : feeding and food relations. *J. Fish. Biol.*, 11 : 1 - 13.

Mann, R. H. K., 1973. Observations on the age, growth, reproduction and food of the roach, *Rutilus rutilus* (L.), in two rivers in southern England. *J. Fish. Biol.*, 5 : 707 - 736.

Marten, G. G., 1978. Calculating mortality rates and optimum yields from length samples. *J. Fish. Res. Bd. Can.*, 35 (2) : 197 - 201.

Martin, W. R., 1949. The mechanics of environmental control of body form in fishes. *Univ. Toronto Stud., Biol. Ser. No. 58. Pub. Ontario Fish. Res. Lab.*, 70 : 1 - 72.

Mathews, C. P. and M. Samuel, 1991. Management and research strategies in Kuwait's trawl fishery. International council for the exploration of the sea. *ICE Mar. Sci. Symp.*, 193 : 330 - 340.

_____, 1991. Growth, mortality and length - weight parameters for some Kuwaiti fish and shrimp. *Fishbyte*, 9 (2) : 30 - 33.

May, R. M. ; J. R. Beddington ; C. W. Clark ; S. J. Holt and R. M. Laws, 1979. Management of multispecies fisheries. *Science*, 205 : 267 - 277.

Mc Comish, T. S., 1966. Food habits of bigmouth and smallmouth buffalo in Lewis and Clarke Lake and the Missouri River. *Trans. Am. Fish. Soc.*, 96 : 70 - 74.

Mc Pherson, G. R. et al., 1985. Great Barrier Reef demersal fish research . South Pacific Commission, 7th Regional Technical Meeting on Fisheries, Noumea, New - Caledonia, August 1985. *SPC / Fisheries* 17 / WP 23.

Mitchell, C. L., 1979. Bioeconomics of commercial fisheries management. *J. Fish. Res. Bd. Can.*, 36 : 699 - 704.

Mohsin, A. M. ; M. A. Ambak and M. N. Abdul Salam, 1993. Malay, English and scientific names of the fishes of Malaysia. *Fuc. Fish. Mar. Sci.*, Universiti Pertanian Malaysia, Selanger Darul Ehsan, Malaysia. *Occasional publication* No. 11.

Monastyrsky, G. N., 1926. Methods of determining growth of fish from scale measurments. In : Instructions for growth rate studies with the Monastyrsky calculating device. *Vses. Nauchno - Issled. Inst. Morsk. Rybn. Khoz. i Okeanogr. (VNIRO)*, Moscow.

Moore, J. W. and I. A. Moore, 1975a. The basis of food selection in flounders in the Severn Estuary. *J. Fish. Biol.*, 9 : 139 - 156.

_____, 1975b. The basis of food selection in some estuarine fishes. Eels, whiting, spart and stickleback. *J. Fish. Biol.*, 9 : 375 - 390.

Morales - Nin, B., 1988. Age determination in a tropical fish, *Lethrinus nebulosus* (Forsskal, 1775) (teleostei : Lethrinidae) by means of otolith interpretation. *Inv. Pesq.*, 52 (2) : 237 - 244.

_____, 1989. Growth determination of tropical marine fishes by means of otolith interpretation and length analysis. *Aquat. Living Resour.*, 22 : 241 - 253.

Morgan, C. R., 1985. Status of the shrimp and fish resources of the Gulf. *FAO Fish. Cir.*, 792 : 49p.

Munro, J. L., 1976. Aspects of the biology and ecology of Caribbean reef fishes : Mullidae (goat fishes). *J. Fish. Biol.*, 9 : 79 - 97.

Neill, R. M., 1938. The food and feeding of the brown trout (*Salmo trutta*, L.) in relation to the organic environment. *Trans. R. Soc. Edinb.*, 59 : 481 - 520.

Nikolsky, G. V., 1963. The ecology of fishes. London, Academic Press : 352p.

Nzioka, R. M., 1979. Observations on the spawning seasons of east African reef fishes. *J. Fish. Biol.*, 14 : 329 - 342.

Pauly, D., 1980. A selection of simple methods for the assessment of tropical fish stocks. *FAO Fish. Circ.*, 729 : 54p.

_____, 1980a. A new methodology for rapidly acquiring basic information on tropical fish stocks : growth, mortality and stock recruitment relationships. In : Stock assessment for tropical small-scale fisheries (eds. S. Saila and P. Roedel). Univ. Rhode Island. Int. Center Mar. Res. Dev., 198p.

Pauly,D., 1980b. On the interrelationships between natural mortality growth parameters and mean environmental temperature in 175 fish stocks. *J. Cons. CDEM*, 39 (2) : 175 - 192.

_____, **1983a.** Length converted catch curves : a powerfull tool for fisheries research in the tropics. Part 1. *Fishbyte*, 1 (2) : 9 - 13.

_____, **1983.** Some simple methods for the assessment of tropical fish stocks. *FAO, Fish, Tech. Pap.*, 234 : 52p.

_____, **1984.** Fish population dynamics in tropical waters : a manual for use with programmable calculators. *ICLARM Stud. Rev.*, (8) : 325p.

_____, **1984a.** Length - converted catch curves. A powerful tool for fisheries research in the tropics. (part II) . *ICLARM Fishbyte*, 2 (1) : 17 - 19.

_____, **1984b.** Length - converted catch curves. A powerful tool for fisheries research in the tropics. (III : conclusion) . *ICLARM Fishbyte*, 2 (3) : 9 - 10.

_____, **1989a.** Biology and management of tropical marine fisheries. *Res. Manag. Optimaz.*, 6 : 253 - 271.

_____, **1989b.** The editor replies. *Fishbyte*, 7 (2) : 3 - 4.

Pauly, D. and G. I. Murphy (eds.), 1982. Theory and management of tropical fisheries. *ICLARM Conf. Proc.*, 9 : 360p.

Pedley, R. B. and J. W. Jones, 1978. The comparative feeding behaviour of brown trout *Salmo trutta* (L.) and Atlantic salmon *Salmo salar* (L.) in Llyn Dwythwch, Wales. *J. Fish. Biol.*, 12 : 239 - 256.

Pella, J. J. and P. K. Tomlinson, 1969. A generalized stock production model. *Bull. Int.-Am. Trop. Tuna Comm.*, 13 : 419 - 496.

Pemberton, R., 1976. Sea trout in North Argyll Sea lochs. II. Diet. *J. Fish. Biol.*, 9 : 195 - 208.

Petersen, C. G. J., 1892. Fiskenes biologiske forhold i Holbaek Fjord, 1890 - 1891. *Beret. Danm. Biol. St.*, 1890 (1) 1 : 121 - 183.

Peterson, I. and S. J. Wroblewski, 1984. Mortality rate of fishes in the pelagic ecosystem. *Can. J. Fish. Aquat. Sci.*, 41 : 1117 - 1120.

Pitcher, J. T. and P. J. B. Hart, 1982. Fisheries ecology. Croom Helm, London, 414p.

Pitt, T. K., 1970. Distribution, abundance and spawning of yellow tail flounder, *Limanda ferriginea* in the new found land area of the north - west Atlantic. *J. Fish. Res. Bd. Can.*, 12 : 2261 - 2271.

Pope, J. G., 1979. Stock assessment in multispecies fisheries with special reference to the trawl fisheries in the Gulf of Thailand. Manila. South China Sea Development and coordinating Programme, SCS/DEV/79/-19 : 106p.

_____, 1980. Assessment of multispecies resources. In : Selected lectures from the CIDA/FAO/CECAF seminar on fishery resource evaluation. Casablanca, Morocco, 6 - 24 March 1978. Rome, FAO, Canada Funds-in-Trust, FAO/TF/INF 180 (c) Suppl. : 93 - 137.

Rafail, S. Z., 1971. A new growth model for fishes and the estimation of optimum age of fish populations. *Mar. Biol.*, 14 : 338 - 340.

_____, 1972. Fitting a parabola to growth data of fishes and some applications to fisheries. *Mar. Biol.*, 15 : 255 - 264.

_____, 1973. A simple and precise method for fitting a von Bertalanffy growth curve. *Mar. Biol.*, 19 : 354 - 358.

Ralston, S., 1982. Influence of hook size in the Hawaiian deep - sea handline fishery. *Can. J. Fish. Aquat. Sci.*, 39 : 1297 - 1302.

Randall, J. E., 1983. Red Sea reef fishes. *Immel publishing, London*. 192p.

Reibisch, J., 1899. Über die Eizahl bei *Pleuronectes platessa* und die Untersbestimmung dieser form aus den otolithen. *Wiss. Meeresuntersuchungen (kiel) N. F.*, 4 : 233 - 248.

Rice, A. L., 1962. The food of the sea scorpion *Acanthocottus bubalis* in Manx waters. *Proc. Zool. Soc. Lond.*, 138 : 295 - 303.

Richards, F. J., 1959. A flexible growth function for empirical use. *J. Exp. Bot.*, 10 : 290 - 300.

Ricker, W. E., 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Bd. Can.*, 191 : 382p.

_____, 1979. Growth rates and models. In : Fish physiology. Bioenergetics and growth (eds. W. S. Hoar, D. J. Randall and J. R. Brett). *London, Academic Press* : 677 - 744.

Rikhter, V. A. and V. N. Efanov, 1976. On one of the approaches to estimation of natural mortality of fish populations. *ICNAF Res. Doc.*, 76/88/8 : 12p.

Robotham, P. W. J., 1977. Feeding habits and diet in two populations of spined loach *Cobitis taenia* (L.). *Freshw. Biol.*, 7 : 469 - 477.

Rounsefell, G. A. and W. H. Everhart, 1953. Fishery science : Its methods and applications. John Wiley and Sons, New York, N. Y. 444p.

Roussow, G., 1957. Some considerations on sturgeon spawning periodicity in northern Quebec. *J. Fish. Res. Bd. Can.*, 14 : 553 - 572.

Saila, S. B. and P. Roedel (eds.), 1980. Stock assessment for tropical small-scale fisheries. Proceedings of an international workshop held 19 - 21 September (1979), at the University of Rhode Island, Kingston R. I., International Center for Marine Resources Development, Univ. Rhode Island. 198p.

Salem, S. A., 1971. Biological studies on family Lethrinidae from the Red Sea. M. Sc. thesis, Cairo University, A.R.E.

_____, 1976. On the biology of two *Lethrinus* species (from the Red Sea). Thesis of Ph. D. Zoology Department, Faculty of Science, Cairo University, A.R.E.

Samboo, R., 1982. The Mauritius banks fishery. *Swio Fish. Bull. Peches Oiso*, 2 : 9 - 12.

Samuel, M. ; C. P. Mathews and A. S. Bawazeer, 1987. Age and validation of age from otoliths for warm water fishes from the Arabian Gulf. In : Age and growth of fish (eds. R. C. Summerfelt and G. E. Hall). Iowa State Univ. Press, Ames., Iowa. 253 - 265.

Sanders, M. J., 1989. An assessment of exploitation levels in the banks hand line fishery and a management plan to improve the fishery performance. *FAO/UNDP RAF/87/008/WP/52/89/E* : 71 - 85.

Sanders, M. J. and S. M. Kedidi, 1984. Catches, fishing effort, catches per fishing effort and fishing locations for the Gulf of Suez and Egyptian Red Sea fishery for reef associated fish during 1979 to 1982. *UNDP/FAO. RAB/83/002/02* : 65p.

Sanders, M. J. ; S. M. Kedidi and M. R. Hegazy, 1984. Stock assessment for the spangled emperor *Lethrinus nebulosus* caught by small scale fishermen along the Egyptian Red Sea coast. *FAO, Rome. RAB/83/023/01* : 41p.

Sanders, M. J. and G. R. Morgan, 1989. Review of the fisheries resources of the Red Sea and Gulf of Aden. *FAO, Fish. Tech. Rep.*, 304 : 138p.

Sano, M. ; M. Shimizu and Y. Nose, 1984. Food habits of teleostean reef fishes in Okinawa Island, southern Japan. *Univ. Mus., Univ. Tokyo Bull.*, 25 : 1 - 128.

Sato, T., 1971. A revision of the Japanese sparoid fishes of the genus *Lethrinus*. *J. Fac. Sci., Univ. Tokyo. Sec.4*, 12 (2) : 117 - 144.

Sato, T. , 1978. A synopsis of the sparoid fish genus *Lethrinus* with the description of a new species. *Univ. Mus., Univ. Tokyo Bull.*, 15 : 1 - 70, 35 Figs., 12 pls.

_____, 1984. Lethrinidae. See Fischer, W. & G. Bianchi (eds.).

_____, 1986. A systematic review of the sparoid fishes of the subfamily Monotaxinae. In : Indo - Pacific fish biology (eds. T. Uyeno et al.). *Proc. Sec. Intl. Conf. Indo - Pac. Fishes, Tokyo. Ichthyological Society of Japan* : 602 - 612.

Sauvonsaari, J., 1971. Biology of the stone loach (*Nemacheilus barbatulus*, L.) in the Lakes Paijanne and Palkanevesi, S. Finland. *Ann. Zool.fenn.*, 8 : 187 - 193.

Saville, A., 1977 (ed.). Survey methods of appraising fisheries resources. *FAO Fish. Tech. Pap.*, 171 : 76p.

Schaefer, M. B., 1954. Some aspects of the dynamics of populations important to the management of commercial marine fisheries. *Bull. inter - Am. Trop. Tuna Comm.*, 1 (2) : 26 - 56.

_____, 1957. A study of the dynamics of the fishery for yellowfin tuna in the eastern tropical Pacific Ocean. *Bull. inter - Am. Trop. Tuna Comm.*, 2 (6) : 247 - 285.

Sharma, S. P., 1990. Growth and mortality of spangled emperor *Lethrinus nebulosus* in Fiji. *Fishbyte* 8 (2) : 3 - 5.

Sheriff, C. W. M., 1922. Herring investigations. Report on the mathematical analysis of random samples of herrings. *Fish. Bd. Scotland, Sci. Invest.* (1922) : 1 - 25.

Sikora, W. B.; R. W. Heard and M. D. Dahlberg, 1972. The occurrence and food habits of two species of hake *Urophycis regius* and *Urophycis floridanus* in Georgia estuaries. *Trans. Am. Fish. Soc.*, 101 : 515 - 516.

Simpson, A. C., 1951. The fecundity of the plaice. *Fish. Invest. Ser.* II, 17 (5) : 1 - 27.

Sissenwine, M. P., 1978. Is MSY an adequate foundation for optimum yield? *Fisheries*, 3 (6) : 22 - 24 and 37 - 42.

Sivasubramaniam, K., 1981. Biological information collected during the demersal resources survey. In : Demersal resources of the Gulf and the Gulf of Oman. Regional fishery survey and development project. *FAO, Rome. FI : DP/RAB/71/278/10* : 57 - 88.

Sivasubramaniam, K. and M. A. Ibrahim, 1981. Fisheries of Qatar : past, present and future possibilities. *MS Univ. of Qatar, Sci. Res. Cen.*, 131p.

Smith, J. L. B., 1955. New species and new records of fishes from Mozambique. Part 1. *Mem. Mus. de Castro*, 3 : 3 - 27.

_____, 1957. Deep - line fishing in northern Mozambique with the description of a new pentapodid fish. *Ann. Mag. Nat. Hist.*, (12) 10 : 121 - 124, 1 pls.

_____, 1959. Fishes of the family Lethrinidae from the western Indian Ocean. *Ichth. Bull. Rhodes Univ.*, 17 : 285 - 295, pls 20 - 25.

Smith, M. M., 1986. Lethrinidae. In : Smith's Sea Fishes (eds. M. Smith and P. C. Heemstra). *J. L. B. Smith Institute of Ichthyology, Grahamstown*, 1047 p., 144 pls.

Smyly, W. J. P., 1952. Observations on the food of fry of perch (*Perca fluviatilis*, Linn.) in Windermere. *Proc. Zool. Soc. Lond.*, 122 : 407 - 416.

Sparre, P. ; E. Ursin and S. C. Venema, 1989. Introduction to tropical fish stock assessment. Part 1. Manual. *FAO Fish. Tech. Pap.*, 306 (1) : 337p.

Ssentongo, G. M. and P. A. Larkin, 1973. Some simple methods of estimating mortality rates of exploited fish populations. *J. Fish. Res. Bd. Can.*, 30 : 695 - 698.

Starostka, V. J. and R. L. Applegate, 1970. Food selectivity of bigmouth buffalo *Ictiobus cyprinellus* in Lake Poinsett, south Dakota. *Trans. Am. Fish. Soc.*, 99 : 571 - 573.

Stearns, S. C. and R. E. Crandall, 1984. Plasticity for age and size at sexual maturity. A life - history response to unavoidable stress. In : reproduction : strategies and tactics (eds. G.W. Potts and R. J. Wootton). *Academic press, U. K.* : 13 - 33.

Stickney, R. R., 1976. Food habits of Georgia estuary fishes. II. *Symphurus plagiatus* (*Pleuronectiformes* : *Cynoglossidae*). *Trans. Am. Fish. Soc.*, 105 : 202 - 207.

Straskraba, M. ; J. Chiar ; S. Frank and V. Hruska, 1966. Contribution to the problem of food competition among the sculpin, minnow and brown trout. *J. Anim. Ecol.*, 35 : 303 - 311.

Swynnerton, G. H. and E. B. Worthington, 1940. Notes on the food of fish in Haweswater (Westmorland). *J. Anim. Ecol.*, 9 : 183 - 187.

Tanaka, S., 1956. A method of analysing the polymodal frequency distribution and its application to the length distribution of porgy, *Taius tumifrons* (T. & S.). *Bull. Tokai Reg. Fish. Res. Lab.*, 14 : 1 - 12.

_____, 1960. Studies on the dynamics and the management of fish populations. *Bull. Tokai Reg. Fish. Res. Lab.*, 28 : 1 - 200.

_____, 1962. A method of analysing a polymodal frequency distribution and its application to the length distribution of the progey *Taius tumiforms* (T. & S.). *J. Fish. Res. Bd. Can.*, 19 (6) : 1143 - 1159.

Tawada, S., 1983. Studies on the seedling production of Lethrinous fish *Lethrinus nebulosus*, Forsskal. The natal spawning in using large - scale out-door tanks. *Aquaculture*, 31 (2) : 60 - 66.

Taylor, C. C., 1960. Temperature, growth and mortality - the Pacific Cockle. *J. Cons. CDEM*, 26 : 117 - 124.

Thompson, W. F. and F. H. Bell, 1934. Biological statistics of the Pacific halibut fishery. 2. Effect of changes in intensity upon total yield and yield per unit of gear. *Rep. Int. Fish. (Pacific Halibut) Comm.*, (8): 49p.

Thresher, R. E., 1984. Reproduction in reef fishes. *IFH Publications*. Neptune City, N. J. : 399p.

Tomlinson, P. K. and N. J. Abramson, 1961. Fitting a von Bertalanffy growth curve by the least squares. *Fish. Bull. Calif.*, 116 : 1 - 69.

Toor, H. S., 1964. Biology and fishery of the pigface bream, *Lethrinus lentjan*. I. Maturation and spawning. *Indian J. Fish.*, 11 (A), 11 (2) : 559 - 596.

_____, 1966. On the occurrence of *Lethrinus mahsenoides* in the Indian waters with a note on *Lethrinus sanguineus*. *Proc. Indian Sci. Congr.*, 53 (3) : 378p.

_____, 1968. Biology and fishery of the pigface bream *Lethrinus lentjan*, Lacepede from Indian waters. III. Age and growth. *Ind. J. Fish. Sect.*, (A), 11 (2) : 597 - 620.

_____, 1973. Taxonomic position of *Lethrinus mahsenoides* and *Lethrinus sanguineus*. *Res. Bull. Panjan Univ. Sci.*, 22 (3-4) : 375 - 378.

Ursin, E., 1967. A mathematical model of some aspects of fish growth, respiration and mortality. *J. Fish. Res. Bd. Can.*, 24 : 2355 - 2453.

Van Oosten, J., 1923. The white fishes *Coregonus clupeaformis*. A study of the scale of white fishes of known ages. *Zoologica*, II, 17 : 380 - 412.

_____, 1929. Life history of the lake herring (*Leucichthys artedi*) Le Sueur, of lake Huron as revealed by its scales, with a critiques of the scales method. *Bull. U. S. Bur. Fish.*, Vol. 44 (1928): 265 - 448.

Verhulst, P. F., 1845. Recherches mathematiques sur la Loi d'accroissement de la population. *Nouv. Mem. Acad. Roy. Sc. et Belles Lettres Bruxelles*, 18 : 1 - 38.

Walford, L. A., 1946. A new graphic method of describing the growth of animals. *Biol. Bull. Mar. Biol. Lab., Woods Hole*, 90 (2) : 141 - 147.

Walker, M. H., 1975. Aspects of the biology of emperor fishes , family Lethrinidae , in north Queensland Barrier Reef waters. Unpubl., Ph. D. Diss., Dept. Zoology. James Cook Univ. N. Queensl.

_____, 1978. Food and feeding habits of *Lethrinus chrysostomus* Richardson (pisces : Perciformes) and other lethrinid in the Great Barrier Reef. *Aust. J. Mar. Freshw. Res.*, 29 (5) : 623 - 630.

Wassef, E. A., 1991. Comparative growth studies on *Lethrinus lentjan*, Lacepede, 1802 and *L. mahsena*, Forsskal, 1775 (pisces, Lethrinidae) in the Red Sea. *Fish. Res.*, 11 : 75 - 92.

Wetherall, J. A., 1986. A new method for estimating growth and mortality parameters from length - frequency data. *Fishbyte*, 4 (1) : 12 - 14.

Windell, J. T. and S. H. Bowen, 1978. Methods for study of fish diets based on analysis of stomach contents. In : Methods for the assessment of fish production in fresh waters. 3 rd edn. (ed. T. Bagnel). Oxford : Blackwell Scientific publications. 219 - 226.

Winsor, C. P., 1932. The Gompertz curve as a growth curve. *Proc. Nat. Acad. Sci.*, 18 (1) : 1 - 8.

Wolfert, P. R. and T. J. Miller, 1978. Age, growth and food of northern pike in Eastern Lake Ontario. *Trans. Am. Fish. Soc.*, 107 : 696 - 702.

Yamamoto, K., 1956a. Studies on the formation of fish eggs. I. Annual cycle in the development of the ovarian eggs in the flounder, *Liopsetta obscura*. *J. Fac. Sci. Hokkaido Univ., Ser. VI*, 12 : 362 - 373.

_____, 1956b. Studies on the formation of fish eggs. II. Changes in the nucleus of the oocyte of *Liopsetta obscura*, with special reference to the activity of the nucleolus. *J. Fac. Sci. Hokkaido Univ., Ser. VI*, 12 : 375 - 390.

Yaroslavtseva, L. D., 1966. Technical and chemical properties of some fishes of the Indian Ocean. *Rybn. Kh. - Vo. No. 1*.

Yesaki, M., 1981. Review of exploratory trawling investigations and commercial fishing ventures in the Gulfs. In : Demersal resources of the Gulf and Gulf of Oman. Regional fishery survey and development project. Rome, FAO, ID : DP/RAB/71/278/10 : 89 - 115.

Yoshida, M. O., 1966. Skipjack tuna spawning in the Marquesas Islands and Tuamatu Archipelago. *Fish. Bull.*, Vol. 65 (2) : 479 - 488.

Young, P. C. and R. B. Martin, 1982. Evidence for protogynous hermaphroditism in some lethrinid fishes. *J. Fish. Biol.*, 21 : 475 - 484.