

INFERENCES

REFERENCES

Alirzaev, G. M. (1986)

Practical tests of dimilin in the control of mosquito larvae in Baku . Meditsinskay a parazitologiya ; Parazitarny Bolezin , (1) : 32 - 35 .

Amalraj, D. D. ; Doss, P.S.B. ; Mariappan, T. & Arunachalam, N. (1995) .

Toxicity of some commercially available insecticides to field strain of *Culex sitiens* wiedemann, 1828 (Diptera : Culicidae) Trop. Biomedicine, 12 (2) : 97 - 99 .

Amin, A. M. & Hemingway, J. (1989) .

Preliminary investigation of the mechanisms of DDT and pyrethroid resistance in *Culex quinquefasciatus* say (Diptera : Culicidae) from Saudi Arabia .

----- **& Peiris, H.T.R. (1990)**

Detection and selection of organophosphate and carbamate resistance in *Culex quinquefasciatus* from Saudi Arabia, 4 (3) : 269 - 273 .

Apperson, C.S. & Georghiou, G.P. (1975) .

Mechanisms of resistance to organophosphorous insecticides in *Culex tarsalis* . J. Econ. Entomol., 68 (2) 153 - 157 .

Ascher, K.R.S. & Nemny, N. E. (1974) .

The ovicidal effect of PH 6040 1-(4-chlorophenyl)-3-(2,6- difluorobenzoyl) Urea in *Spodoptera littoralis* (Boisd) . Phytoparasitica, 2 (2) : 131 - 133 .

Azmi, M.A. ; Nagvi, S.N.H. & Akhtar, K. (1993) .

Comparative study of toxicity of DDT, malathion and solfac against various strains of *Culex fatigans* by WHO method .

Bakr, R.F.A. ; Abo Gabal, N.M. & Hussein, M.A. (1989) .

Insect growth regulators : 1 . Biological activity of some IGR's against the susceptible and resistant strains of *Culex pipiens* larvae . II . pattern of cross resistance to IGR's in carbaryl-resistant strain . J. Egypt. Soc. parasit., 19 (2) : 589 - 596 .

Beard , C. B. ; Kloter , K. O. ; Carroll , M. K. ; Lisa , J. M. & Trapido , H. (1985) .

Response of domestic and peridomestic strains of *Aedes Aegypti* (Diptera : Culicidae) in New Orleans , Louisiana , USA , to organophosphate , organochlorine and pyrethroid insecticides . J. Med. Entomol., 22 (3) : 276 - 280 .

Betzios, B.C. (1977) .

Development of resistance to dieldrin in a laboratory colony of the mosquito *Culex pipiens molestus* forsk . (Lab. insecticides of publ. health) import . Dep. phytopharm., Inst. phytopathol . Benaki , 11 (4) : 284 - 301 .

Bilbie, I. ; Guirca, I. ; Velehorsi, N. ; Ceianu, G. ; Tacu, V. & Dragon, D. (1984) .

The larvicidal effect on culicidae of microencapsulated malathion laboratory and field trials . Archives Roumaines de pathologie Experimentale et de Microbiologie, 43 (2) : 185 - 191 .

Brown, A.W.A. & Tadano, T. (1965) .

Studies on insecticide resistance in *Culex pipiens fatigans* .
World Health Organization Info. Cric. Insecticide resistance .
Insect behavior and vector genetics 52 : 4.5 .

----- ; Lewallen, L. L. & Gillies, P. A. (1963) .

Organophosphorous resistance in *Aedes nigromaculi* in California . Mosq. News, 23 (4) : 341 - 345 .

Buei, K. ; Ito, S. ; Yamoda, T. ; Gamo, S. & Kato, M. (1975) .

The effect of a juvenile hormone mimic , methoprene, against mosquito larvae . Jap. J. sanit. Zool., 26 (213) : 105 - 111 .

Busvine, J.R. (1970) .

A critical review of the technique for testing insecticides .
Comm. Agric. Bull. England, pp. 345 .

Casida, J. E. ; Engel, J. ; Essac, E.G. ; Kamienski, F. X. & Kawatsuke, S. (1966) .

Metabolism of methylene-C¹⁴-dioxyphenyl compounds in relation to their synergistic action . Science , 153 , 1130 .

Colwell, A. E. & Schaefer, C. H. (1981) .

Effects of the insect growth regulator bay sir 8514 on Diptera and non-target aquatic organisms . Can. Entomol., 117 : 185 - 191 .

Crow, J.F. (1954) .

Genetics of insect resistance to chemicals . Ann. Rev. Entomol., 2 : 226 - 246 .

Curtis, C. F. & pasteur, N. (1981) .

Organophosphate resistance in vector populations of the complex of *Culex pipiens* L. (Diptera : Culicidae) . Bull. Ent. Res., 71 : 153 - 161 .

Dame, D. A. ; Lowe, R.E., ; Wichterman, G. J. ; Cameron, A. L. ; Baldwin, K.F. & Miller, T. W. (1976) .

Laboratory and field assessment of insect growth regulators for mosquito control . Mosq. News., 36 (4) : 462 - 472 .

Dash, A. P. & Hazra, R. K. (1995) .

A small scale field experiment on comparative efficacy of three mosquito larvicides in a filarial endemic locality of Drissa, India . Trop. Biomed., 12 (1) : 69 - 72 .

Davis, A. N. & Gahan, J. B. (1961) .

New insecticides for the control of salt-marsh mosquitoes . The Florida Entomologist, 44 (1) : 11 - 14 .

Dorta, D. M. ; Vasuki, V. & Rajavel, A. (1993) .

Evaluation of organophosphorous and synthetic pyrethroid insecticides against six vector mosquito species . Revista de Saude Publica, 27 (6) : 391 - 397 .

Dubois, K. P. (1961) .

Potential of the toxicity of organophosphorous compounds . Advances pest Cont. Res., 4 : 117 - 151 .

Salama, E. M. (1988) .

Interaction between chemical and biological larvicides used in mosquito control . M Sc. thesis, Faculty of Science, Zagazig University, Benha Branch, pp. 135 .

----- (1994) .

Effect of surfactants on the histology, histochemistry and ultrastructure of *Culex pipiens* resistant to insecticides . Ph. D. thesis, Faculty of Science, Zagazig University, Benha Branch, pp. 106 .

El-Sebae, A. H. ; Metcalf, R. L. & Fukuto, T. R. (1964) .

Carbamate insecticides synergism by organothiocyanates . J. Econ. Entomol., 57 : 478 - 482 .

Failloux, A. B. ; Raymond, M. & Pasteur, N. (1994) .

Insecticide susceptibility in mosquitoes (Diptera : Culicidae) from French Polynesia . J. Med. Entomol., 31 (5) : 639 - 644

Farghal, A. L. & Temerak, S. A. (1981) .

Effect of the juvenile hormone analogue altosid on some culicine mosquitoes and their associated insects under field and laboratory conditions . Zeitschrift fur Angewandte Entomol .. 92 (5) : 505 - 510 .

Gaaboub, L. A. ; El-Gayar, F. M. & Abdel-Gawaad, A. A. (1973)

Comparative studies on the sensitivity of *Culex pipiens fatigans* wiled mosquito larvae and microcrustacean adults *Daphnia magna*, strains as microbioassay test organisms for screening certain soil insecticides applied to cotton cultivations in Egypt .

Ganguly, S. S. ; Dutta, G. K. K. ; Dutta, P. K. & Gupta, K. K. D. (1994) .

Evaluation of susceptibility status of *Culex quinquefasciatus* larvae to few organophosphorous insecticides based on logistic regression analysis . Indian J. of Public Health, 38 (1) : 8 - 13 .

Georghiou, G. P. (1962) .

Carbamate insecticides : the cross-resistance spectra of four carbamate resistant strains of housefly after protracted selection pressure . J. Econ. Entomol ., 55 : 494 - 497 .

----- **& Calman J. R. (1969) .**

Results of fenitrothion selection of *Culex pipiens fatigans* wield. and *Anopheles albimanus* wield . Bull. WHO, 40 : 97 - 101 .

----- ; Metcalf, R. L. & Gidden, F. E. (1966) .

Carbamate-resistance in mosquitoes, selection of *Culex pipiens fatigans* wiedemann (*C. quinquefasciatus* say) for resistance to baygon . Bull. WHO, 35 : 691 - 708 .

----- ; Ariaratnam, V. ; Pasternak, M. E. & Lin, C. S. (1975) .

Organophosphorous multiresistance of *Culex pipiens quinquefasciatus* in California . J. Econ. Entomol., 68 (4) : 461 - 467 .

Gillies, P. A. ; Womeldorf, D. J. & White, K. E. (1967)

Cross-tolerance of California *Aedes nigromaculis* Larvae to EPN, abate and dursban . Proc. Calif. Mosq. contr. Ass. 36 : 85.

Gordon, H. T. & El-Defrawi, M. E. (1960) .

Analog-synergism of several carbamate insecticides . J. Econ. Entomol., 53 : 1004 - 1009 .

Hamed, M. S. ; Ramzi, A. & El-Said, S. (1983) .

Susceptibility status of mosquitoes in Egypt to commonly used insecticides . J. Egypt . Publ. Hlth. Assoc. Vol. Lv III ; No. 1, 2 : 160 - 167 .

Hamon, J. (1953) .

Apparition a la Reunion D'une resistance au DDT chez *Culex fatigans* wiedman ; principal vector de la filariose a *wucheria bancrofti* . Bull. Soc. Path. Exot., 64 , 454 .

- Hatakashi, M. ; Ohsumi, T. ; Kasida, H. ; Itoyg, M. & Nakayama, I. (1986) .
Mosquito larvicidal activity of juvenile hormone active oxime-
ether compounds . J. Jap. Sani. Zool., 37 (2) 99 - 104 .
- Hoogstraal, H. ; Meegan, J. H. ; Khalil, G. M. & Adham, F.K. (1979)
The Rift valley fever epizootic in Egypt 1977-1978 : 2 .
Ecological and entomological studies . Trans. R. Soc. Trop.
Med. Thyg., 73 : 624-629 .
- Hoskins, W.M. & Gordon, H. I. (1956) .
Arthropod resistance to chemicals . Ann. Rev. Ent. 1 : 89 - 122
- Hossain, M. I. ; Ameen, M.U. & Ahmed, A. K.M. (1995) .
Efficacy of two pyrethroid insecticides against *Culex quinque-*
fasciatus say larvae in Dhaka City. Bangladesh J. zoolo., 23
(2) : 187 - 192 .
- Ittycheriah, P. I. ; Quraishi, M. S. & Marks, E. P. (1974) .
Effects of ecdysones, Juvenile hormone analogs, and 6- oxo-
octanoic acid on the development of the mosquito *Culex tarsalis*
(Diptera : Culicidae) . Can. Entomol., 106 : 79 - 85 .
- Jakob, W. L. & Schoof, H. F. (1972) .
Mosquito larvicide studies With MON 585, a juvenile hormone
mimic . Mosq. News, 32 (1) : 6 - 10 .
- Kamal, A. Z. (1991) .

Laboratory evaluation of IGR_s, insecticides and bacterial agents against *Culex pipiens* . M. Sc. thesis, Faculty of science, Ain Shams University .

Kamel, O. M. ; Mahdi, A. H. ; Merk , W. & Beckmann, K. (1972) .

Ultra low volume aerial spraying of iodofenphos against mosquitoes over rice fields and villages in the Arab Republic of Egypt in 1971 . Mosq. News, 32 (4) : 514 - 519 .

Kenawy, M. A. ; Soliman, S. A. ; Kaschef, A. H. & schanbaky, N. M. (1985) .

Effect of organophosphorous insecticides on cholinesterase activity in mosquitoes . J. Egypt. Soc. parasit., 15 (1) : 201 - 212 .

Kottkamp, W. B. & Meisch, M. V. (1985) .

Efficacy of bay sir 5814, an insect growth regulators against *Posrophora columbiae* and *Anopheles quadrimaculatus* in small plot and field trials. L. Am. Mosq. control Ass., 1 (1) : 17 - 19 .

Lale, N. E. S. (1991) .

The biological effects of three essential oils on *Callosobruchus maculatus* (Coleoptera : Bruchidae) . J. Afric. Zool., 105 (5) : 357 - 362 .

Lari, F. A. (1962) .

Estimation of susceptibility of mosquito larvae to insecticides in Karachi (Pakistan) . Pakist. J. Hlth., Lahore, 12 : 1 - 14 .

Losada, E. O. ; Lazcano, Z. B. ; Coto, M. M. R. & Pantoja, C. D. (1991) .

Variability and resistance to organophosphorous and carbamate insecticides and its relationship to the activity of esterase enzymes in *Culex quinquefasciatus* say 1823 (Diptera : Culicidae) . Revista Cubana de Medicina Tropical, 43 (3) : 171 - 174 .

Lyarskii, P. P. ; Dremova, G. M. ; Zubova, M. A. ; Baidarovtseva & Tsetlin, V. M. (1983) .

Insecticidal properties of permethrin and permethrin-based agents . J. Hyg. Epidemiol . Microbiol. Immunol., 27 (2) : 173 - 178 .

Mansour, N. A. ; Eldefrawi, M. E. ; Tappozada, A. & Zeid, M. (1966)

Toxicological studies on the Egyptian cotton leaf worm, *Prodenia litura* VI. Potentiation and antagonism of organophosphorous and carbamate insecticides . J. Econ. Entomol., 59 : 307 - 311 .

Marchi, A. & Addis, G. (1990) .

Resistance to organophosphorous insecticides in Sardinian populations of *Culex pipiens* (Diptera : culicidae) XVI Congresso Nazionale della societa italiana di parassitologia, Cagliari, S. Margherita di pula, Parassitologia . 32 ; supplemento, 177 - 178 .

- Meegan, J. M. ; Khalil, G. M. ; Hoogstraal, H. & Adham, F. K. (1980) .
Experimental transmission and field isolation studies implicating *Culex pipiens* as a vector of Rift Valley fever virus in Egypt . Am. J. Trop. Med. Hyg., 29 : 1405 - 1410 .
- Metcalf, K. L. (1955) .
Physiological basis for insect resistance to insecticides .
physiol. Rev., 35 : 197 - 232 .
- Mohamed, M. A. (1986) .
Susceptibility and resistance of some culicine and Anopheline mosquitoes to some insecticides in Sharquia Governorate . M. Sc. Thesis Faculty of Science, Ain Shams Univ.
- Mosna, E. (1948) .
Culex pipiens (autogenicus) DDT- resistance loro controllo con octaklore esachorocicl oesano. Riv. Parasit., 9 : 19 - 25 R. C.
- Mouchet, J. (1964) .
Resistance of *Culex fatigans* to organophosphorous insecticides, WHO / Vector control 82 .
- Moustafa, F. I. & Georghiou, G. P. (1989) .
Extension of resistance to adults as a results of selection in mosquito larvae with temphos and the cross-resistance to other organophosphorous insecticides . International pest control, 31 (3) : 61 - 65 .

Mulder, R. & Gijswijt, M. J. (1973) .

The laboratory evaluation of two promising new insecticides which interfere with cuticle deposition. *Pestic. Sci.* 4 : 737 - 745 .

Mulla, M. S. & Darwazeh, H. A. (1976) .

Evaluation of aliphatic amine concentrates against immature mosquitoes. *Mosq. News*, 36 (2) : 160 - 165 .

----- & ----- (1979) .

New insect growth regulators against flood and stagnant water mosquitoes-effects on nontarget organisms . *Mosq. News*, 39 : 746 - 755 .

----- ; ----- & Norland, R. L. (1974) .

Insect growth regulators ; evaluation procedures and activity against mosquitoes . *J. Econ. Entomol.*, 67 (3) : 210 - 212 .

Novak, D. (1972) .

Granular Abate for control of mosquitoes larvae . *Pathol. Exp. Microboil.*, 31 (4) : 579 - 582 .

Okpara, C. C. & Iwuala, M. O. E. (1993) .

Observations on the comparative toxicities of abate, malathion and baygon against non-target immature stages of mosquitoes, *Aedes aegypti* and *Culex pipiens quinquefasciatus* (Diptera : Culicidae) *Indian J. Entomol.*, 55 (4) : 428 - 432 .

Pener, H. (1983) .

Organophosphorous multiresistance in *Culex pipiens molestus* in Israel . Rivista di parassitologia, 43 (3) : 409 - 414 .

Plapp, F. W. ; Bigley , W. S. ; Darrow, D. I. & Eddy , G. W. (1963) .

Synergism of malathion against resistant houseflies and mosquitoes . Ibid., 56 : 643 - 649 .

Post, L. C. & Vicent, W. R. (1973) .

A new insecticide inhibits chitin synthesis . Naturwissenschaften, 60 : 431 - 432 .

Pridantseva, E. A. ; Kuprianova, E. S. ; Aksenova, A. S. & Vorotnikov, L. M. (1976) .

Trials of altosid SR-10, a preparation with juvenile activity, against the mosquito *Culex pipiens molestus* (Fork, 1775) breeding in basement premises . Meditsinskaya parazitologiya i parazitarnye Bolenzni, 48 (1) : 31 - 34 .

Priester, M. T. & Goerghious, P. G. (1978) .

Induction of high resistance to permethrin in *Culex pipiens quinquefasciatus* . J. Econ. Entomol., 71 (2) : 197 - 200 .

----- & ----- (1980) .

Cross-resistance spectrum in pyrethroid-resistant *Culex pipiens quinquefasciatus* . Pestic. Soc., II (6) : 617 - 624 .

Pushpalatha, N. & Vijayan, V. A. (1994) .

Adult population differences in response to six insecticides in *Culex fuscocephala* theobald . South East Asian J. Trop. Med. & Publ. Hlth., 25 (3) : 532 - 536 .

Raghavendra, K. ; Subbarao, S. K. ; Vasantha, k ; Pillai, M. K. K. & Sharma, V. P. (1992) .

Differential selection of malathion resistance in *Anopheles culicifacies* A and B (Diptera : Culicidae) in Haryana state , India . J. Med. Entomol., 29 (2) : 182 - 187 .

Rajagopalam, P. K. (1980) .

Insecticide resistance in *Culex pipiens fatigans* wield . Bull. WHO, 22 : 587 - 590 .

Rathburn, C. B. & Boike, A. H. (1975) .

Laboratory and small plot field tests of altosid and dimilin for the control of *Aedes taeniorhynchus* and *Culex nigripolus* larvae . Mosq. News, 35 (4) : 540 - 546 .

Raymond, M. & Marquine, M. (1994) .

Evaluation of insecticide resistance in *Culex pipiens* populations . The corsican paradox . J. Evolutionary Biology, 7 (3) : 315 - 337 .

Rettick, F. (1978) .

Effect of diflubenzuron on four species of mosquitoes (Diptera : Culicidae) . Aceta Entomol . Bohemoshav ., 75 (5) : 312 - 315 .

Revanna, M. A. ; Vijayan, V. A. & Gowda N. N. (1991) .

Larval susceptibility status of *Culex gelidus* Theobald , against four organophosphorous compounds in Mysore City . J. of communicable Diseases , 23 (3) : 202 - 203 .

Rohani, A. ; Syuriza, A. S. ; Lee, H. L. ; Abdallah, A. G. ; Ong, Y. F. Chan, S. T. & Ong, K. K. (1995) .

Tropical, Biomedicine, 12 (1) : 39 - 44 .

Salama, S. A. ; Hussein, R. P. ; El-Husseini, M. M. & Abdallah, S. K. (1984) .

Antimoulting insecticide for mosquito control and its effect on transmission of African horse sickness virus by treated *Culex pipiens* L. J. Vet. Med. Ass., 13 (25) : 97 - 112 .

Saleh, M. S. & Aly, M. I. (1987) .

The biological effects for three insect growth regulators on *Culex pipiens* (L.) Anzeiger fur shadlings Kunde, pflanzen-schatz, Umweltschutz ., 60 (2) : 34 - 37 .

----- ; Kelada, N. L. & Abdeen, M. I. (1990) .

Factors affecting of *Bacillus thuringiensis* H. 14 against mosquito larvae with special reference to the joint action of the pathogen with three chemical insecticides . Anzeiger fur schadling skunde, Pflanzenschutz, Umweltschutz, 63 : (11) 10 - 13 .

Sawicki, R. M. (1979) .

Detection of pyrethroid resistance in houseflies in Britain .
Rothamsted report for 1978, part 1 : 133 .

Saxena , S. C. ; Kaushik, R. & Simh, R. (1992) .

Growth regulating activity of bendiocarb, a carbamate compound against mosquito vectors, *Anopheles stephensi* and *Culex quinquefasciatus* . Entomol, 17 (1 & 2) : 131 - 134 .

Schaefer, C. H. (1979) .

Effects of bay sir 8514 on the clear lake gnat and non-target organisms . Pap. Proc. Annu. Conf. Calif. Mosq. Vector control Ass., 47 : 33 .

----- & Wilder, W. H. (1972) .

Insect developmental inhibitors, a practical evaluation, as mosquito control agents (Diptera : Culicidae) . J. Econ. Entomol., 65 (4) : 1066 - 1071 .

----- & ----- (1973) .

Insect developmental inhibitors . 2. Effects on target mosquito species . J. Econ. Entomol., 66 (4) : 913 - 916 .

Self, L. S. ; Tonn, R. J. ; Bai, D. H. & Shin., H. K. (1973) .

Toxicity of agricultural pesticide applications to several mosquito species in South Korean rice . Trop. Med., 15 (4) : 177 - 188 .