

Summary

Psammophytes are plants that inhabit the sand formations including: sand bars, sand hillocks and sand dunes in the coastal and inland deserts of Egypt. They are specialized plants due to their ecological and economical values. The present investigation deals with the ecological and taxonomical studies of seven psammophytic species categorized under two main groups:

I- Gramineaceous species:

The three selected Gramineae species are: *Ammophila arenaria*, *Elymus farctus* and *Halopyrum mucronatum*.

II- Non- Gramineaceous species:

The four selected Non- Gramineae species are: *Atriplex farinosa*, *Euphorbia paralias*, *Lotus polyphyllus* and *Populus euphratica*.

The studied psammophytic plants are characterized by a particular combination of morphological and anatomical adaptations allowing them to survive under water stress, high salt concentration, high temperature, and high irradiances. These adaptations include: Rolling or shedding of leaves to reduce transpiring area (*Ammophila arenaria*, *Elymus farctus* and *Halopyrum mucronatum*). Presence of salt bladder, white silvery hairs and wax layer on leaf surface (*Atriplex farinosa*, *Lotus polyphyllus*, *Euphorbia paralias* and *Populus euphratica*) which increase reflectance of solar

radiation and decrease cuticular transpiration. Presence of sand sheath (fine sand grains around root), increasing in root/ shoot ratio and long fibrous roots in the studied grass species to maintain and retain any moisture content around root. Deposition of thick cuticle, dense macro-hairs (unicellular hairs), leaf furrows, sunken stomata, narrow intercellular spaces, size and density of vascular system, CAM photosynthetic pathway (C_4 plants) and stem assimilants are internal characters helping in suppression of the internal evaporation, increase the efficiency of photosynthetic activity and enable plants to quickly counteract water loss.

A total of fifty two (52) stands were sampled representing the studied plants (12 stands for *Ammophila arenaria*, 10 stands for each of *Elymus farctus* and *Atriplex farinosa* two stands for each of *Halopyrum mucronatum* and *Populus euphratica* and 8 stands for each of *Euphorbia paralias* and *Lotus polyphyllus*). Stands and sites were selected on the basis of the field visual variation of vegetational coverage, climatic and edaphic characteristics prevailing in the study area.

This investigation shows that climatic factors, soil salinity, calcium carbonate ratio, , and ratio of silt and clay are the most important factors affecting in morphological, anatomical, distribution and vegetation cover of the studied psammophytic species.

Four of the studied plants, namely: *Ammophila arenaria*, *Elymus farctus*, *Lotus Polyphyllus* and *Euphorbia paralias* are restricted to the

Mediterranean coastal sand dunes, where the climate is mild and less arid favorable for plant life than in the inland extreme arid desert. While *Atriplex farinosa* and *Halopyrum mucronatum* are restricted to the Red Sea coastal sand dunes. On the other hand *Populus euphratica* is restricted only to the sand dunes around the lakes of Siwa Oasis, Western Desert.

Fifty three (53) plant species were associated to the studied species at different sites. These species are belonging to forty six (46) genera and 25 families. This low and limited number of species may be due to the instability of the soil of the sand dunes.

The main families of the associated species are: Gramineae (16.98 %), Chenopodiaceae (13.2 %) and Compositae (11.32 %). Cruciferae, Umbelliferae, Boraginaceae, etc are present in low ratio. The chorology of the studied associate species revealed that: most of the recorded species are: Mediterranean (30%), Saharo-Arabian (26.41%) and Mediterranean - Saharo-Arabian (9.43 %), elements.

Few species are belonging to the Australian, Sudanian, Irano-Turanian, and Pluriregional elements. The life forms are: Chamaephyte (32 %), Hemicryptophyte (24.52 %), Phanerophyte (22.64 %), Therophyte (15.0%) and Geophyte (5.84%).

Vegetation analysis were carried out quantitatively and qualitatively (by quadrat and line intercept methods) to determine the relative frequency, relative density and relative cover for each species.