

REFERENCES

- Abd- El- Shafy, E. and El- Saadawi, W. (1982) : Macroflora from the Jurassic of Ras El-Abd , Gulf of Suez, Egypt. Bull. Fac. Sci., Zagazig Univ., 4: 33-40.*
- About-El- Ela, N , El-Saadawi , W. E., Darwish, M. H. (1989) : Some Lower Cretaceous Microflora from Abu Darag area, western side of the Gulf of Suez Egypt . Egypt J. Geol, 33, 1-2 : 347-361.*
- Barron, T. (1905) : On the age of Gebel Ahmer Sands and Sandstone, the petrified forest, and the associated Lavas between Cairo and Suez. Geol. Mag. London, N. S. dec. v, 2,58-62. (Reviewed by Allen, H. A., 1907, in : Geol. Cent., Leipzig, Bd. 9, 7,316 - 317).*
- _____ (1907) : The topography and geology of the district between Cairo and Suez. Cairo.*
- Barthel, K. W., and Bottcher, R. (1978) : Abu Ballas Formation : A significant lithostratigraphic unit of the former Nubian Series : Mitt. Bayer. Staats Paleontol. Hist. Geol. 18: 155-166 .*
- Barthoux J. C. and Fritel. P. H. (1912) : Sur des empreintes (Meduses, Algues) recueillies dans le Carbonifere de environ de Suez c.r. Acad. Sci., Paris,155, 795-796 .*
- _____ (1925): Flore cretacee du gres de Nubie. Mem. Inst. Eg.7,65-119. (Reviewed by Lorin, H., 1925 , in : Bibliogr Geogr.,Paris, Annee 35, p. 434).*

- Blanckenhorn, M. (1901):** Geologie Agyptens , Fuhrer durch die geologische Vergangenheit Agyptens von der Steinkohlenperiode bis zur Jetztzeit- Berlin .
- Bonnet, E. (1939) :** *Nipadites sickenbergeri*, Gebel Giuschi, Untermokattam, Mideocene. (From Kaul, K. N., 1960)
- Bottcher, R. (1985) :** Environmental model of the shallow marine Abu Ballas Formation (Aptian, Nubia Group) In South-Western Egypt. N. Jb. Geol. Palaot. Abh., 169 (3), 261-283, Stuttgart.
- Boureau, E., Lapparent, A. F. (1951) :** Decouverte de *Weichselia reliculata* Stokes et WEBB = *Paradoxopteris stromeri* Hirmer dans le Jurassique due Sud de la Tunisia, C. R. Som. Soc. Geol. Fr., 7, 108-109, Paris.
- Carlquist, S. (1977) :** Wood anatomy of Onagraceae, additional species and conceptans Missouri Bot. Gard. 64 : 627-637.
- _____ **(1983):** Wood anatomy of *bubbia* (Winteraceae), with comments on origin of vessels in dicotyedons. Amer. J. Bot. 70 (4): 578-590.
- _____ **(1985) :** Wood and stem anatomy of Misodendraceae; systematic, and ecological conclusions. Brittonia, 37 (1) : 58-75.
- _____ **(1987a) :** Wood anatomy of Nolanaceae. Aliso. 11(4): 463 - 471.
- _____ **(1987 b):** Wood anatomy and relationships of Stackhousiaceae. Bot. Jahrb. Syst. 108 (2,3) : 473 - 480

- _____ (1987 c) : Wood anatomy of noteworthy species of *Ludwigia* (Onagraceae) with relation to Ecology and systematics. Ann. Missouri. Bot. Gard . 74: 889-896.
- _____ (1987d) : Wood anatomy of Martyniaceae and Pedallaceae. Aliso 11(4) : 473 - 483.
- _____ (1988a) : Wood anatomy and relationships of Duckeodendraceae Goetzeaceae. IAWA Bulletin N.S., Vol. 9 (1) : 3-12 .
- _____ (1988 b) : Wood anatomy of Cytopetalacea e. Aliso. 12(1) : 63-76.
- _____ (1988c) : Wood anatomy of Acanthaceae. Asurvey. Aliso 12(1) : 201 - 227 .
- _____ (1989a) : Wood anatomy of *Cercidium* (Fabaceae), with emphasis on vessel wall sculpture- Aliso. 12(2):235-255.
- _____ (1989b) : Wood anatomy relationships of *Montinia* . Alsio 12(2) 369 - 378 .
- _____ & Zonas , S. (1984) : Wood anatomy of Malesherbiaceae. Phytomorphology , Vol. 34 (1-4) : 180 - 190 .
- Carruthers, W. (1870) :** On the Petrified forest near Cairo. Geol. Mag. London, 7, 306-310.
- Chattaway , M. M. (1949) :** The development of tyloses and secretion of gum in heartwood formation. Aust. J. Sci. Res., Ser. B2, 227-240.

- Chiarugi, A. (1933)** : Tronchi silicizzati di un'alga arborea silurico - devoniana "*Nematophyton saharianum*" n. sp., nel Deserto Libico presso le Oasi di Cufra Giorn. Bot. Ital., Firenze, (n.s.) 40, 590-594. (Reviewed by Tavani, G., 1933-1934, in: Rev. Geol., Liege, 14, 237-238).
- _____ **(1934)** : Una Tallofita arborea silicizzata del deserto Libico: *Nematophyton saharianum* n. sp. Missione, Sci. R. Acc. d' Italia a Cufra (1931), Roma, 3, 291-319.
- Cotta, K., (1832)** : Die Dendrolithen in Beziehung auf ihren innern Bau. Dresden u. Leipzig 1832. 4 : 89 pp., 20 tab .
- Couvert, M. (1977)** : Atlas D'anatomie des charbons de foyers prehistoriques afrique du nord Temperee. Mem. No. xxvi. C. R. A. P. E.
- Cuvillier, J. (1928)** : Les vegetaux fossiles d, Egypte. Soc. Royale Geogr. Bull. le Caire, (n.s.), 15, 289-305.
- Darwish, M. H. (1990)** : Studies on some Egyptian fossil plants - p.H. D. Thesis, Dept. Bot. Fac. of Girls, Ain Shams Univ. Cairo .
- Dominik, W. (1985)** : Stratigraphie und Sedimentologie (Geochemie, Schwermineralanalyse) der Oberkreble von Bahariya und ihre korrelation zum Dakhla Becken (Western Desert, Agypten). Berl. Geowiss. Abh. 62(A) : 1-173 .
- Edwards, W. N. (1932)** : Some Mesozoic plants from Africa. Ann. and Mag. Hist. 10, 406 - 411.

- El-Ramly , M. F. (1973) :** Final report on geomorphology ,
hydrogeology , planning for groundwater resouces and
land reclamation in Lake Nasser region and its environs
Desert Research Institute , Cairo .
- El-Saadawi, W. E. (1979) :** A bibliography concerning fossil
plants of Egypt. *Phytologia* , 43(2) : 353 - 69 .
- _____, **Badawi, A.A. and El-Awamri, A.A. (1975) :**
On silicified rhizome fragments of *phragmites*
communis Trin. From the Pleistocene J. El- Fayum.
Palaeoentographica Abt. 13 . 154, 172-178.
- _____, **Badawi, A. A., and El-Awamri , A.A. (1979) :**
Silicified root fragment of *Tamarix* L. from the pleiso-
cene of El-Fayum. *Gir's coll. Ain Shams Univ.* 12 :
322-335.
- _____, **and Farag, E. (1972) :** Some Mesozoic plants
from Abu - Darag, western side of Gulf of Suze. Egypt.
J. Bot., 15 : 121 - 130 .
- _____, **and Kedves, M. (1991) :** Palaeobotanical
investigations on palnt impressions and sporomorphs
from Egypt . *Plant Cell Biology and Development*, 2 :
8- 33.
- _____, **Shaaban, A. A., and El-Awamri, A. A. (1979)**
Pleistocene diatoms from El-Fayum. *Proc. Egypt. Acad*
. Sci. 31 : 257 - 263.
- Fahn, A., Ella, W. & Pieter, B. (1986) :** Wood anatomy and
identification of trees and and shrubs from Israel and
adjeacent regions. Israel. Jerusalem.
- Felix , J. (1882) :** Studien uber fossile holzer. Diss. Leipzig.

- _____ (1883) : Die fossilen Halzer westindiens . Cassel.
- Fontaine, w N. (1889) : The potomac or yaunger Mesozoic flora - U.S. Geol. Survey, VI, 1-144, 44 pl., Washington.
- Fontaine , W. N. (1889) : The Potomac or younger Mesozoic flora - u.s. Geol. Survey, 15 (I, II), 1-377, 180 pl., Washington.*
- Fourtau, R. (1898) : Note sur l'age des forets petrifiees des, deserts d'Egypte : Bull . Soc. Khed. de Geogr . 8 pp.*
- Fraas, O.F. (1867): Geologisches aus dem Orient : Sinai, Palastina, und Agypten. Jahresh. Ver. Naturk. Wurt., Stuttgart, Bd. 23 , 145-362.*
- Frenguelli, G. (1927) : Diatomee dei travertini del Uadi 'Refuf, presso l'Oasi di Kharga nell'Alto Egitto.Boll. Soc. Geol. Ital., Roma , 46, 1-12.*
- Ghous, A. K., M. & Yunus, M. (1972): Research trends in plant anatomy. TATA. McGraw. Hill.*
- Gothan, W. and Weyland, H. (1973) : Lehrbuch der pataobtanik Boraginaceae (S. I.). I cordioideae.IAWA Bull . n.s., 4: 161-178.*
- Gottwald, H. (1983) : Wood anatomy studies of Boraginaceae (s.I.) I. Ordioideae . IAWA Bull n.s. 4, 161-178.*
- Gregor, H.J. and Hagn (1983) : Fossil fructifications form the Cretaceous- Paleocene boundary of SW Egypt (Danian, Bir Abu. Mungar). Tert. Res. 4:121 -147.*
- Greguss, P. (1955) : Identification of living Gymnosperms on the basis of xylotomy : Budapest .*

- Hanelt, P. (1976)** : Diapensiales - Ebenales und Tubiflorae (Polmoniaceae, Callitrichaceae) In : *Urania Pflanzenreich Hohere, Pflanzenreich. Hohere Pflanzen* 2,s,169-193 und s 220-238. Leipzig- Jena-Berlin.
- Hartig, T. (1848)** : Beitrage zur Geschichte der Pflanzen und zur Kenntniss der norddeutschen Braukohlen- Flora : Bot. Zeitung, Band 6, p. 166-172.
- Heer, O. (1876)** : Uber fossile Fruchte der Oase Chargeh, Denksch. schweiz. naturf. Ges ., Bd. 27, 11s. Zurich .
- Ibrahim, M. M. (1943)**: The petrified forest. Inst. Egypt Bull., 25 159-182.
- _____ **(1953)** : The petrified forest, Pt. II. Inst. Egypt Bull., 34, 317-328.
- Issawi, B., and Jux, U. (1982)** : Contribution to the stratigraphy of the Paleozoic rocks in Egypt. Geol. Surv. Egypt, paper No. 64, 28 p.
- Itier, J. (1974)** : Desfore'ts P'etrifi'es del, Egypte et de la Libye, 16pp Montpellier.
- Joy, K. W., Willis, A. J., & Lacey, W.S. (1956)** : A rapid cellulose peel technique in Palaeobotany. Ann. Bot., N. S. 20 : No. 80, 635-637.
- Jux, U. and Issawi, B. (1983)** : Cratonic Sedimentation in Egypt during the Paleozoic. Ann. Geol. Surv. Egypt., XIII, 223-245.
- Kaul, K. N. (1964)** : The Anotomy of the stem of palms and the problem of the artificial genus *Palmoxylon Schenk*. NBC. Lucknow, India
- Klitzsch, E. (1978)** : Geologische bearbeitung sudwest- Agyp'tens. Geologisch rundschau , 67, (2), 509-520. Stuttgart.

- _____ (1984) : North western Sudan and bordering areas, geological development since Cambrian time. Berliner Geowiss Abh., A/50, 23-43. Berlin.
- _____ *Harms, J.c. Lejal-Nicol & List, F.F. (1979) :*
Major subdivisions and depositional environments of Nubia strata southwestern Egypt. Bull. Am. Assoc. Petrol. Geol. 63 (6), 974-976.
- _____ & *Lejal - Nicol, A. (1984):* Flora and faune from strata in southern Egypt and northern Sudan (Nubia & Surrounding areas). Berliner geowiss. Abh. (A), 50 ; 47-79, Berlin.
- Krausel, R. (1924) :* Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wusten Agyptens. IV., (1), (2), (3); (A) Fungi, Algae; (B) Gymnospermae, Coniferae; (c) Angiospermae, Moncotyledoneae; Abh. Bayer. Akad. Wiss. Munchen, 30, 1-48.
- _____ (1939) : Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wusten Agyptens, 3. Die fossilen Pflanzen Agyptens, EL Abh. Bayer. Akad. Wiss., Munchen (N.F.) 47,1-140.
- Lacey, W.S. (1963) :* Palaeobotany Technique, View Points in Biology, (Editors J. D. Carthey and .I. Duddington) 2 : 202 - 243).
- Lejal - Nicol, A. (1981) :* A propos de nouvelles flores paleozoiques et mesozoiques de l'Egypte due Sud-Ouest. C.R. Acad. Sci. Paris, 292 D: 1337-1340, Paris.

- _____ (1987) : New Paleozoic and Mesozoic Floras from Egypt and Northern Sudan. Berl. Geowiss. Abh. 75(A): 151 -248.
- _____ & Dominik, W. (1987) : Flores nouvelles du Paleozoique et du Mesozoique D' Egypte et du Soudan septentrional Berl. Geowiss. Abh. 75 (A): 151 - 248.
- Linant, de B. (M. A.) (1840)** : Notice sur la Fore'ts petrifie'e des environs du Caire. Bull. Soc. Geogr., Paris, (ser.2), 13, 97-107.
- Louvet, P. (1973)** : Sur les affinites des flores tropicales ligneuses africaines tertiaires et actuelles. Bull. Soc. Bot. France. 120: 385-396.
- Mair, M. N., B. (1988)** : Wood anatomy and heartwood formation in Neem (*Azadirachta indica* A. Juss.) Botanical J. of the Linnean Society 97 : 79-90.
- Manchester, S. R. (1979)** : *Triplochitoxylon* (Sterculiaceae) . A new genus of wood from the Eocene of Oregon and its bearing on xylem evolution in the extant genus *Triplochiton*. Amer. J. Bot. 66 (6) : 699-708 .
- Mayer -Eymar, K. (1893)** : Le Ligurien et le Tongrien en Egypte. Bull. Soc. Geol. France, ser. 3, 7-43. (Appeared also in Bull. Inst. Eg. 1893 et 1896, according to Cuvillier, 1928).
- Metcalf, C.R. and Chalk, L. (1950)** : Anatomy of the dicotyledons. Oxford.
- _____ (1961): Anatomy of the Monocotyledons. Oxford

- Montagne, J. F. (1844)** : Extrait de deux lettres adressés par M. le Dr. Montagne à M. Jomard sur un phénomène observé dans la Mer Rouge. Bull. Soc. Geogr., Paris, (ser. 3), 1, 149-153.
- Muhammad, A. F., & Satteler, R. (1982)** : Vessel structure of *Gnetum* and the origin of angiosperms. Amer. J. Bot. 69 (6) : 1004-1021.
- Muller-Stoll, W. R. & Madel, E. (1967)** : Die fossilen Leguminosen - Holzer. Palaeontographica B : 95-174.
- Osman, R. (1992)** : Contribution to the geology of the area southwest of Aswan. Ph. D. Thesis. Geo. Dept., Fac. Sci., Benha University .
- Petresu, J. (1978)** : Studiul lemnoșilor fosile din oligocenul din Nord-Vestul Transilvaniei. Memoires (Bucarest), 27, 113-184.
- Pigg, B. & Rothwell, G. W. (1979)** : Stem root transition of an Upper Pennsylvanian woody lycopsid. Amer. J. Bot. 66 (8) : 914-924.
- Prakash, U. Q., Barghoorn, E.S. & Scott, R. A. (1968)** : Fossil wood of *Robinia* and *Gleditsia* from the Tertiary of Montana. Amer. J. Bot. 49(7) : 692-696.
- Renner, O. (1907)** : *Teichosperma*, eine Monokotylenfrucht aus dem Tertiär Ägyptens. Beitr. z. Pal. u. Geol. Osterr. - Ung. u. d. Orients, Bd. 20, 217-220.

- Roziere, F. (1813-1824) :** De la constitution physique de l' Egypte et de ses rapports avec les anciennes institutions de cette contree. Description de l' Egypte (Histoire Naturelle), Paris. 1st Edn. 1813, tome 21, 407-732.
2nd. Edn. 1824, tome 20, 211-523, and tom 21 in 1826, 1-324.
- Sahni, B. (1931) :** Material for a Monograph of the Indian petrified palms. Proc. Acad. Sci. United Provinces of Agra and Oudh,1.
- Said, R. (1962):** The geology of Egypt. Elsevier, 377 pp.
_____ (1990) : The geology of Egypt. Published for the Egyptian General Petroleum Corporations Conoco Hurghada Inc. and repsol exploracion, S.A. By A. A. Balkema , Rotterdam/Brookfield /1990 .
- Schenk, J. H. A. (1880) :** Ueber fossile Holzer aus der Libyschen Wuste. Bot. Zeit., Berlin, Bd. 38, 657-661.
_____ (1888) : Fossil Holzer aus Ostasien und Aegypten. Bih. Kgl. Srwensk. Vet. Akad. Handl. 14, 111.
- Schrank, E. (1982) :** Kretazische Pollen und Sporen aus dem "Nubischen Sandstein" des Dakhla Beckens (Agypten). Berl. Geowiss. Abh. 40 (A) : 87-109.
- Schweinfurth, G. (1882) :** Zur Beleuchtung der Frage uder den versteinerten Wald. Zeitschr. D. Geol. Ges., Bd. 34 , 139-145, Berlin.
_____ (1887) : Sur une recente exploration geologique de l' Ouadi Araba. Bull. Inst. Egypte, 18.

- Schwob, G. (1863) :** Analyse du Memoire de M. Unger sur la
Foret petrifiee pres du Caire. Bull. Inst. Eg., ser. 1,
no. 8 (1862-1863), Fevr. 1863, 71-73.
- Scurfield, G. A., & Segnit, E.R. (1984) :** Petrification of wood
by silica minerals. Sediment. Geol. 39: 149 - 167.
- Seward, A. C. (1935) :** Leaves of dicotyledons from the Nubian
Sandstone of Egypt. Ministry of Finance, Geological
Survey of Egypt, Cairo., (Reviewed by Cuvillier, J.
1936. In : Rev. Geol., Liege, 16, 1936-1937, p . 412).
- Shukri, N. M. (1953) :** The geology of the desert east of Cairo.
Bull. Inst. Desert, Egypt 3 (2) : 89 - 105.
- Stein, W., E. and Beck, C. B. (1978) :** *Bostonia perplexa* gen.
Et-sp. Nov. , a *Calamopityan* axis from the new Albany
shale of Kentucky. Amer. J. Bot. 65 (4) : 459-465.
- Stenzel, K. G. (1904) :** Fossile Palmenholzer. Beit. z. Pal. u. Geol
Osterr.- Ung. u.d. Orients. Bd. 16, 107-287. Wien.
- Stern, W. L. (1967) :** *Kleinodendron* and xylem anatomy of
Cluytieae (Euphorbiaceae) Amer. J. Bot. 54 (6) :
663-676.
- Suss, H. and Gregor, H. J. (1987) :** Two new silicified woods
Cordioxydon barthelli new-genus, new-species and
Enebenoxydon saharicum new-genus, new-species from
the southwestern Sahara, Egypt. Feddes repert 98-(9-10) :
521-536.
- Unger , F.J. A. N.(1845) :** " De Palmis Fossilibus" chap. 2;
Liii Martius Genera Palmarum Vol. 1(1831 - 1850).
- Unger, F.J. A. N. (1858-1859) :** Der versteinerte Wald bei Kairo

und einige andere Lagerverkieslten Holzes in Agypten.
Sitz Ber. K. Akad. Wiss. M. N. K. Bd. 33, 1858, 209 -
233, Wien 1859 (Also reviewed in Q. J. G.S., Lond.,
15, p. 13).

Vaudois - Mieja, N., Lejal - Nicol, A. (1987): Paleocarpologie
africaine : apparition des l'Aptien en Egypte d' un
Palmier (*Hyphaenocarpon aegyptiacum n. sp*). - C. R.
Acad. Sci. Paris .

Zittel, K. A. (1874) : Etudes geologiques ; constitution
geologique du desert Libyque et des Oasis, indication
des divers terrains et fossiles. Bull. Inst. Eg., No.
13,75-83.