

# CONTENTS

|   | Page      |
|---|-----------|
| <b>ABBREVIATIONS</b>  | <i>m</i>  |
| <b>1. INTRODUCTION</b>  | <b>1</b>  |
| <b>2. LITERATURE REVIEW</b>   | <b>4</b>  |
| <b>2.1 Occurrence of The Tested Fungal Pathogens in Soybean And Sunflower Plants.</b>   |           |
| 2.1.1 Occurrence of <i>Macrophomina phaseolina</i> in Soybean   | 6         |
| 2.1.2 Occurrence of <i>Fusarium solani</i> in Soybean   | 7         |
| 2.1.3 Occurrence of <i>Macrophomina phaseolina</i> in Sunflower   | 9         |
| 2.1.4 Occurrence of <i>Fusarium solani</i> in Sunflower   |           |
| <b>2.2 Bacteria, Streptomycetes And Trichoderma As Antagonists Against Soil Borne Fungal Plant Pathogens</b>                                | <b>9</b>  |
| 2.2.1 <i>In vitro</i>   | 9         |
| 2.2.1.1 Bacteria ( <i>Bacillus</i> spp.) As An Antagonist Against Soil Borne Fungal Plant Pathogens.  | 9         |
| 2.2.1.2 Another Type of Bacteria (Other Than The Genus <i>Bacillus</i> ) Antagonist Against Soil-Borne Fungal Plant Pathogens.              | 10        |
| 2.2.1.3 <i>Streptomyces</i> as an Antagonist Against Soil-Borne Fungal Plant Pathogens.   | 12        |
| 2.2.1.4 <i>Trichoderma</i> spp. ( <i>T. harzianum</i> ) As An Antagonist Against Soil-Borne Fungal Plant Pathogens ( <i>M. phaseolina</i> ) | 13        |
| 2.2.1.5 <i>T. Harzianum</i> as an Antagonist Against Soil-Borne Fungal Plant Pathogens (Other Than <i>M. phaseolina</i> )                   | 14        |
| 2.2.1.6 Other Species of <i>Trichoderma</i> . as an Antagonists Against Soil-Borne Fungal Plant Pathogens.                                  | 17        |
| 2.2.2 <i>In vivo</i>  | 18        |
| 2.2.2.1 Seed Treatment With Bacteria as Biocontrol Agents   | 18        |
| 2.2.2.2 Soil Treatment With Bacteria <i>B. subtilis</i> As Biocontrol Agent   | 20        |
| 2.2.2.3 Seedling Treatment With Bacteria As Biocontrol Agents   | 21        |
| 2.2.2.4 Seed Treatment with <i>Streptomyces</i> spp. as Biocontrol Agents.  | 21        |
| 2.2.2.5 Soil Treatment With <i>Streptomyces</i> as Biocontrol Agent   | 23        |
| 2.2.2.6 Seedling Treatment with <i>Streptomyces</i> as Biocontrol Agent   | 24        |
| 2.2.2.7 Seed Treatment With <i>Trichoderma</i> as Biocontrol Agent  | 24        |
| 2.2.2.8 Soil Treatment With <i>Trichoderma</i> spp. As Biocontrol Agent   | 25        |
| <b>2.3 Isolation of Actinomycetes Form Natural Habitats</b>   | <b>26</b> |
| 2.3.1 Isolation of <i>Streptomyces</i> from Natural Habitats  | 27        |
| 2.3.1.1 Biochemical Activities of <i>Streptomyces</i>   | 28        |
| 2.3.1.2 Production of Antibiotic  | 28        |
| <b>2.4 Phytoalexins Production</b>  | <b>28</b> |
| 2.4.1 <i>Phytoalexins</i> Production in Soybean   | 29        |
| 2.4.1.1 Elicitation of <i>Phytoalexins</i> Production   | 29        |
| 2.4.1.2 Accumulation of <i>Phytoalexins</i> Production  | 30        |
| 2.4.1.3 Mechanism(s) of <i>Phytoalexins</i> Production  | 33        |
| 2.4.2 <i>Phytoalexins</i> production in sunflower   | 35        |
| 2.4.2.1 Elicitation of <i>phytoalexins</i> Production   | 35        |
| 2.4.2.2 Accumulation of <i>Phytoalexins</i> Production  | 35        |
| 2.4.2.3 Mechanism of <i>Phytoalexins</i> Production   | 35        |
| <b>2.5 Scanning Electron Microscopy</b>   | <b>36</b> |

|   |           |
|---|-----------|
| <b>3. MATERIALS AND METHODS</b>   | <b>38</b> |
| <b>3.1 Source and Origin of Cultures of Microorganisms Used in This Study</b>   | <b>38</b> |
| 3.1.1 Culture of The Tested Phytopathogenic Fungi   | 38        |
| 3.1.2 Culture Of The Tested Non Phytopathogenic Microorganisms Served As Biocontrol   | 38        |
| 3.1.3 Culture of The Other Tested Non Phytopathogenic Microorganisms Other Than the Ones Served as Biocontrol Agents  | 39        |
| <b>3.2 Isolation of <i>Streptomyces</i> species antagonistic to <i>M. phaseolina</i> And <i>F. solani</i> f. sp. <i>phaseoli</i> at a distance From the Soil - Rhizosperic Region</b> | <b>39</b> |
| 3.2.1 Collection of Soil Samples  | 39        |
| 3.2.2 Isolation Technique Employed  | 40        |
| 3.2.3 Maintenance and preservation of cultures  | 41        |
| <b>3.3 Media used</b>   | <b>42</b> |
| <b>3.4 Plant Host Material</b>  | <b>42</b> |
| 3.4.1 Source of Seeds   | 42        |
| 3.4.2 Growth of Plants  | 43        |
| <b>3.5 Methods of Inoculation</b>   | <b>43</b> |
| 3.5.1 Inoculum Preparation  | 43        |
| 3.5.2 Inoculum Application Techniques:  | 43        |
| 3.5.2.1 Using Excised Hypocotyls Under Controlled Environment (excised hypocotyl inoculation technique)   | 43        |
| 3.5.2.2 Using Artificial Seed Infestation By Fungal Spore Suspension, Under Greenhouse Conditions.  | 44        |
| <b>3.6 Disease severity assessment</b>  | <b>47</b> |
| <b>3.7 Re-Isolation And Pathogenicity Tests</b>   | <b>48</b> |
| <b>3.8 Biocontrol</b>   | <b>48</b> |
| 3.8.1 In vitro Tests Using The Dual Culture Technique   | 48        |
| 3.8.1.1 Antagonism Studies  | 48        |
| 3.8.1.1.1 Antagonistic Ability of <i>B. subtilis</i> against <i>M. phaseolina</i> And <i>F. solani</i> f.sp. <i>phaseoli</i> .  | 48        |
| 3.8.1.1.2 Antagonistic Ability of Various <i>Streptomyces</i> Isolates Against <i>M. phaseolina</i> and <i>F. Solani</i> f.sp. <i>phaseoli</i> .                                      | 49        |
| 3.8.1.1.2.1 Solid Media   | 49        |
| 3.8.1.1.2.1.1 Streak Method.  | 49        |
| 3.8.1.1.2.1.2 Reversed Layer Method.  | 50        |
| 3.8.1.1.2.2 Liquid Media  | 50        |
| 3.8.1.1.2.2.1 Bore Diffusion Method.  | 50        |
| 3.8.1.1.2.2.2 Antibiotic Production on Agar Plates.   | 51        |
| 3.8.1.1.3 Antagonistic Ability of <i>T. harzianum</i> Against <i>M. phaseolina</i> And <i>F. solani</i> f.sp. <i>phaseoli</i>   | 52        |
| 3.8.2 In vivo   | 52        |
| 3.8.2.1 Biocontrol of Soybean Damping off Caused Separately by <i>M. phaseolina</i> and <i>F. solani</i> f.sp. <i>phaseoli</i>  | 52        |
| 3.8.2.1.1 Seed Treatment Using Various Biocontrol Agents Under Greenhouse Conditions.   | 52        |
| 3.8.2.2 Biocontrol of Sunflower Damping off Caused Separately by <i>M. phaseolina</i> And <i>F. solani</i> f.sp. <i>phaseoli</i>  | 52        |
| 3.8.2.2.1 Seed Treatment Using Various Biocontrol Agents Under Greenhouse Conditions  | 52        |
| <b>3.9 Effect of Culture Filtrates of The Biocontrol Agents Used in This Study on Seed Germination (In Vitro Culture Phytotoxicity Assay)</b>   | <b>53</b> |
| <b>3.10 Anatomical Examination By Light Microscopy</b>  | <b>54</b> |
| 3.10.1 Preparation of Epidermal Strips  | 54        |
| 3.10.2 Toluidin Blue O Stain  | 54        |

|   |            |
|---|------------|
| <b>3.11 Phytoalexin Techniques</b>  | <b>54</b>  |
| 3.11.1 <i>Extraction Method For Phenolic Compounds</i>  | 54         |
| 3.11.2 <i>Chromatographic Technique</i>   | 55         |
| 3.11.2.1 <i>TLC</i>   | 55         |
| 3.11.2.1.1 <i>Location And Detection of Compounds</i>   | 55         |
| 3.11.2.1.1.1 <i>Ultraviolet radiation (U.V)</i>   | 55         |
| 3.11.2.1.1.2 <i>Elution Procedure</i>   | 56         |
| 3.11.2.1.1.3 <i>Spectrophotometry</i>   | 56         |
| 3.11.2.2 <i>HPLC Technique</i>  | 56         |
| <b>3.12 Electron Microscopy</b>   | <b>56</b>  |
| 3.12.1 <i>Scanning Electron Microscopy</i>  | 56         |
| <b>4. RESULTS</b>   | <b>57</b>  |
| <b>4.1 Isolation And Differentiation of Streptomyces</b>  | <b>57</b>  |
| <b>4.2 Antagonistic Activity of Two Groups Isolates of <i>Streptomyces</i> spp.</b>   | <b>62</b>  |
| 4.2.1 <i>Antagonistic Properties of Red Group Isolates of Streptomyces</i>  | 62         |
| 4.2.2 <i>Antagonistic Properties of Gray Group Isolates of Streptomyces</i>   | 64         |
| 4.2.2.1 <i>Detailed Culture Properties of The Two Selected Isolates of Streptomyces spp. (No. 112 &amp; 729) on Different Media</i>             | 66         |
| 4.2.2.2 <i>Antimicrobial potentialities of the selected isolates of Stryptomyces spp. (No. 112 &amp; 729)</i>                                   | 68         |
| 4.2.2.3 <i>Identification of The Selected Isolates of Streptomyces spp. (No. 112 &amp; 729)</i>   | 71         |
| <b>4.3 Phytotoxicity Bioassay</b>   | <b>73</b>  |
| 4.3.1 <i>Phytotoxicity in Soybean</i>   | 73         |
| 4.3.2 <i>Phytotoxicity in Sunflower</i>   | 75         |
| <b>4.4. Biocontrol Studies</b>  | <b>77</b>  |
| 4.4.1 <i>In Vitro</i>   | 77         |
| 4.4.1.1 <i>Antagonistic Studies</i>   | 80         |
| 4.4.2 <i>In Vivo</i>  | 80         |
| 4.4.2.1 <i>Pathogenicity Test And Disease Evaluation</i>  | 80         |
| 4.4.2.2 <i>Pathogenicity Test By Drop Technique In Growth Cabinet.</i>  | 80         |
| 4.4.2.3 <i>Evaluation of Disease Symptoms Development in Soybean (Excised) Hypocotyls in Growth Cabinet.</i>                                    | 83         |
| 4.4.2.4 <i>Pathogenicity Assessment Under Greenhouse.</i>   | 85         |
| 4.4.3 <i>Efficacy of Several Biocontrol Agents on Disease Severity And Plant Growth</i>   | 91         |
| 4.4.3.1 <i>Effect of Biocontrol (Seed Treatment) on Soybean Macrophomina Charcoal Rot Severity And Plant Growth Under Greenhouse Conditions</i> | 95         |
| 4.4.3.2 <i>Effect of Biocontrol (Seed Treatment) on Soybean Fusarium Root Rot Severity And Plant Growth Under Greenhouse Conditions</i>         | 98         |
| 4.4.3.3 <i>Effect of Biocontrol (Seed Treatment) on Sunflower Charcoal Rot Severity And Plant Growth Under Greenhouse Conditions</i>            | 103        |
| 4.4.3.4 <i>Effect of Biocontrol (Seed Treatment) on Sunflower Fusarium Root Rot Severity And Plant Growth Under Greenhouse Conditions</i>       | 107        |
| <b>4.5 Phytoalexins</b>   | <b>111</b> |
| 4.5.1 <i>Phytoalexins Detection by TLC</i>  | 111        |
| 4.5.1.1 <i>Screening of Phytoalexins Production in Soybean (Hypocotyls)-Pathogens/Biocontrol Agents Interaction Systems, in Greenhouse.</i>     | 111        |
| 4.5.1.2 <i>Screening of Phytoalexins Production in Soybean (Leaves)-Pathogens/ Biocontrol Agents Interaction Systems in Greenhouse.</i>         | 130        |
| 4.5.1.3 <i>Screening of Phytoalexins Production in Sunflower (Hypocotyl)-Pathogen/ Biocontrol Agents Interaction Systems in Greenhouse.</i>     | 140        |
| 4.5.1.4 <i>Efficiency of TLC Used in This Study</i>   | 145        |

|  |            |
|--|------------|
| 4.5.2 Phytoalexins Detection by HPLC   | 150        |
| 4.5.2.1 Screening of Phytoalexins Production in Soybean (Hypocotyl)-M. phaseolina/Biocontrol Agents Interaction Systems in Greenhouse  | 151        |
| 4.5.2.2 Screening of Phytoalexins Production in Soybean (Hypocotyl) F. solani. f.sp. phaseoli/ Biocontrol Agents Interaction Systems in Greenhouse.                                  | 160        |
| 4.5.2.3 Screening of Phytoalexins Production in Soybean (Leaves) M. phaseolina/ Biocontrol Agents Interaction Systems in Greenhouse.   | 166        |
| 4.5.2.4 Screening of Phytoalexins Production in Soybean (Leaves) F. solani f.sp. phaseoli/ Biocontrol Agents Interaction System in Greenhouse.                                       | 174        |
| 4.5.3 Characterization, Distribution and Identification of Phytoalexins Produced in Soybean Hypocotyls and Leaves - Pathogen(s)/ Biocontrol Agents Interaction Systems in Greenhouse | 181        |
| 4.5.3.1 Characterization, Distribution and Identification of Phytoalexins Produced in Soybean (Hypocotyls) - M. phaseolina/Biocontrol Agents Interaction Systems in Greenhouse       | 181        |
| 4.5.3.2 Characterization, Distribution and Identification of Phytoalexins In Soybean Hypocotyls - F. solani f.sp. phaseoli/Biocontrol Agents Interaction Systems in Greenhouse.      | 190        |
| 4.5.3.3 Characterization, Distribution And Identification of Phytoalexins In Soybean (Leaves)- M. phaseolina/Biocontrol Agents Interaction Systems In Greenhouse                     | 198        |
| 4.5.4 Confirmation of the Identity of one of the Major Known Phytoalexin Compounds   | 215        |
| 4.6 Streoscaning Electron Microscopy   | 219        |
| 4.6.1 Interaction Between Soybean- M. phaseolina/Biocontrol Agents Systems: Hypocotyls as Revealed by S.E.M.   | 219        |
| 4.6.1.1 B. subtilis system   | 219        |
| 4.6.1.2 S. roseodiataticus (isolate No. 112) system  | 222        |
| 4.6.1.3 T. harzianum system  | 227        |
| 4.6.2 Interaction Between Soybean- F. solani f. sp. phaseoli/ Biocontrol Agents Systems: Hypocotyls as Revealed by S.E.M.  | 230        |
| 4.6.2.1 B. subtilis system   | 230        |
| 4.6.2.2 S. roseodiataticus (isolate No. 112) system  | 232        |
| 4.6.2.3 T. harzianum system  | 235        |
| <b>5. DISCUSSION</b>   | <b>238</b> |
| 5.1 Impact On Plant Development  | 238        |
| 5.2. Impact on Host Metabolism   | 247        |
| 5.2.1 Impact on Soybean Metabolism as Revealed From TLC Analysis   | 247        |
| 5.2.2 Impact on Sunflower Metabolism as Revealed From The TLC Analysis   | 251        |
| 5.3 Mechanism of Biocontrol.   | 256        |
| 5.3.1 In vivo  | 256        |
| 5.3.1.1 Role of Aromatic Secondary Metabolites as Revealed by HPLC   | 256        |
| 5.3.1.2 Role of Hyperparasitism  | 262        |
| 5.4 Further work   | 265        |
| <b>6. SUMMARY</b>  | <b>266</b> |
| <b>Appendix</b>  |            |
| <b>7. REFERENCES</b>   | <b>275</b> |