

7. REFERENCE

- Abdalla, A.M.; Rizk, F.A. and Adam,S.M. (2001). The productivity of pepper plants as influenced by some biofertilizer treatments under plastic house condition. Bulletin of Faculty of Agricultural, Cairo. University. 52 : 4, 625-639.
- Abdalla, S.O.M. (2002). Effect of organic and biofertilization sources on *cenchrus ciliaris* (L.) at the north western coast of Egypt. Arab Universities Journal of Agricultural sciences. 10(2) :607-617.
- Abdel Azeem, H.H. (2001). Some factors affecting the composting process of rice straw. Annals of Agricultural. Science. 2, 525-542.
- Abdel Ghany, B.F. (1996). Influence of different bacterial strains as biofertilizes on wheat crop production in new cultivated land. . Desert institute- Bulletin, Egypt. 46(2): 215-226.
- Abdel Ghany, B.F. (1997) .New bio-organic approach and their correlation with maize production in new cultivated calcareous soil. : Desert-Institute-Bulletin,-Egypt. 47(2): 363-378..
- Abdel Hafez, A.M. (1966). Some studies on acid Producing microorganisms in soil and rhizosphere with special reference to phosphate dissolvers. Ph. D. Thesis. Fac. Agric., Ain Shams Univ. Cairo, Egypt.
- Abdel Malek, Y. and Ishac., Y.Z. (1968). Evaluation of methods used in counting *Azotobacter* . J. App. Bacteriol., 31, 267- 275.
- Abdel Sabour, M.F.; El Seoud, M.A.A .and Rizk, M. (1999). Physiological and chemical response of sunflower to previous organic waste composts application to sand soil. Egyptian. Journal of soil science. 39 (4): 407-420.

- Abou Bakr, M.A. and Omar, A..M. (1996). Contribution of municipal waste as organic fertilizer to sunflower production. Egyptian – Journal of soil Science. 36 (1/4): 69-82.
- Adediran, J.A ; Taiwo, L.B.; Akande, M.O. ; Sobulo, R.A. and Idowu, O.J. (2004) . Application of organic and inorganic fertilizer for sustainable maize and cowpea yields in Nigeria. . Journal-of-plant-nutrition.; 27(7): 1163-1181
- Adhikary, R.K.; Barua, P. and Bordoloi, D.N. (1991).Influence of microbial pretreatment on degradation of muschorrom nutrient substrate . International- Biodeterioration and Biodegradation 30: 4, 233-241.
- Aleksandrova, G.P.; Medvedeva, S.A.; Babkin, V.A.; Solovev-V.A.; Malysheva, O.N. and Ivanova, S.Z (1989). Effect of composition of the nutrient medium on the ligninase activity of the basidiomycetes phanerochate sanguinea. Khimiya- Drevesiny. 6 (77):80- 126 .
- Allen, I.N. (1959): Experiments in Soil Bacteriology. Burgess Publishing. Co., Minneapolis. Minnesota, USA.
- Anonymous, (1989). The biocycle guide to composting municipal wastes. The JGC press, Inc., Emmaus, Pa. 195 pp.
- Atif, H.; Yasmeen, A. and Rajoka, M.1. (2004). Induction production, repression and de-repression of exoglucanase synthesis in *Aspergillus niger*. Bioresource- Technology. 94 (3) 311-: 319.
- Bakare, M. K.; Adewale, I. O.; Ajayi, A. O.; Okoh, A. and Shonukan, O.O. (2005).Regulatory mutations affecting the synthesis of cellulase in *Pseudomonas fluorescens* .African Journal of Biotechnology.4(8):838-843.

- Barkai, G. R. and Karadavid, R. (1996) . Cellulolytic activity of *Penicillium digitatum* and *P. italicum* related to fungal growth and to pathogenesis in citrus fruit. *Journal-of-Phytopathology*. 131(1): 65-72.
- Barsukov, S.S. and Vasheprudov V.F. (1991) Nutreint uptake by maize. *Khimiza - siya- Sel skogo- khozyaistva*. 8, 80-82.
- Begain, F. (1983): Detection of cellulose activity in polyacrglamide gles using congored stained agar replicas. *Anal. Bioch emistry*. 131: 333- 336.
- Beguin, P. and N.R. Gilkes (1987): Cloning of Cellulase Genes. *CRC Critical Reviews in Biotechnology*, 6 (2): 129- 162.
- Black, C. A. (1965): Methods of soil analysis part 1. Physical and Mineralogical properties.Including statistics of meansurmenets and sampling. Amer. Soc. of Agronomy, Inc., Publisher Madison, Wisconsin, U.S.A.
- Briton, W.F. (1985). Nitrogen response of maize to fresh and composted manure. *Biological Agriculture and Horticulture*. 3:1 , 55- 64.
- Bunt, J.S. and Rovira, A.D. (1955): Microbiological studies of some subantaretic soils, *J. Soil Sci.*, 6: 119- 128.
- Businell , M.; Giglotti, G. and Giusquiani, P.L (1990). The use of compost from town waste in agriculture. *Agrochimica*. 34 : 5-6, 454-466.
- Cahyani, V.R.; Watanabe, A.; Matsuya, K ; Asakawa, S. and Kimura, M .(2002). Succession of microbiota estimated by phospholipid fatty acid analysis and changes in organic consitutents during the composting process of rice straw. *Soil science and plant Nutrition*. 48(5): 735-743.

- Chapman, S.J. and Lynch, J.M. (1985) .Polysaccharide synthesis by capsular microorganisms in coculture with cellulolytic fungi on straw and stabilization of soil aggregates. *Biology-and-Fertility-of-Soils*. 1(3): 161-166 .
- Chung, R.; Wang, C. H.; Wang, C.; Wang, Y. P. and Chung, R.S. (2000). Influence of organic matter and inorganic fertilizer on the growth nitrogen accumulation of corn Plants. *Journal of Plant Nutiriton*. 23(3) : 297- 311.
- Cochran, W.G. (1950): Estimation of bacterial densities by means of the (Most probable Number). *Biometrics*. 6: 105- 115.
- Comstock, V.E. and Gulbeton, J.O. (1958): Methods of determining the oil of seed value of oil from small samples of flax. *Agron. J*. 50, 113.
- Coughlan, M. R. (1985): The properties of fungal and bacterial cellulases with comment on their production and application. *Biotechnol. Genet. Eng. Rev.*, 3: 39.
- Couresty (1921): *Journal of biological chemistry of* 49. P. 183.
- El- Borollosy, M.A and Refaat A.A. (1982). Phyllosphere inoculation with a symbiotic nitrogen fixing bacteria. *Res. Bull (1716) Fac. pf Agric., Ain Shams Univ., Cairo, Egypt*.
- El- Demerdash, M.E.; Abdel- Hafez, A.E. Mostafa; M. and Ishac, Y.Z. (1992). Response of wheat plants to inoculation with rhizobia and associative diazotrophs in the response of rock. Phosphate as a P. fertilizer. *Annals of Agricultural science Cairo* 37: 2, 379- 388.
- El-Gindy,A..A.. (1991) . Production of cellulase(s) by *Myriococcum albomyces* . *Zentralblatt-fur-Mikrobiologie.*, 146 (3) :193-196.

- El-Kina, G. Y. and Konstantinova, T. P. (1998).The effect of large applications of organic fertilizers on ameliorated pod zolic soils. *Agrokhimiya*. 5:68-75.
- El-Lateef, M.E. ; Selim, M.M. and Behairy, T.G. (1998). Response of some oil crops to biofertilization with phosphate dissolving bacteria. *Bulletin of the National Research Center..* 23(2): 193-202.
- El-Sersawy, M.M.; Abdel- Ghany, B.F.; Khalil, K.W. and Awadalla , S.Y (1997).. Interaction between organic manure mixtures, applied N-level and biofertilization on calcareous soil properties and wheat production in Wadi Sudr, South Sinai. *Egyptian – Journal of soil science*, 37: 3, 367-397
- El-Sharawy, M.A.O.; Aziz ,M..A.; Laila, A and Ali, K.M. (2003). Effect of the application of plant residues composts on some soil properties and yield of wheat and corn plants. *Egyptian Journal of soil science* . 43 (3): 421- 434.
- Folke, T. (2004). Degradation of cellulose . The Research of Folke Tjerneld's Group. Dept of Biochemistry , Lund university , Sweden. E-mail : Folke.Tjerneld@biokem .lu.se
- Gadgil, N.J.; Dagainawala, H.F.; Chakrabarti, T. and Khanna, P. (1995) .Enhanced cellulase production by a mutant of *Trichoderma reesei*. *Enzyme and Microbial Technology*. 17 (10): 942-946.
- Gallagher, J. ; Winters, A. ; Barron, N.; McHale, L. and McHale, A.P. (1996). Production of cellulase and beta-glucosidase activity during growth of the actinomycete *Micromonospora chalybeata* on cellulase-containing media. *Biotechnology-Letters*. 18: 5, 537-540 .

- Gomes, I.; Gomes, J.; Steiner, W. and Esterbauer, H. (1992) . Production of cellulase and xylanase by a wild strain of *Trichoderma viride* . *Applied-Microbiology-and-Biotechnology*. 36: 5, 701-707 .
- Griech, J. and Janicke, G. (1988): pot experiments comparing effects on yield of untreated household refuse and slurry with those of farmy and manure and mineral nitrogen. *Archiv- fur- Acke und pflanzenbau und Bodenkunde*. 32(3) : 197-203.
- Hassen, A ; Kaouala, B.; Jedidi, N.; Cherif, A.; Cherif, M. and Boudabous, A (2001) .Microbial characterization during composting of municipal solid waste. *Bioresource-Technology*. 80 (3): 217-225
- Hussein, H.F. and Radwan, S.M.A. (2002). Bio-orgnaic fertelization of potato under plastic mulches in relation to quality of production and associated weeds. *Arab University Journal of Agricultural Sciences*. 10(10) :287-309.
- Hussein, A.H.A. and Abdel-Nasser, G .(2001) Effect of different manure sources on some soil properties and sunflower plant growth. Part II. Growth, yield, leaf water contents and nutrients status. *Alexandria-Journal-of-Agricultural-Research*. 46(1): 205-225
- Indira, S.; Singh, N.G. and Singh, A.C. (2004) Studies on the nutrient content and quality assessment of compost enriched with biofertiliser. *Journal-of-Interacademia*. 8(4): 571-574
- Ivanitza, V.A.; Bagaeva, O.S.; Khodyrev, V.I. and Piven, T.V. (1990) , Method for testing the fungal resistance of material containing flax. *Mikologiya- I- fitopatologiya* .24(1): 44-50.

- Jackson, M.L. (1958): Soil Chemical Analysis. Prentice - Hall, Inc., England, Cliffs. 219- 221.
- Jah, K.; Khare, S.K. and Gandhi, A.P. (1995) .solid state fermentation of soy hull for the production of cellulase . Bioresource – Technology. 5 (3)321-322.
- Juange, T.C. (1993) : combined use of organic and chemical fertilizers. Extension Bulltein of food and fertilizer technology center. 371: 14-31.
- Kalogeris, E.; Iniotaki, F.; Topakas, E.; Christakopoulos, P.; Kekos, D. and Macris, B.J. (2003) . Performance of an intermittent agitation rotating drum type bioreactor for solid-state fermentation of wheat straw. Bioresource Technology. 86(3) : 207-213.
- Kanda, T.; Amano, Y.; Shiroishi, M. and Hoshino, E. (1994): Mode of action of exco- and endo - type cellulose from fungi in the hydrolysis of various substrates. Oyo Toshitsu Kagaku, 41 (2): 273- 282.
- Kang, H. W.; Nam, M.; Rhee, I.; Ko, J.; Park, K. B. and Kim, J. (1995). Journal of Agricultural Science – soil and fertilizer.
- Khatik, S.K. and Dikshit, P.R .(2001) Integrated use of organic manures and inorganic fertilizers on yield, quality, economics and nutrition of sunflower grown in Haplustert clay soil. Agricultural-Science-Digest. 21(2): 87-90 .
- Kim, T.I.; Jeong, K. H. ; Ham, J.S.; Yang, C.B.; Chung, I.B.; Kim, M.K. and Kim, K.N. (2004) Isolation and characterization of cellulase secreting bacterium from cattle manure: application to composting. Compost-Science-and-Utilization.; 12(3): 242-248

- Kluge, R.; Mokry, M. ; Timmermann. F and Braun. C (1997). Effect of composts on the yield and heavy metal content in soil and plant. VDLVFA- congresses leipzig, 6(7): 650.
- Kumari, C.P.; lachanna, A.; and Satyanarayana, V. (2004). Effect of biofertilizers on seed yield and quality of sun flower, *Helianthus annuus*. Journal of Oil seeds- Research. 21 (1): 183- 184.
- Lee, N.E. ; Lima, M. and Woodward, J. (1988); Hydrolysis of cellulose by a mixture of *Trichoderma reesei* cellobiohydrolase and *Aspergillus niger* endoglucanase. Biochem. Biophys. Acta. 967.
- Li, G. ; Zhang, Z.; Bai, Y.; Li, G.; Zhang, Z. and Bai, Y. (1995). The different effects of HTC and WC on C,N transformation and pathogenic bacterial population. Acta- Agriculture- Universitatis- pekinensis . 21: 3, 286-290.
- Limtong, P.; Vangnai, S. ; Leaungvutivirog, C. ; piriyaprin, S.; Sunan, T.V. and Thunyadee, P. (1994). Special – publication – Taichung-District- Agricultural- Improvement station. 35, 905-911.
- Limtong, P.; Vangnai, S. ; Sunathapongsuk, V. and Piriyaprin, S. (1990). Isolation and selection of thermophilic cellulolytic microorganisms for compost production in Thailand. Kasetsart - Journal .24: 1, 108-115.
- Li, R.; Xi,Y.; Yingo, C.Z. ; Han, W.; Liu, J. Q. ; LiRQ, X.; Cao, Z.; Han, W.H. and Liu, J.Q. (2002). Complex use of wastes containing cellulose .China – Environmental- Science, 22:1, 24-27.

- Loecke, T.D, libeman, M., Cambardella , C.A. and Richard, T.L. (2004): Corn response to composting and time of application of solid swine manure. *Agronomy. Journal* 20, 96 (1): 214-233.
- Lu,W.J.; Wang, H.T.; Nie, Y.F.; Wang, Z.C.; Huang, D.Y.; Qiu, X.Y. and Chen, J.C. (2004) .Effect of inoculating flower stalks and vegetable waste with ligno-cellulolytic microorganisms on the composting process. *Journal-of-Environmental-Science-and-Health-Part-B,-Pesticides,-Food-Contaminants,-and-Agricultural-Wastes*. 39(5/6): 871-887
- Maccheroni, J.W.; Araujo, W.L. and Azevedo, J. L. (2004) . Ambient pH-regulated enzyme secretion in endophytic and pathogenic isolates of the fungal genus *Colletotrichum* . *Scientia-Agricola* . 61(3): 298-302
- Mackie,T. J. and McCartney, J.E. .(1956). *Hand Book of Practical Bacteriology* 9th ed .
- Mandels, M.; Andreotti, R. and Roche, C. (1976): Measurmeent of saccharifying cellulose. In *Enzymatic conversion of cellulosic materials : technology and applications*. Edited by E.L. Gaden, M.C. Mandlels, E.T. Reese and L.A. Spano. John Wiley & Sons, New York. pp. 21- 33.
- Manna, M.C.; Ghosh, P.K and Ganguly, T.K. (2003) . Comparative performance of four source of enriched phosphocompost and inorganic fertilizer application on yield, uptake of nutrients and biological activity of soil under soybean- wheat- rotation. *Journal of food Agriculture and environment*. 1 (2): 203-208.

- Mekki, B.B.; Selim, M.M. and Saber, M.S.M.. (1999).. Utilization of biofertilizers in field crop production. Egyptian- Journal of Agronomy 199 (21) : 113-124.
- Mihalache G. (1999). Probleme- de- Agrofitotehnie- Teoretica- Si- Aplicata, . The influence of treatment with atrazine on cellulose degradation in soil. 21(1-2): 87-91.
- Miller, G.L. (1959): Use of dinitrosally cyclic acid reagent for determination of reducing sugars Anal, Chem., 31, 426- 428.
- Muniswaran, P.K.A. and Charyulu, N. (1994). Solid state fermentation of coconut coirpith for cellulase production. Enzyme and Microbial Technology. 16(5): 436-440 .
- Nakasaki, K. ; Yaguchi, H. ; Saski,Y. and Kubota, H. (1991). Effects of C/N ratio on thermophilic composting of garbage. Journal-of-Fermentation-and-Bioengineering. 1991, 73: 1, 43-45
- Nandi, N.; Hajra, J. N.and Sinha, N.B. (1996). Microbial synthesis of humus from rice straw following two- step composition process. Journal of the Indian society of soil science. 44: 3, 413-416.
- Omar, S.A. and Abdalla, M.H . (2000) .Physiological aspects of fungi isolated from root nodules of faba bean (*Vicia faba* L.). Microbiological-Research. 154(4): 339-347.
- Oxoid (1982). Oxoid Manual of Culture Media Ingredient and other Laboratory Surface, Turner graphic Ltd. England fifth. ed.
- Parameter, J.R. Jr. and H.S. Whitney (1970): Taxonomy and Nomenclature of the imperfect state In: *Rhizoctonia solani*. Biology and Pathology. Parameter. J.R. (Ed.) Univ. of California Press. USA: 7- 19.

- Pardo, A.G. and Forchiassin, F. (1999) . Influence of temperature and pH on cellulase activity and stability in *Nectria catalinensis*. *Revista-Argentina-de-Microbiologia*. 31(1): 31-35.
- Parkinson, R.J.; Fuller, M.P and Groenhof, A.C (1999): An evaluation of greenwast compost for the production of fograge maize (zea mays) . *Compost science and vtilization*. 7(1): 72-80.
- Pera, A.; Vallini, G.; Sireno, I.; Bianchin, M.L ; and Bertoldi, M.D. (1983) . Effect of organic matter on rhizosphere microorganisms and root development of sorghum plants in two different soils. *Plant-and-Soil*. 74(1): 3-18
- Ponce, N. T. and De la Torre, M. (2001). Regulation of cellulases and xylanases from a mutant of *Cellulomonas flavigena* growing on sugar-cane bagasse in continuous culture. *Bioresource Technology*. 78(3): 285-291.
- Prasetran, P. and Doelle, H.W. (1987) . Nutrient optimization of cellulase biosynthesis by a newly isolated cellulomonas sp. *MIRCEN – Journal – of – Applied- Microbiology and Biotechnology*, 3 (1): 33-44 .
- Radia, K.M.; Tahir Abbas, F.R.; Usmani , N.F. and Hussain. S.S . (1988) The effect of cotton waste compost, zarkhez and cow dung manure on sunflower yield. *Paksitan- Journal of scientific and industrial Res*. 31 (7) : 512- 515
- Radwan, S.M.A.; Hussein, H.F.; Rubio, J.L.; Morgan, R.P.C.; Asins, S. and Andreu, V. (2002). Man and soil at the third Millennium, proceedings International congress of the European society for soil conversation, Valencia, spain, 2002, 1015-1023.

- Radwans, S.M.A.; Saber, M.; Ana, C. and Martinprevel, P. (1999): Response of corn to bio and organic fertilizers in a newly reclaimed sand Soil. Improved crop quality by nutrient management. 253- 257.
- Ramadan, H.M.; Koreish, E.A.; Gaber, H.M. and El-Fayoumy, M.E. (2002). Assessment and comparison of bio and mineral fertilization of farm profitability in different newly- reclaimed soils. Alexandria Journal of Agricultural Research. 47: 1, 133-146.
- Reese, E.T.; Situ, R.G.H. and Levinson, H.S. (1950): The biological degradation of soluble cellulose derivatives and its relationship to the mechanism of cellulose hydrolysis. J. Bacteriol., 59: 458- 497.
- Robson, L.M. and Chambliss, G.H. (1984): Characterization of the cellulolytic activity of a *Bacillus* isolate. Appl. Environ. Microbiol. 47: 1039- 1046.
- Saha, B.C. (2004). Production, purification and properties of endoglucanase from a newly isolated strain of *Mucro circinelloides*. Process, Biochemistry. 39 (12): 1871-1876.
- Sandark, N.A. (1977). Cellulose decomposition by *Micromonospora*. Mikrobiolgiya, 46: 478- 481.
- Sarkar, A. and Varma, A. (1988): Influence of cellulolytic organisms associated with termite, *Odontotermes obesus* on carbon mobility in a semiarid ecosystem. Arid- Soil- Research and Rehabilitation 2 (2): 75- 84.

- Sarkar, A.; Paul, J.; Saxena, S. and Varma, A . (1985). Microbes and cellulose biotechnology. Indian-Journal-of-Agricultural-Chemistry.; 18(2): 221-238
- Sarojin, K. and Mathur, R.S. (1994). Biodegradation of paddy straw with cellulolytic fungi and its application on wheat crop . Bioresource – Technology. 47:2, 185:188.
- Schaffner, D.W. and Toledo, R.T. (1991). Cellulase production by *Trichoderma reesei* when cultured on xylose-based media supplemented with sorbose. Biotechnology-and-Bioengineering. 37(1): 12-16.
- Schitzer, M. (1982) Extraction, fractionation and characterization of soil organic matter. Methods of soil analysis, part.2. Chemical and microbiological properties, 2.ed. SSSA, Madison,. p.539-579
- Selby, K. and Matiland, C.C. (1967): The cellulose of *Trichoderma viride*: Separation of the components involved in the solubilization of cotton. Biochem., J. 104: 716- 724.
- Sharma, N. and Bhalla, T.C .(1992) . Bacteria from forest litter. Indian-Journal-of-Microbiology., 32: 2, 213-215.
- Sharma S.K.; Kalra., K.L. and Grewal, H.S. (2002) .Fermentation of enzymatically saccharified sunflower stalks for ethanol production and its scale up.Bioresource-Technology. 85(1): 31-33.
- Shash, S.M. (2002): Environmental nutritiaonl parameters . controlling the prdocutivity of cellulase (s) by *Asperigillus flavus* S-7 attacking water hyacinth Egyptian. Journal of Microbiology. 37 (2): 101-115.

- Shindi, A.A. (1995). Studies on pectin-degrading fungi in compost. *Egyptian-Journal-of-Microbiology* 30(1): 85-99.
- Sinha, R.K.; Sunil, H. and Heart, S. (2002). A cost- effective microbial slurry technology for rapid composting municipal solid wastes in waste dump sites in India and its feasibility for use in Australia. *Environmentalist* . 22: 1, 9-12.
- Snedecor, G. W. (1966): Statical methods. The state Univ. Press. Ames, Iowa U.S.A. PP. 534.
- Spiridonov, N.A. and Wilson, D.B. (1998). Regulation of biosynthesis of individual cellulases in *Thermomonospora fusca*. *Journal of Bacteriology*. 180 (4): 3529-3532.
- Suzuki, T; Ikumi, Y.; Okamoto, S.; Watanabe, I.; Fujitake, N. and Otsuka, H. (2004) . Aerobic compositing of chips from clear . cut trees with various co materials. *Bioresource- Technology*, 95 (2): 121-128.
- Taj, A.S. and Alkenany, K.I. (1993) . Properties of the cellulolytic system from *Aspergillus niveus* .*Mycological-Research*. 97: 1, 15-22
- Tepper, E.Z.; Sidoren, K.D and Mohamed, A. (1987). *Biology – Bulltein*, 8, 1: 82-90. Rice straw as source of humus and microflora involved in its transformation.
- Timinon, A. (1940): The interaction of higher plants and soil microorganisms. II. study of the microbiological population of the rhizosphere in relation to resistance of plants to soil - borne diseases. *Can. J. Res. B.*, 18 (444- 456).
- Tripathi, B.P. (1997): Report on third cycle maize and wheat synchrony in maize -wheat system. Working paper -Lume- Agrciutylrual Research center. 60-97 .

- Tweddle , R.J.; Marshall, J. and Jabaji, H.S. (1996). Endo- 1,4- beta. glucanase production by *stachybotrys elegans*, a mycoparasite of *Rhizoctonia solan*. *Mycologia* , 88 (3): 410-415.
- Urmila, G. ; Roma, K.; Phutela, R.P.; Gupta, U. and Kalra, R. (1996). Factors affecting cellulase production in *Volvariella*, the straw mushroom. *Mushroom-Research*. 1996, 5: 1, 29-32.
- Valino, E.; Medina, I. and Dustet, J. (2000) .Characterization and optimization of the exo beta 1-4 glucanase enzyme of the *Aspergillus fumigatus* 155 v cellulolytic fungus strain. *Cuban-Journal-of-Agricultural-Science*.
- Velasco, V.J.; Figueroa, S.B.; Ferrera, C.R.; Trinidad, S.A.; and Gallegos S. J. (2004) .CO₂ and microbial population dynamics in manure and straw compost under aeration. *Terra*. 22(3): 307-316
- Verma, L.N.; Rawat, A.K.; Dubey, S.B. and Rathore, G.S.(1999). Variation in temperature and shrinkage of *Trichoderma harzianum*-inoculated compost heaps and quality of compost prepared from crop and poultry wastes: *Indian-Journal-of-Agricultural-Sciences*. 69(5): 340-344
- Vorogen, A.G.J.; Beldman, G. and Rombouts, F.M. (1988): Cellulases of a mutant strain of *Trichoderma viride*. *Methods Enzymol.*, 160: 243- 251.
- Waller, R.A. and Duncan, B.D. (1969): Away for the symoretric multiple comparison Problem. *Amer. Stat. Assoc. J.* 3, 1485- 1503.
- Wang, Q. and Zhao, X. (1994): Chemical composition of cellulose from *Aspergillus niger*. *Weishengwuxue Zazhi*, 14 (1): 6- 8.

- Waszczynskyj, N.; Feitosa, F.R.F.; Hishida, D.O. and Nascimento, K.H (1990) . Production of cellulolytic enzymes by *Pycnoporus cinnabarinus* . *Arquivos-de-Biologia-e-Tecnologia*. 33 (1): 215-225.
- Watanabe, F.S. and Olsen, N. (1965) .Test of an ascorbic acid method for determining phosphorus in soil, water and NaHCO₃ extracts from soil. *Soil Sci. Soc. Amer. Proc.*, 29: 677- 678.
- Watanabe, N.; Kamei, S.; Onkubo, A., Yamanaka, M. and Oshawa, S. (1986): Assay of Protein by Pyrogallolred reagent . *Clin. Chemistry*, 32; 1551-1553
- Windle, T. E. (1958): *The Examination of Waters and Water supplies* 7th ed., Church; Ltc. London.
- Wood, T.M. and McCrae, S. I. (1972): The purification and properties of the CI component of *Trichoderma koningii* cellulose. *Biochem. J.*, 128: 1183- 1192.
- Woodward, J.; Lima, M. and Lee, N.E. (1988): Hydrolysis of cellulose by a mixture of *Trichoderma ressei* cellobiohydrolase and *Aspergillus niger* endoglucanase. *Biochem. J.*, 225: 895- 899.
- Xia, L.; Yu, S. and Cheng, Z. (1993). Cellulase production by immobilized *Trichoderma ressei* in fed- batch fermentation. *Linchan Huaxue Yu Gongye*, 13 (3): 217- 222.
- Yamaguchi, H. and Yoshino, K. (2001). Influence of tannin-copper complexes as preservatives for wood on mechanism of decomposition by brown-rot fungus *Fomitopsis palustris*. *Holzforschung*-. 55(5): 464-470

- Ye, G.S. and Fields, M.L. (1989) . Cellulolytic enzyme production by three fungi grown in a ground corn cob medium. *Journal-of-Food-Protection*. 1989, 52: 4, 248-251 .
- Zhang, Y. ; LuXue, B.; Wan, X.; Dan, H.; Wang, H. and Sun, Y. (2005) Isolation of a cellulose-decomposing microorganisms and investigation of its crude enzyme properties. *Journal-of-South-China-Agricultural-University*. 26(2): 69-72
- Zhu-N.W.; Deng, C.Y.; Xiong, Y.Z. and Qian, H.Y. (2004) .Performance characteristics of three aeration systems in the swine manure composting. *Bioresource-Technology*. 95(3): 319-326