

References

LITERATURE CITED

Abott, W. S. 1925. A method of computing the effectiveness of an insecticide. J. Econ. Entomol., 18: 265-267.

Abou-Bakr, H.; T. A. El-Kifl; and M. F. Haidar 1989. laboratory Evaluation of certain Entomogenous nematodes against the hairy beetle Tropinota Squalida scop. Adults (Coleoptera- Scarabaeidae). Proc. 1st.Int. Conf. Econ. Ent. Vol. II: 47.

Ahmad, S.; H. T. Stren; and L. M. vasvary 1983. The japanese beetle: a major pest of turfgrass, Am. lawn Appl., Vol. 4, No. 2: 2.

Alfieri, A. 1976. Coleoptera of Egypt. Mem. Soc. Ent. Egypte vol. 5: 361 pp.

Ali, M. A. and I. L. Ibrahim 1988. Preliminary study on occurance and abundance of Tropinota squalida Scop. (coleoptera-scara-baeidae) infesting cartain ornamental plants and fruit crops. Al-Azhar J. Agric. Ras., 9: 1-14.

Bishara, L.S. (1958). Comparative Morphology of some Dynastid larvae injurious to agriculture in Egypt. Bull.Soc. entomol. Egypte XIII, 501-513.

Brandt, C. R.; M. J. Adang and K. D. Spence. 1977. The peritrophic membrane: ultrastructure analysis and function as a mechanical barrier to microbial infection in Prqyia pseudotsugata. J. Invertebr. Pathol. 3.

Byers, J. A. and G. O. poinar, Jr. 1982. Location of insect hosts by the nematode, Neoaplectona carpocapsae, in response to temperature. Behavior, 79: 1-10.

Comprage, D. 1977. Pests attacking the underground parts of field crops. Stetocine podzenih organa ratarskin Kultura Novi Sad, Yugoslavia, Institut za Zastitu Bilja 1 - 210.

Croll, N. A. and B. E. Matthews. 1977. Survival of nematodes: in "biology of nematodes", pp. 152 - 165. Wiley and sons, New York.

Donalson, J. M. I. 1979. Laboratory rearing technique for pachnoda sinuata Flaventriss (Coleoptera-Scarabaeidae) and other cetoniid larvae. I. Ent Sth Afr. 42 (1): 137-147.

Douce, G. K. and R. M. Mc pherson [eds.] 1989. Summary of losses from insect damage and costs of control in Georgia, 1988. Univ. of Georgia College of Agriculture Experiment Station special publication No. 64.

El-Kifl, T. H. 1984. Factors affecting potentialities of Entomogenous nematodes in the Biological control of insect pests. Ph. D. thesis, Fac. Agric., Cairo Univ. 160 pp.

El-Kifl, T. H. and M. A. El-sherif 1986.
Dispersal pontentiality of the infective
juveniles of the Entomogenous nematodes,
Neoaplectana carpocapsae. Bull. Fac. Agric.,
Univ. of Cairo, 37 (2): 1085-1093

Forschler, B. T. and W. A. Gardner 1991a.
Concentration-Mortality response of
phyllophaga hirticula (Coleoptera-
scarabaeidae) to three entomogenous nematodes.
J. Econ. Ento-mol. 84 (31): 841-843.

Forschler, B. T. and W. A. Gardner 1991b.
Field efficacy and persistence of entomogenous
nematodes in the management of white grubs
(Coleoptera-Scarabaeidae) in the turf and
pasture. J. Econ. Entomol. 84: 1454-1459.

Forschler, B. T., and W. A. Gardner 1991c.
Parasitism of Phyllophaga hirticula
(Coleoptera: Scarabaeidae) by Heterorhabditis
heliothidis and Steinernema carpocapsae. J.
Invertebr. Pathol., 58: 396-407.

Garcia-vidal, M. 1984. The genus Phyllophaga Harris 1826 (Coleoptera-Scarabaeidae) in Cuba. II Description of eleven new species and illustration of female genitalia of twelve other Phyllophaga pan-pacific Entomologist 60 (4): 313-325.

Gaugler, R. 1981. The biological control potential of neoaplectanid nematodes. J. Nematol. 13: 241-249.

Gaugler, R. 1988. Ecological considerations in biological control of soil-inhabiting insects with entomopathogenic nematodes, Agric. Ecosys. Environ., 24, 351.

Gaugler, R. and H. K. Kaya [eds]. 1990. Entomopathogenic nematodes in biological control. CRC press, Boca Raton Fl, USA.

Georgis, R. and G. O. Poinar, Jr., 1983a.
Effect of soil texture on the distribution and
infectivity of Neoaplectana carpocapsae
(Nematoda: Steinernematidae). J. Nematol., 15,
219.

Georgis, R. and G. O. Poinar, Jr. 1983b. Effect
of soil texture on the distribution and
infectivity of Neoaplectana glaseri (Nematoda:
Steinernematidae) in sandy loam soil. J.
Nematol., 15, 652.

Georgis, R., and G. O. Poinar, Jr. 1983c.
Vertical migration of Heterorhabditis
bacteriophora and H. heliothidis (Nematoda :
Heterorhabditidae) in sandy loam soil. J.
Nematol., 15, 652.

Georgis, R., and G. O. Poinar, Jr. 1983D.
Effect of soil texture on the distribution and
infectivity of Neoaplectana carpocapsae
(Nematoda: Steinernematidae). J. Nematol. 15:
308-311.

Glazer, I. and A. Gol'Berg 1989. Laboratory evaluation of steinernematid and Heterorhabditid nematodes for control of the beetle Maladera matrida. phytoparasitica 17 (1): 3-12

Hale, P. R. and J. T. Hale 1975. Insect pests of rice in the New Guinea Islands. Rice Ent. Newsletter No. 3: 4.

Heller, H. R. and S. Kellogg 1989. Fall control of japanese beetles. Insectic. Acaricide tests 14: 313-314

Ishibashi, N. and E. Kondo. 1986. Steinernema feltiae (DD-136) and S. glaseri persistence in soil and bark compost and their influence on native nematodes. J. Nematol., 18: 310.

Kamel, A. S. 1988. Biology and morphology of Pachnoda fasciata. M. Sc. thesis, Fac. Scien. Zag. Univ., Benha branch.

Kard, B. M. R.; F. P. Hain and W. M. Brooks
1988. Field suppression of three white grub
species (Coleoptera: Scarabaeidae) by the
entomogenous nematodes S. feliae and H.
heliethidis J. Econ. Entomol. 81: 1033-1039.

Kaya, H. K. 1985. Entomogenous nematodes for
insect control in IPM system, In (M. A. Hoy
and D. C. Herzog, eds.) Biological control in
agricultural IPM system. Academic press, New
York, PP. 283-302.

Kaya, H. K.; C. Mannion; T. M. Burlando and C.
E. Nelson. 1987. Escape of Steinernema feltiae
from alginate capsules containing tomato
seeds. J. Nematol., 19: 287.

Kondo, E. and N. Ishibashi. 1989. Non-oral
infection of Steinernema feltiae (DD-136) to
the common cutworm, Spodoptera litura
(Lepidoptera: Noctuidae). Appl. Entom. Zool.,
19: 85-95.

Kung, S. P.; R. Gaugler and H. K. Kaya. 1990a. Soil type and entomopathogenic nematode persistence. J. Invertebr. Pathol., 55: 401 - 406.

Kung, S. P.; R. Gaugler and H. K. Kaya. 1990b. Effect of soil temperature, moisture and relative humidity on entomopathogenic nematode persistence. In R. Gaugler and H. K. Kaya [eds.]. Entomopathogenic nematodes in biological control. CRC press, Boca Raton, Fl., USA pp. 101 - 111.

Mamiya, Y. 1989. Heterorhabditid nematodes for three different insects. Appl. Entomol. & Zool. 24 (3) : 302-308.

Molyneux, A. S. 1985. Survival of infective juveniles of Heterorhabditis spp. and Steinernema spp. (Nematoda: Rhabditida) at various temperature and their subsequent infectivity for insects. Rev. Nematol., 8: 165.

Molyneux, A. S. and R. A. Bedding 1984. Influence of soil texture and moisture on the infectivity of Heterorhabditis sp. DI and Steinernema glaseri for larvae of the sheep blowfly Lucilia cuprina Nematologica 30: 358-365.

Moyle, P. L. and H. K. Kaya 1981. Dispersal and infectivity of the entomogenous nematode Neoaplectana carpocapsae Weiser (Rhabditida: Steinernematidae) in sand. J. Nematol. 13. 295-300.

Nabil, M. A. 1982. Comparative studies on the effect of pesticides and the entomogenous nematodes in controlling some economic insects. Ph. D. thesis, Fac. Agric., Cairo Univ. 157 pp.

Nguyen, K. B. and G. C. Smart 1990. Vertical dispersal of Steinernema scapterisci. Journal of Nematology 22 (4): 574-578.

Nickle, W. R. 1963. Observation on the effect of nematodes on Ips confusus (Leconte) and other bark beetle. Ibid. 5: 386-389.

Passlow T. 1969. Insect pests of peanuts in southern Queensland. Qd. Agric. J. 95 (7): 449-451.

Poinar, G. O., Jr. 1971. Use of the nematodes for microbial control of insects. PP 181-203 in H. D. Burges and H.W. Hussey, eds. Microbial control of insects and mites. New York: Academic press.

Poinar, G. O., Jr. 1978. Generation polymorphism in Neocaplectana glaseri Steiner (Steinernematidae: Nematoda) described from Strigoderma arboricola (Fab.) (Scarabaeidae: Coleoptera) in Northern California. Nematologica 24: 105 - 114.

Poinar, G. O., Jr. and G. M. Thomas 1966.
Significance of Achromobacter nematophilus
Poinar and Thomas (Achromobacteriaceae:
Eubacteriales) in the development of the
nematode DD-136 (Neoaplectana sp.,
Steinernematidae). Parasitology 56: 385-390.

Poinar, G. O., Jr.; G. M. Thomas; G. V.
Veremchur and D. E. Pinnock 1971. Further
characterization of Achromobacter nematophilus
from American and Soviet population of the
nematode Neoaplectana carpocapsae Weiser.
International Journal of Systematic
Bacteriology 21: 78-82.

Reed, E. M. and P. B. Carne 1967. The
suitability of a nematode (DD-136) for the
control of some pasture insects. J.
Invertebrate Pathol. 9: 196-204.

Schroeder, W. J. and J. B. Beavers 1987.
Movement of the Entomogenous nematodes of the
families Heterorhabditidae and
Steinernematidae in soil. J. Nematol., 19,
257.

Shetlar, J. D., D. E. Suleman and R. Georgis
1988. Irrigation and use of Entomogenous
nematodes, Neoaplectana spp. and
Heterorhabditis heliothidis (Rhabditida:
Steinernematidae), for control of japanese
beetle (Coleoptera: Scarabaeidae) grubs in
turfgrass. J. Econ. Entomol. 81: 1318-1322.

Simoes, N.; C. Laumond and F. Bonifassi, F.
1993. Effectiveness of Steinernema spp. and
Heterorhabditis bacteriophora against Popillia
japonica in the Azores. J. Nematol. 25 (3):
480 - 485.

Simons, W. R. and G. O. Poinar, Jr. 1973. The ability of Neoaplectana carpocapsae (Steinernematidae: Nematoda) to survive extended periods of dessication. J. Invertebr. Pathol. 22: 228-230.

Sosa O., Jr. and Beavers S. B. 1985. Entomogenous nematodes as biological control organisms for ligyrus subtropicus (Coleoptera: Scarabaeidae) in sugar-cane Environ. Entomol. 14: 80-82.

Sosa, O., jr. and D. G. Hal 1989. Mortality of Ligyrus subtropicus (Coleoptera: Scarabaeidae) by entomogenous nematodes in field and laboratory trials, J. Econ. Entomol. 82, 740.

Steven, R. A.; Y. Tamson; L. H. Janes and G. Ramon 1992. Biological control of japanese, oriental and Black turfgrass Ataenius beetle (Coleoptera-Scarabaeidae) larvae with Entomogenous nematodes (Nematoda-Steinernematidae, Heterorhabditidae). J. Econ. Entomol. 85 (5): 1660-1665.

Tamson, Y. and R. A. Steven 1992. Effect of entomopathogenic nematode species, rate, soil, moisture and bacteria on control of Japanese beetle (Coleoptera: Scarabaeidae) larvae in the laboratory. J. Econ. Entomol. 85 (6): 2144-2148.

Tashiro, H. 1987. Turfgrass insects of the U.S. and Canada, Cornell University Press, Ithaca, NY.

Thomas, G. M. and G. O. Poinar, Jr. 1979. Xenorhabdus gen. nov., a genus of entomopathogenic nematophilic bacteria of the family: Enterobacteriaceae. International Journal of Systematic Bacteriology. 29: 352-360.

Villani, M. G. and R. J. Wright 1988. Entomogenous nematodes as biological control agents of European chafer and Japanese beetle (Coleoptera-Scarabaeidae) larvae infesting turfgrass J. Econ. Entomol. 81: 484-487.

Wang, J. X. and L. Y. Li 1987. Entomogenous nematode research in China, Rev. Nematol., 10: 483.

White, G. F. 1927. A method for obtaining infective nematode larvae from cultures. Science 66: 302-303.

Wood, B. J.; R. H. V. Corley and K. H. Goh 1973. Studies on the effect of pest damage on oil palm yield. Communication, chemora research station No. 58: 20 pp.