

REFERENCES

Abd El-Gawad, K. H.; Farghal, A. I.; Salman, A. G. A. and Ali, M. K. (1987).

Survey and population density of rodent species in some urban, rural and cultivated areas in Sohag Governorate, Upper Egypt. *Assiut J. Agric. Sci.*, 18(1): 259-266.

Abd El-Gawad, K. H.; Salit, A. M. and Ali, A. M. (1982).

Population density of rodent species in agricultural and semi-arid areas. *Assiut J. Agric. Sci.*, 13: 27-37.

Abdel Aziz, H. A. (2004).

Ecological studies of phlebotomine sand flies (Diptera: Psychodidae) in Southern Sinai. *Ph. D. Thesis, Fac. Sci., Ain Shams Univ. Cairo, A.R.E.*

Aguiar, G. M.; Vilela, M. L.; Schubach, P. D.; Soucasaux, T. and Azevedo, A. C. R. (1985).

Aspectos da ecologia dos flebotomos de Parque Nacional da Serra dos Orgaos. Rio de Janeiro. IV- Frequencia mensal em armadilhas luminosas (Diptera: Psychodidae: Phlebotominae). *Mem. Inst. Oswaldo Cruz.*, 80: 465-482.

Aldawood, A. S.; Alahmed, A. M.; Kheir, S. M. and Hussein, S. M. (2004).

Population dynamics of sand flies (Diptera: Psychodidae) in Hanifah Valley Riyadh, Saudi Arabia. *Pakistan J. Bio. Sci.*, 7(4): 464-467.

Alexander, J.B. (1987).

Dispersal of phlebotomine sandflies (Diptera: Psychodidae) in a Colombia coffee plantation. *J. Med. Entomol.*, 4: 552-558.

Alptekin, D.; Kasap, M.; Luleyap, U.; Kasap, H.; Aksoy, S. and Wilson, M. L. (1999).

Sand flies (Diptera: Psychodidae) associated with epidemic cutaneous leishmaniasis in Sanliurfa, Turkey. *J. Med. Entomol.*, 36(3): 277-281.

Alves, J. C. M.; Pacheco, R. S. F. and Brazil, R. P. (2001).

Biological aspects of *Lutzomyia renei* (Martins, Falcao & Silva, 1957) (Diptera: Psychodidae) under laboratory and field conditions. *Rev. Brasil. Entomol.*, 45(1): 17-21.

Aly, M. Z. (1996).

Effect of moon light and lunar periodicity on the light trap catches of the order Lepidoptera at two different areas in Upper Egypt (Qena). *J. Egypt. Ger. Soc. Zool.*, 21(E): 297-309.

Ashford, R.W. (2001).

Phlebotomus fevers. *The Encyclopedia of Arthropod-transmitted Infection* (ed. By M.W. Service), pp. 104-106. CABI publishing, Wallingford, U.K.

Azevedo, A. C.; Vilela, M. L.; Souza, N. A.; Andrade-Coelho, C. A.; Barbosa, A. F.; Firmo, A. L. and Rangel, E. F. (1996).

The sand fly fauna (Diptera: Psychodidae:Phlebotominae) of focus of cutaneous leishmaniasis in Ilheus, State of Bahia Brazil. *Mem. Inst. Oswaldo Cruz.*, 91(1): 75-79.

Barata, R. A.; da Silva, J. C. F.; da Costa, R. T.; Fortes-Dias, C. L.; da Silva, J. C; de Paula, E. V.; Prata, A.; Monterio, E. M. and Dias, E. S. (2004).

Phlebotomine sand flies in Porteirinha, an area of American visceral leishmaniasis transmission in the state of Minas Gerais. *Mem. Inst. Oswaldo Cruz.*, 99(5): 481-487.

Barros, V. L.; Rebelo, J. M. M. and Silva, F. S. (2000).

Sand flies (Diptera: Psychodidae) in a secondary forest area in the county Paco do lumiar, Maronhao Brazil: a leishmaniasis transmission area. *Cad. Saude publica.*, 16(1): 265-270.

Bidlingmayer, W. I. (1974).

The influence of environmental factors and physiological stage on flight patterns of mosquitoes taken on vehicle

aspirator and truck, suction, bolt and New Jersey light traps.
J. Med. Entomol., 15: 119-146.

Birtles, R. J. (2001).

Carrion's disease. *The encyclopedia of Arthropod-transmitted Infection* (ed. By M.W. Service), pp.104-106. CABI Publishing, Wallingford, U.K.

Bishop, A. L.; Mckenzie, H. J.; Barchia, I. M. and Spohr, L. J. (2000).

Moon phase and other factors affecting light-trap catches of *Culicoides brevitarsis* Kieffer (Diptera: Ceratopogonidae).
Austr. J. Entomol., 39(1): 29-32.

Bowden, J. (1973).

The influence of moonlight on catches of insects in light-traps in Africa. Part 1. *Bull. Entomol. Res.*, 63: 115-128.

Brinson, F. J.; Hagan, D. V.; Comer, J. A. and Strohlein, D. A. (1992).

Seasonal abundance of *Lutzomyia shannani* (Diptera: Psychodidae) on Ossabaw Island, Georgia. *J. Med. Entomol.*, 29(2): 178-182.

Carrasco, J.; Morrision, A. and Ponce, C. (1998).

Behavior of *Lutzomyia longipalpis* in an area of southern Honduras endemic for visceral typical cutaneous leishmaniasis. *Ann. Trop. Med. Parasitol.*, 92(8): 867-876.

Carvalho, M. L.; Rebelo, J. M. M.; Arouje, J. A. C.; Barros, V. L. L. and de Araujo, J. A. C. (2000a).

Ecological study of sand flies (Diptera: Psychodidae) in the municipality of Sao Jose de Ribamar, Sao Luis Island MA, Brazil. Endemic area of leishmaniasis. *Entomol. Vectors*, 7(1): 19-32.

Carvalho, R. W.; Souza, M. B.; Serra-Freire, N. M.; Pontes. C. S.; Conceicao, N. F.; Almeida, A. B.; Campos, V. S.; de-Carvalho, R. W.; de-Souza, M.B.; da-Conceicao, N. F. and de-Almeida, A. B. (2000b).

Sand flies (Diptera: Psychodidae) of Araujo Island, municipal district of Paraty, Rio de Janeiro State, Brazil. 2-Behavior in relation to time of day and the lunar cycle. *Entomol. Vectores*, 7(2): 143-153.

Chaniotis, B. N.; Correa, M. A.; Tesh, R. B. and Johnson, K. M. (1974).

Horizontal and vertical movement of phlebotomine sand flies in a Panamanian rain forest. *J. Med. Entomol.*, 11: 369-375.

Chaniotis, B. N.; Tesh, R. B.; Correa, M. A. and Johnson, K. M. (1972).

Diurnal resting sites of phlebotomine sand flies in a Panamanian tropic forest. *J. Med. Entomol.*, 9: 91-98.

Charlwood, J. O.; Lopes, J. and Whalley, P.C. (1982)

Light intensity measurement and the biting behavior of some sylvatic mosquitoes of the Amazon basin (Diptera: Culicidae). *Acta Amazonica*, 12: 61-64.

Clements, A. N. (1999).

The biology of the mosquitoes. Vol. 2 *Sensory Reception and Behavior*, CABI publishing, Oxon, 740 pp.

Costantini, C.; Sagnon, N. F.; Sanogo, E.; Merzagora, L. and Coluzzi, M. (1998).

Relationship to human biting collection and influence of light and bednet in CDC light-trap catches of West African malaria vector. *Bull. Entomol. Res.*, 88: 503-511.

Croset, H.; Rioux, J. A.; Maistre, M. and Bayer, N. (1978).

Les phlebotomes de Tunisie (Diptera: Phlebotomidae) mise au point systematique, chorologie et ethologique. *Ann. Parasitol. Hum. Comp.*, 53: 711-749.

Danthanarayana, W. (1976).

Diel and lunar flight periodicities in the light brown apple moth *Epiphyas postvittana* Tortricidae and their possible adaptive significance. *Austr. J. Zool.*, 24(1): 65-73.

Darwish, A. B. (2004).

Ecological studies of leishmaniasis vectors and the reservoir host(s) in some areas in Suez Governorate. *M. Sc. Thesis, Zool. Dept., Fac. Sci., Suez Canal Univ., Ismailia, Egypt.*

Das, S. B. and Katyar, N. (2001).

Effect of moon light and lunar periodicity on the pheromone trap catches of tobacco caterpillar *Spodoptera litura* moths. *Crop Res. (Hisar)*, 21(2): 229-236.

De Queiroz, R. G.; Vasconcelos, A. W.; Vasconcelos, I. A.; De Sousa, R. N.; Pessoa, F. A.; De Alencar, J. E. and David, J. R. (1991).

Phlebotomine sandfly (Diptera: Psychodidae) fauna survey in an American cutaneous leishmaniasis (ACL) focus in Baturite, Ceara State, Northeast Barzil. *Parasitol.*, 33: 159-167.

Desjeux, P. (2001).

Worldwide increasing risk factors for leishmaniasis. *Med. Micro. Immunol., (Berlin)*, 190: 77-79.

Devi, D. A.; Thomas, G. and Manu, P. S. (2000).

Effect of lunar cycle on light trap catches of all midge, *Orseolia eryzae* wood masan in Kerala. *J. Appl. Zool. Res.*, 11(2.3): 93-95.

Dinesh, D. S.; Ranjan, A.; Palit, A.; Kishore, K. and Kar, S. K. (2001).

Seasonal and nocturnal landing/ biting behavior of *Phlebotomus argentipes* (Diptera: Psychodidae). *Ann. Trop. Med. Parasitol.*, 95: 197-202.

Doha, S. (1986).

Sand fly ecology in relation to leishmoniasis in El Agamy, Alexandria Governorate. *M.Sc. Thesis, Dept. Entomol., Fac. Sci., Ain Shams Univ.*

Dyer, F. C. (1985).

Nocturnal orientation by the Asian honey bee, *Apis dorsata*. *Anim. Behav.*, 33(3): 769-774.

Eflatoon (1922).

Record of the Museum of Entomological Society of Egypt.

El-Desokay, M. H. M. (1991).

Studies on some biological aspects of common rats in Egypt. *Ph. D. Thesis, Fac. Sci., Benha Branch. Zagazig Univ., Qalyubya, A.R.E.*

El Hossary, S. S.; Shehata, M. G.; Helmy, N.; Lane, R. P. and El Sawaf, B. M. (2000).

Studies on the phlebotomine sand flies (Diptera: Psychodidae) and rodents in cutaneous leishmaniasis focus in

Nekheli, North Sinai, Egypt. *J. Union. Arab. Biol. Zool.*, (13A): 53-64.

El Naggar, M. H.; Shoukry, N. M.; Soliman, B. A.; Darwish, A. B. and El Sawaf, B. M. (2006).

Ecology, biology and susceptibility of *Phlebotomus papatasi* to leishmania experimental infection in Suez Governorate. *J. Egypt. Soc. Parasitol.*, 36(1): 127-138.

Elnaiem, D. A.; Connor, S. J.; Thomson, M. C.; Hassan, M. M.; Hassan, H. K.; Aboud, M. A. and Ashford, R. W. (1998).

Environmental determinants of the distribution of *Phlebotomus orientalis* in Sudan. *Ann. Trop. Med. Parasitol.*, 92(8): 877-887.

El Said, S.; Beier, J. C.; El Sawaf, B. M.; Doha, S. and El Kordy, E. (1986).

Sand flies (Diptera: Psychodidae) associated with visceral leishmaniasis in El Agamy, Alexandria Governorate, Egypt. II- Field behavior. *J. Med. Entomol.*, 23(6): 609-615.

El Said, S. M.; Kenawy, M. A.; El Sawaf, B. M.; Beier, J. C. and El Sawy, F. M. (1985).

Seasonal abundance and distribution of *Phlebotomus papatasi* (Diptera: Psychodidae) inside houses in Aswan Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 15(2): 371-380.

El Sawaf, B. M.; Beier, J. C.; Hussein, S. M.; Kassem, H. A. and Sattar, S. A. (1984).

Phlebotomus langeroni, a potential vector of Kala-azar in the Arab Republic of Egypt. *Trans. Roy. Soc. Trop. Med. Hyg.*, 78(3): 421.

El Sawaf, B. M.; Shoukry, A.; El Said, S.; Lane, R. P.; Kenawy, M. A.; Beier, J. C. and Abdel Sattar, S. (1987).

Sand fly species composition along an altitudinal transect in Southern Sinai, Egypt. *Ann. Parasitol. Hum. Comp.*, 62(58): 467-473.

El Sayed, S. A.; El-Raaba, F. M. and Abd El-Nur, O. (1991).

Daily and seasonal activities of some sand flies from Surrugia village, Khartoum, Sudan. *Parasitol.*, 33(1): 205-215.

El-Ziady, S. and Osman, M. F. H. (1961).

Investigations into the relation between lunar periodicity and the activity of Diptera in Egypt. *Bull. Soc. Entomol. Egypte*, XLV: 287-295.

Flores-Mendoza, C. and Lourenco-de-Oliveria, R. (1996).

Bionomics of *Anopheles aquasalis* Curry 1932 in Guarai, State of Rio Janeiro Southeastern Brasil. *Mem. Inst. Oswaldo Cruz.*, 91(3): 265-270.

Fouad, L. M. and Rashed, F. F. (1986).

Incidence of some domestic and commensal rats trapped during June 1980 – May 1981 in Alexandria. *Alex. J. Agric. Res.*, 13(1): 431-435.

Fryauff, D. J.; Modi, G. B.; Mansour, N. S.; Kreutzer, R. O.; Soliman, S. and Youssef, F. G. (1993).

Epidemiology of cutaneous leishmaniasis at a focus monitored by the multinational force and observers in the Northeastern Sinai Desert of Egypt. *Am. J. Trop. Med. Hyg.*, 49(5): 598-607.

Gad, A. M. and Saad, A. (1974).

Insects of medical importance in Aswan and Lake Nasser area. *A booklet of Aswan Regional Planning, Lake Nasser Development Center, Aswan, A.R.E.*, 106 pp.

Ghosh, K.; Mukhopadhyay, J.; Desai, M. M.; Senroy, S. and Bhattacharya, A. (1999).

Population ecology of *Phlebotomus argentipes* (Diptera: Psychodidae) in West Bengal, India. *J. Med. Entomol.*, 36(5): 588-594.

Gil Collado, J. (1977).

Phlebotomes et les leishmanioses en Espagne. *Colloques Int. Cent. Natr. Res. Scient.*, 239: 177- 189.

Gonzalez, R.; De-Souza, L.; Devera, R.; Jorquera, A.; Ledezma, E. and Sousa. L. (1999).

Seasonal and nocturnal domiciliary human landing / biting behavior of *Lutzomyia* (*Lutzomyia*) *evansi* and *Lutzomyia* (*Psychodepygus*) *panamensis* (Diptera: Psychodidae) in a periurban area of a city on the Caribbean coast of eastern Venezuela (Barcelona; Anzoategui state). *Trans. Roy. Soc. Trop. Med. Hyg.*, 93(4): 361-364.

Goodwin, S. and Danthanarayana, W. (1984).

Flight activity of *Plutella xylostella* (L.) (Lepidoptera: Yponomeutidae). *J. Aust. Entomol. Soc.*, 23(3): 35-240.

Guimaraes, A. E.; Gentile, C.; Lopes, C. M. and Mello, R. P. (2000).

Ecology of mosquitoes (Diptera: Culicidae) in areas of Serra do Mar State Park, State of Sao Paulo, Brazil. III- daily biting rhythms and lunar cycles influence. *Mem. Inst. Oswaldo Cruz.*, 95(6): 753- 760.

Hamilton, J. G. C. and Ramsoondar, T. M. C. (1994).

Attraction of *Lutzomyia longipalpis* to human skin odours. *Med. Vet. Entomol.*, 8: 375-380.

Hanafi, H. A.; El Sawaf, B. M. and Beavers, G. M. (1999).

The effect of *Leishmania major* on some biological parameters of *Phlebotomus papatasi* (Diptera: Psychodidae)

from endemic and non-endemic areas in Egypt. *J. Egypt. Soc. Parasitol.*, 29 (2): 293-305.

Harrat, Z.; Izerouel, S.; Belkaid, M. and Tabet-Derraz, O. (1993).

Seasonal fluctuations of phlebotominae in Algiers and its suburbs. *Archives de inst. Pasteur d'Algerie*, 59: 245-255.

Hassan, A. N.; Kassem, H. A.; El Hossary, S. I. and Hermas, E. M. (2000).

The use of geographic information system to identify environmental factors associated with sand flies prevalence and cutaneous leishmaniasis transmission risk in Northeastern Sinai, Egypt. *J. Egypt. Ger. Soc. Zool.*, 31(E): 13-27.

Hassan, A. N.; Kassem, H. A. and El Sawaf, B. M. (1999).

Species composition and abundance of sand flies at natural desert and modified agricultural lands at Nekhel, Sinai, Egypt. *J. Egypt. Ger. Soc. Zool.*, 28(6): 37-47.

Hati, A. K.; Sur, S.; De, N.; Dwivedi, H. N.; Bhattacharyya, J.; Mukherjee, H. and Chandra, G. (1991).

Longitudinal study on distribution of *Phlebotomus argentipes* sand flies at different heights in cattleshed. *Indian J. Med. Res.*, 93: 388-390.

Hervy, J. P.; Legros, F. and Ferrara, L. (1986).

Influence of moonlight on the biting activity of *Aedes taylori* (Diptera: Culicidae). *Cahiers Orstom., Entomol. Med. Parasitol*, 24(1): 59-65.

Holck, A. R. and Meek, C. L. (1991).

Nocturnal distribution of Louisiana mosquito adults. *J. Am. Mosq. Control Assoc.*, 7(4): 628-632.

Hussein, M. S. (1974).

Studies on the Phlebotominae sandflies the vector of kala azar in the Sudan. *M. Sc. Thesis, Entomol. Dept., Fac. Sci., Ain Shams Univ.*

Ibrahim, A. A. and Abdoon, A. M. (2005).

Distribution and population dynamics of *Phlebotomus* sand flies (Diptera: Psychodidae) in an endemic area of cutaneous leishmaniasis in Asir Region, Southwestern Saudi Arabia. *J. Entomol.*, 2(1): 102-108.

Janini, R.; Saliba, E. and Kamhawi, S. (1995).

Species composition of sand flies and population dynamics of *Phlebotomus papatasi* (Diptera: Psychodidae) in the Southern Jordan Valley, an endemic focus of cutaneous leishmaniasis. *J. Med. Entomol.*, 32(6): 822-826.

Janousek, T. E. and Olson, J. K. (1994).

Effect of a lunar eclipse on the flight activity of mosquitoes in the upper Gulf Coast of Texas. *J. Am. Mosq. Control Assoc.*, 10(2): 222-224.

Johnson, R. N.; Lawyer, P. G.; Ngumbi, P. M.; Mebrahtu, Y. B.; Mwanyumba, J. P.; Monsonik, N. C.; Makasa, S. J.; Githure, J. I. and Roberts, C. R. (1999).

Phlebotomine sand fly (Diptera: Psychodidae) seasonal distribution and infection rates in a defined focus of *Leishmania tropica*. *Am. J. Trop. Med. Hyg.*, 60(5): 854-858.

Kansouh, A. S.; El-Zemaity, M. S. and Mikhail, M. W. (1990).

Population density of domestic rodents and associated fleas at Dahshour, Giza Governorate. *Ann. Agric. Sci., Fac. Agric., Ain Shams Univ. Cairo, Egypt*.

Kassem, H. A.; Abdel Sattar, S.; Shehata, M. G. and El Sawaf, B. M. (1988).

Abnormalities in the sand fly *Phlebotomus papatasi* (Scopoli) (Diptera: Psychodidae) of Egypt. *J. Egypt. Pub. Hlth. Assoc.*, Lxiii, (3,4): 209-213.

Kassem, H. A.; Hassan, A. N. and Kamal, H. A. (1999).

Spatial distribution of *Phlebotomus papatasi* (Scopoli) and *P. bergeroti* Parrot in Sinai, Egypt in relation to some landscape variables. *J. Egypt. Ger. Soc. Zool.*, 28(E): 189-201.

Kelly, D. W. and Dye, C. (1997).

Pheromones, kairomones and the aggregation dynamic of the sand fly. *Anim. Behav.*, 53: 721-731.

Kenawy, M. A.; El Sawaf, B. M. and Doha, S. (1990).

Sand flies (Diptera: Psychodidae) in Mersa Matruh and Siwa Oasis, A. R. Egypt. *J. Egypt. Pub. Hlth. Assoc.*, lxx (3,4): 283-289.

Khalid, M. L.; Morsy, T. A.; El-Shennawy, S. F.; Farrag, A. M.; Sabry, A. H. and Mostafa, H. A. (1992).

Studies on flea fauna in El-Fayoum Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 22: 783-799.

Khalil, M. B. (1934).

Dermal leishmaniasis: A study of an endemic focus in Egypt. *Archiv fur. Schiff. Tropenhygiene*, 38: 417-433.

Kravchenko, V.; Wasserberg, G. and Warburg, A. (2004).

Bionomics of phlebotomine sand flies in the Galilee focus of cutaneous leishmaniasis in northern Israel. *Med. Vet. Entomol.*, 18(4): 418-428.

Lane, R. P. (1986).

The sand flies of Egypt (Diptera: Phlebotominae). *Bull. Br. Mus. Nat. Hist. (Ent.)*, 52(1):1-36.

Lewis, D. J. and Buttiker, W. (1982).

Insects of Saudi Arabia the taxonomy and distribution of Saudi Arabia phlebotomine sand flies (Diptera: Psychodidae). *Fauna of Saudi Arabia* 4.

Lillie, T. H.; Kline, D. L. and Hall, D. W. (1987).

Diel and seasonal activity of *Culicoides* spp. (Diptera: Ceratopogonidae) near Yankeetown. Florida, monitored with a vehicle-mounted insect trap. *J. Med. Entomol.*, 24(4): 503-511.

Linhares, A. X. and Anderson, J. R. (1990).

The influence of temperature and moonlight on flight activity of *Culicoides variipennis* (Coquillett) (Diptera: Ceratopogonidae) in northern California. *Pan Pacific Entomologist*, 66(3):199-207.

Lourenco-de-Oliveira, R.; Silva, T. F.; Heyden, R. De-Oliveira, R. L.; Oliveira, R. L.; Fernandes, S. T. and Da-Silva, T. F. (1985).

Some aspects of the ecology of the mosquitoes (Diptera: Culicidae) of a lowland area (Granjas Calabria), in Jacarepagua, Rio de Janeiro. II. Monthly frequency and in relation to the lunar cycle. *Mem. Inst. Oswaldo Cruz.*, 80(2): 123-133.

Mahadevan, N. R.; Chelliah, S.; Regupathy, A. and Jayaraj, S. (1985).

Population dynamics of the sorghum stem borer, *Chilo partellus* (Swinhoe) in light trap. *Behavioral and physiological approaches in pest management*, 104: 106.

Mansour, N.S.; Awadalla, H.N.; Youssef, F.G. and Tewfik, S. (1984).

Characterization of leishmania isolates from children with visceral infection contracted in Alexandria, Egypt, *Trans. Roy. Soc. Trop. Med. Hyg.*, 78: 704.

McHugh, C. P.; Ostrander, B. F.; Raymond, R. W. and Kerr, S. F. (2001).

Population dynamics of sand flies (Diptera: Psychodidae) at two foci of leishmaniasis in Texas. *J. Med. Entomol.*, 38(2): 268-277.

Mellor, H. E. and Hamitton, J. G. C. (2003).

Navigation of *Lutzomyia longipalpis* (Diptera: Psychodidae) under or starlight conditions. *Bull. Entomol. Res.*, 93: 315-322.

Merdan, A. I.; Morsy, T. A.; Shoukry, A. and Wahba, M. M. (1992).

Two successive years study on *Phlebotomus papatasi* in North Sinai Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 22(1): 91-100.

Mohanraj, D.; Zanarthanan, R. and Suresh, S. (1989).

Effect of lunar phase an attraction of rice pests to black light trap. *International Rice Research, Newsletter*, 14(4): 36.

Morillas-Marquez, F.; Guevarra-Benitez, D.; Ubeida-Ontiveros, J. and Gonzalea-Castro, J. (1983).

Fluctuacions annuelles des populations de phlebotomes (Diptera: Phlebotomidae) dans la province de Granade (Espagne). *Ann. Parasitol. Hum. Comp.*, 58, 625-632.

Morrison, A. C.; Ferro, C.; Pardo, R.; Torres, M.; Devlin, B.; Wilson, M. L. and Tesh, R. B. (1995a).

Seasonal abundance of *Lutzomyia longipalpis* (Diptera: Psychodidae) at an endemic focus of visceral leishmaniasis in Colombia. *J. Med. Entomol.*, 32(4): 538-548.

Morrison, A. C.; Ferro, C.; Pardo, R.; Torres, M.; Willson, M. L. and Tesh, R. B. (1995b).

Nocturnal activity patterns of *Lutzomyia longipalpis* (Diptera: Psychodidae) at an endemic focus of visceral leishmaniasis in Colombia. *J. Med. Entomol.*, 32(5): 605-617.

Morsy, T. A. (1983).

Cutaneous leishmaniasis in Egypt. Historical background and current knowledge. *J. Egypt. Soc. Parasitol.* 13: 597-611.

Morsy, T. A.; Aboul-Ela, R. G.; Sarwat, M. A. A.; Arafa, M. A. S. and El-Gozamy, B. M. R. (1993).

Some aspects of *Phlebotomus papatasi* (Scopoli) in greater Cairo, Egypt. *J. Egypt. Soc. Parasitol.*, 23(2): 399-416.

Morsy, T. A; El Ela, R. G. A. and El Gozamy, B. M. R. (1988).

The commensal rodents and their flea fauna in Alexandria city, Egypt. *J. Egypt. Soc. Parasitol.*, 18: 11-28.

Morsy, T. A.; El-Ela, R. G. A.; Rifaat, M. M. A. and Dakhil, M. A. A. (1995).

The seasonal and daily activities of *Phlebotomus papatasi* in Riyadh, Saudi Arabia. *J. Egypt. Soc. Parasitol.*, 25(3): 699-711.

Morsy, T. A.; El-Missiry, A. G.; Kamal, A. M.; Fayad, M. E. and El-Sharkawy, I. M. (1990).

Distribution of *Phlebotomus* species in the Nile Delta, Egypt. *J. Egypt. Soc. Parasitol.*, 20(2): 589-597.

Morsy, T. A; Hamadto, H. A.; Rashed, S. M.; El Fakahany, A. F. and Abdalla, K. F. (1990).

Animals as reservoir hosts for leishmania in Qalyobia Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 20: 779- 788.

Morsy, T. A; Micheal, S. A.; Bassili, W. R. and Saleh, M. S. M. (1982).

Studies on rodents and their zoonotic parasites, particularly leishmania in Ismailia. *J. Egypt. Soc. Parasitol.*, 12: 568-585.

Morsy, T. A.; Naser, A. M.; El-Gibali, M. R.; Anwar, A. M. and El-Said, A. M. (1995).

Studies on zoonotic cutaneous leishmaniasis among a group of temporary workers in North Sinai Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 25: 99-106.

Mourad, M. G.; Abd El-Gawad, K. H. and Maher, A.A. (1982).

Population density of rodent species in some urban areas in Minia Governorate. *Assiut J. Agric. Sci.*, 13(2): 19 – 26.

Mutinga, M. J.; Basimike, M.; Kamau, C. C. and Mutero, C. M. (1990).

Epidemiology of leishmaniasis in Kenya. Natural host preference of wild caught phlebotomine sandflies in Baringo District, Kenya. *East. Afr. Med. J.*, 67(5): 19-27.

Nag, A. and Nath, P. (1991).

Effect of moon light and lunar periodicity on the light trap catch of cutworm *Agrotis ipsilon* (Hufn.) moths. *J. Appl. Entomol.*, III: 555-560.

Nemec, S. J. (1971).

Effect of lunar phases on light-trap collections and population of Bollworm Moths. *J. Econ. Entomol.*, 64: 860-864.

Neumann, D. (1995).

Physiological clocks of insects-ecophysiology of reproductive activities controlled by lunar cycles. *Naturwissenschaften*, 82: 310-320.

Nowinszky, L. (2004).

Nocturnal illumination and night flying insects. *Appl. Eco. Envi. Res.*, 2(1): 17-52.

Okka, M. A. (1992).

Studies on rodents and their parasites in Kafr El-Sheikh, A.R.E. *Ph. D. Agric. (Ent.)*, *Fac. Agric., Kafr El- Sheikh, Tanta Univ.*

Osborn, D. J. and Helmy, I. (1980).

The contemporary land mammals of Egypt (including Sinai). *Fieldiana Zoology, Field Museum of Natural History, new series*, No. 5.

Ostfeld, R. S.; Roy, P.; Haumaier, W.; Conter, L.; Keesing, F. and Rowton, E. D. (2004).

Sand fly (*Lutzomyia vexator*) (Diptera: Psychodidae) populations in upstate New York: abundances microhabitat, and phenology. *J. Med. Entomol.*, 41(4): 774-778.

Pandya, H. V.; Shah, A. H. and Purohit, M. S. (1989).

Influence of lunar cycle on light trap catches of rice stem borer, *Scirpophaga incertulas*. *Oryza*, 26(1-2): 108-109.

Papadopoulos, B. and Tselentis, Y. (1994).

Sand flies in the Greater Athens Region, Greece. *Parasite*, 1(2): 131-140.

Paras, N.; Rakesh, R.; Nath, P. and Rai, R. (2000).

Effect of moon light and lunar periodicity on the light trap catches of gram pod borer, *Helicoverpa armigera* (Hubner). *Shashpa*, 7(1): 25-27.

Pedgley, D. E. (1986).

Windborne migration in the Middle East by the moth *Heliothis armigera* (Lepidoptera: Noctuidae). *J. Ecol. Entomol.*, 11: 467-470.

Quinnell, R. J.; Dye, C. and Shaw, J. J. (1992).

Host preference of the sand fly *Lutzomyia longipalpis* in Amazonian Brazil. *Med. Vet. Entomol.*, 6: 195-200.

Rebelo, J. M. M. (2001).

Hourly frequency and seasonality of *Lutzomyia longipalpis* (Diptera: Psychodidae, Phlebotominae) on Sao Luis Island, Maranhao, Brazil. *Cad. Saude Publica.*, 17(1): 221-227.

Rebelo, J. M. M.; De Oliveira, S.T.; Barros, V. L. L.; Silva, F. S. and De-Oliveira, S. T. (2001a).

Seasonal fluctuation of phlebotomine sand flies (Diptera: Psychodidae) in a recent colonization area in the Municipal District of Buriticupu, Amazonian Maranhao, Brazil. *Rev. Brasil Entomol.*, 45(1): 11-16.

Rebelo, J. M. M.; De Oliveira, S. T.; Silva, F. S.; Barros, V. L. L. and Costal, J. M. (2001b).

Sand flies (Diptera: Psychodidae) of the Amazonia of Maranhao. V. Seasonal occurrence in ancient colonization area and endemic for cutaneous leishmaniasis. *Brazil J. Biol.*, 61(1): 107-115.

Rebollar-Tellez, E. A.; Andrade-Narvaez, F. J.; Fernandez-Salas, I. and Reyes Villanueva, F. (1996a).

Collections of sand flies (Diptera: Psychodidae) from mammal burrows in an area of cutaneous leishmaniasis in Campeche, Mexico. *Entomological News*, 107(5): 317-321.

Rebollar-Tellez, E. A.; Reyes-Villanueva, F.; Fernandez-Salas, I. and Andrade Narvaez, F. J. (1996b).

Population dynamics and biting rhythm of the anthropophilic sand fly *Lutzomyia cruciata* (Diptera: Psychodidae) in Southeast Mexico. *Rev. Inst. Med. Trop.*, 38(1): 29-33.

Rifaat, M. A.; Morsy, T. A. and Abdel-Mawla, M. M. (1981).

Seasonal activity of *Rattus norvegicus* and flea index on Port Said Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 11: 525-532.

Rifaat, M. A.; Morsy, T. A. and Hassan, Z. A. (1968).

Visceral leishmaniasis in UAR.

Roberts, D. M. (1994).

Arabian sand flies (Diptera: Psychodidae) prefer the hottest nights. *Med. Vet. Entomol.*, 8(2): 194-198.

Rubio-Palis, Y. (1992).

Influence of moon light on light trap catches of the malaria vector *Anopheles nuneztovari* in Venezuela. *J. Am. Mosq. Control Assoc.*, 8(2): 178-180.

Salomon, O. O.; Rossi, G. C.; Cousino, B.; Spinelli, G. R.; de Arias, A. R.; Del Puerta, D. A. L. and Artiz, A. J. (2003).

Phlebotominae sand flies in Paraguay, abundance distribution in the Southeastern Region. *Mem. Inst. Oswaldo cruz.*, 98(2): 185-190.

Santos-De-Marco, T.; Gaia, M.C.D. and Brazil, R. P. (2002).

Influence of the lunar cycle on the activity of phlebotomine sand flies (Diptera: Psychodidae). *J. Am. Mosq. Control Assoc.*, 18(2): 114- 118.

Shehata, M.G.; Doha, S.; El Hossary, S.; Swalem, M.; Abd El Mohsen, A. and El Kadi, G. (1995).

Field behavior of sandflies in Wadi Feiran, Sinai- Egypt. *J. Egypt. Ger. Soc. Zool.*, 16(E): 29- 52.

Sheng, C. F.; Wang C. Z.; Zhong J.; Gac Z. X.; Wang, H. T. and Xuan, W. J. (2003).

Relationship between lunar phase and capture rate of cotton bollworm moths using poplar twig bundle traps in Tongzhou, Jiangsu Province. *Entomological Knowledge*, 40(6): 512-514.

Sherlock, I. A.; Maia, H. and Dias-Lima, A. G. (1996).

Preliminary results of a project on the ecology of phlebotomine vectors of cutaneous leishmaniasis in the state of Bahia. *Rev. Soc. Brasil. Med. Trop.*, 29(2): 207-214.

Shoukry, A.; Morsy, T. A.; Abu Hashish, T. A. and El Kady, G. A. (1986).

Seasonal activities of two commensal rats and flea index in north Sinai Governorate, Egypt. *J. Egypt. Soc. Parasitol.*, 16: 385- 393.

Shrivastava, S. K.; Shukla, B. C. and Shastri, A. S. (1987).

Effect of lunar cycle on light trap catches of *Spodoptera litura*. *Indian J. Agri. Sci.*, 57(2): 117-119.

Singh, K. V. (1999).

Studies on the role of climatological factors in the distribution of phlebotomine sand flies (Diptera: Psychodidae) in semi-arid areas of Rajasthan, India. *J. Arid. Enviro.*, 42: 43-48.

Soliman, B. A.; Hassan, M. M. and El Hossary, S. S. (2001a).

Species composition of phlebotomine sand fly (Diptera: Psychodidae) in the Southeastern transect of Upper Egypt. *Proceedings of "Elflatoun's Conference of Entomol. 1st 199-210.*

Soliman, S.; Marzouk, A. S.; Main, A. J. and Montasser, A. A., (2001b).

Seasonal studies on commensal rats and their ectoparasites in a rural area of Egypt: the relationship of ectoparasites to the

species, locality and relative abundance of the host. *J. Parasitol.*, 87: 545-553.

Souza, N. A.; Andrade-Coelho, C. A.; Silva, V. C.; Peixoto, A. A. and Rangel, E. F. (2005).

Moonlight and blood-feeding behavior of *Lutzomyia intermedia* and *Lutzomyia whitmani* ((Diptera: Psychodidae: Phlebotominae), vectors of American cutaneous leishmaniasis in Brazil. *Mem. Inst. Oswaldo Cruz.*, 100(1): 39-42.

Souza, N. A.; Andrade-Coelho, C. A.; Vilela, M. L.; Peixoto, A. A. and Rangel, E. F. (2002).

Seasonality of *Lutzomyia intermedia* and *Lutzomyia whitmani* (Diptera: Psychodidae: Phlebotominae), occurring sympatrically in area of cutaneous leishmaniasis in the state of Rio de Janeiro, Brazil. *Mem. Inst. Oswaldo Cruz.*, 97(6): 759-765.

Spiegel, M. R. (1961).

Schaum's outline of theory and problems of statistics. *Schaum Publ. Co., New York*, 359 pp.

Srinivasa, N.; Viraktamath, C. A. and Thontadarya, T. S. (1990).

Influence of moon phase and weather on the light trap catches of insect pests of rice. *Oryza*, 27(2): 183-190.

Srinivasan, R.; Panicker, K. N. and Dhando, V. (1993).

Population dynamics of *Phlebotomus papatasi* (Diptera: phlebotominae) in Pondicherry, India. *Acta Trop.*, 54(2): 125-130.

Svobodova, M.; Sadlova, J.; Chang, K. P. and Volt, P. (2003).

Distribution and feeding preference of the sand flies *Phlebotomus sergenti* and *P. papatasi* in a cutaneous leishmaniasis focus in Sanliurfa, Turkey. *Am. J. Trop. Med. Hyg.*, 68 (1): 6-9.

Teodoro, U.; Filho, V. L.; Lima, E. M.; Spinoso, R. P.; Barbosa, O. C.; Ferreira, M. E. M. C. and Lonardnni, M. V. C. (1993).

Observacoes sobre o comportamento de flebotomineos em ecotopos florestais, em area endemica de Leishmaniose Tegumentar Americana, do norte do Estado do Parana, Sul do Brasil. *Rev. Saude Publ.*, 27: 242-249.

Tesh, R.B. (1988).

The genus phlebovirus and its vectors. *Ann. Rev. Entomol.*, 33: 169-181.

Tewfik, S.; Kassem, S.A.; Aref, M.K. and Awadalla, H.N. (1983).

A preliminary report on two cases of visceral leishmaniasis in Egypt. *Trans. Roy. Soc. Trop. Med. Hyg.*, 77: 334-335.

Theodor, O. (1948).

On some sand flies (*Phlebotomus*) of the *sergenti* group in Palestine. *Bull. Entomol. Res.*, 39: 85-111.

Theodore, P. S. (1983).

The dynamic universe, an introduction to astronomy, pp. 13-15.

Toprak, S. and Ozer, N. (2005).

Sand fly species of Sanliurfa province Turkey. Short communication. *Med. Vet. Entomol.*, 19(1): 107-110.

Toth, G. and Nowinszky, L. (1988).

A connection between the illumination of environment the twilight polarization phenomena and the catches of turnip moth *Scotia segetum* Schiff by light-trap. *Acta phytopathologica et entomologica hungarica*, 23(1-2): 167-186.

Travi, B. L.; Montoya, J.; Gallego, J.; Jaramillo, C., Llano, R. and Velez, I. O. (1996).

Bionomics of *Lutzomyia evansi* (Diptera: Psychodidae) vector of visceral leishmaniasis in Northern Colombia. *J. Med. Entomol.*, 33(3): 278-285.

Trouillet, J.; Ba, Y.; Traore-Lamizana, M.; Zeller, H. G. and Fontenille, D. (1995).

Phlebotomine sand flies (Diptera: Psychodidae) from Senegal. Communities from the Ferlo area. Arboviruses isolation. *Parasite*, 2(3): 289-296.

Wahba, M. M.; Schnur, L. F.; Morsy, T. A. and Merdan, A. (1990).

The characterization of *leishmania major* from *Phlebotomus papatasi* (Scopoli) caught in northern Sinai, Egypt. *Trans. Roy. Soc. Trop. Med. Hyg.*, 84(6): 785-786.

Wasserberg, G.; Yaram, I. and Warburg, A. (2003).

Seasonal abundance patterns of the sand fly *Phlebotomus papatasi* in climatically distinct foci of cutaneous leishmaniasis in Israeli deserts. *Med. Vet. Entomol.*, 17(4): 452-456.

Willcocks, F. C. (1917).

Notes on some insects found in Egypt of medical and veterinary interest. *Bull. Soc. Entomol., d' Egypte*, 10: 79-90.

Williams, P. (1970).

Phlebotomine sand flies and leishmaniasis in British Honduras (Belize). *Trans. Roy. Soc. Trop. Med. Hyg.*, 64(2): 317-368.

World Health Organization (WHO), (1990).

Control of the leishmaniasis, World Health Organization Technical Report Series, 793.

Ximenes, M. F.; Souza, M. F. and Castellon, E. G. (1999).

Density of sand flies (Diptera: Psychodidae) in domestic and wild animal shelters in an area of visceral leishmaniasis in the state of Rio Grande do Norte Brazil. *Mem. Inst. Oswaldo Cruz.*, 94(4): 427-432.

Yaghoobi-Ershadi, M. R. and Javadian, E. (1997).

Studies on sand flies in a hyperendemic area of zoonotic cutaneous leishmaniasis in Iran. *Indian J. Med. Res.*, 105: 61-66.

Yela, J. L. and Holyoak, M. (1997).

Effects of moon light and meteorological factors on light and bait trap catches of noctuid moths (Lepidoptera: Noctuidae). *Envir. Entomol.*, 26(6): 1283-1290.

Young, D. G. and Duncan, M. A. (1994).

Guide to the identification and geographic distribution of *Lutzomyia* sand flies in Mexico, the West Indies, Central and South America (Diptera; Psychodidae). *Mem. Am. Entomol. Inst.*, 54: 1-881.

Yuval, B. (1991).

Population of *Phlebotomus papatasi* (Diptera: Psychodidae) and the risk of *Leishmania major* transmission in three Jordan Valley habitats. *J. Med. Entomol.*, 28(4): 492-495.

Yuval, B. and Schlein, Y. (1991).

Leishmaniasis in the Jordan Valley. III. Nocturnal activity of *Phlebotomus papatasi* (Diptera: Psychodidae) in relation to nutrition and ovarian development. *J. Med. Entomol.*, 23: 411- 415.

Zanuncio, T. V.; Zanuncio, J. C.; Aroujo; M. S. and Evaristo, F. C. (1995).

Effect of the phases of the moon on Lepidoptera, in Eucalyptus plantations, in Acailandia, Maranhao, Brazil. *Revista Arvore*. 19(1): 100-109.

Zeese, W.; Khalaf, S. A.; Morsy, T. A. and Abou-El Ela, R.G. (1990).

Rodents and their ectoparasites in Sharkia Governorate, Egypt. *J. Egypt Sec. Parasitol.*, 20: 827-835.

Zein El- Dine, K. (1972).

Phlebotominae (Diptera: Psychodidae) of Egypt. *J. Egypt. Publ. Hlth. Assoc.*, 47(5): 269-272.