

REFERENCES

- Abd El-Rehim, M.A.; Fawaz, K. and Gobrial, W.N. (1976).** Differentiation between *Rhizobium* species and *Agrobacterium* by serological means. Alex. J. Agric. Res. 24 (1):133-140.
- Abd El-Rehim, M.A.; Hassouna, M.S. and El-Badry, H.M. (1995).** Serological relationships between virulent and avirulent mutants of *Pseudomonas solanacearum*. J. Agric. Sci., Mansoura Univ. 20 (12): 5073-5090.
- Abd El-Zaher, H.M.A. (2004).** Occurrence of damping-off of wheat caused by *Pythium diclinum tokunaga* in El-Minia, Egypt and its possible control by *Gliocladium roseum* and *Trichoderma harzianum*. Phytopathology and Plant Protection, 37(2): 147-159.
- Abou-Elseoud, M.S. and Saeed, F.A. (1990).** Relation of virulence to the intracellular soluble protein of *Cephalosporium maydis* (the incidence of late wilt disease of corn). Assiut Journal of Agric. Science, 21 : 165-178.
- Aghighi, S.; Shahidi-Bonjar, G. H.; Rawashdeh, R.; Batayneh, S. and Saadoun, I. (2004).** First report of antifungal spectra of activity of Iranian Actinomycetes strains against *Alternaria solani*, *Alternaria alternata*, *Fusarium solani*, *Phytophthora megasperma*, *Verticillium dahliae* and *Saccharomyces cerevisiae*. Asian Journal of Plant Sciences, 3(4): 463-471.
- Ala El-Dein, O. and El-Kady, S. (1985).** Crossed-immunoelectrophoretic studies on *Botrytis cinerea* Pers and other *Botrytis* species. Acta. Phytopathologica Academiae Scientiarum Hungaricae, 20: 291-301.
- Allam, A.D. (1994).** Reaction of some wheat cultivars to root rot disease caused by *Cochliobolus sativus*. Assiut Journal of Agricultural Sciences, 25 (1):249-264.
- Allam, A.D.; El-Zawahry, A.M.; Mohamed, M.S. and Bakr, D.W. (2000 b).** Reaction of certain wheat cultivars in different soil textures to root-rot disease caused by *Fusarium moniliforme*, *Fusarium graminearum* and /or *Drechslera sorokiniana*. Assiut Journal of Agricultural Sciences, 31(3):213-226.
-

REFERENCES

- Allam, A.D.; El-Zawahry, A.M.; Mohamed, M.S. and Bakr, D.W. (2000 a).** Fungi associated with wheat foot and root-rot diseases in Assiut governorate and some factors affecting disease incidence. Assiut Journal of Agricultural Sciences, 31(4): 41-57.
- Amer, G.A. and El-Shennawy, R.Z.(2003).** Field application of powder formulation of *Bacillus subtilis* for controlling damping-off and root rot of Soybean. Minufiya J.Agric. Res., 28 (4):1079-1091.
- Amer, G.A. and Utkhede, R.S. (2000).** Development of formulations of biological agents for management of root rot of lettuce and cucumber. Can. J. Microbiol., 46: 809-816.
- Andres-Ares, J.L.; Alonso-Ferro, R.C.; Campo-Ramirez, L. and Moreno-Gonzalez, J.(2004).** *Fusarium graminearum* schwabe, a maize root and stalk rot pathogen isolated from lodged plants in Northwestern Spain. Spanish Journal of Agricultural Research, 2 (2): 249-252.
- Axelsen, N.H.;Krou, J. and Weeke, B.(1973).** A manual of quantitative immunoelectrophoresis. Methods and applications. Oslo, Bergen and Tromso Univ.
- Babadoost, M.(1995).** *Fusarium* species in wheat seeds and plants in East Azerbaijan and Ardabil Provinces. Iranian Journal of Plant Pathology,31(1/4): 33-36.
- Balmas, V.(1994).** Root-rot of wheat in Italy caused by *Fusarium graminearum* Group 1. Plant Disease,78 (3): 317.
- Balmas, V.; Burgess, L.W. and Summerell, B.A. (1995).** Reaction of durum wheat cv.Yallaro to crown and root-rot caused by *Fusarium graminearum* group 1 and *Fusarium crookwellense*. Australasian Plant Pathology, 24(4):233-237.
- Banu, S.P. and Deverall, B.J.(1996).** Pathogenicity assays with *Bipolaris sorokiniana* for common root-rot of wheat. Bangladesh Journal of Plant Pathology,12 (1-2): 55-57.
- Barnett, H.J.(1960).** Illustrated genera of imperfect fungi. Burgeess, Minneaplis, USA,226 pp.
-

REFERENCES

- Basilio, A.;Gonzalez, I.;Vicente, M.F.;Gorrochategui, J.;Cabello, A.; Gonzalez, A. and Genilloud, O.(2003).**Patterns of antimicrobial activities from soil Actinomycetes isolated under different conditions of pH and salinity. Journal of Applied Microbiology, 95 (4) : 814-823.
- Bertagnolli, B.L.;Daly, S. and Sinclair, J.B. (1998).** Antimycotic compounds from the plant pathogen *Rhizoctonia solani* and its antagonist *Trichoderma harzianum*.ISSN 0931-1785,Phytopath. 146, 131-135.
- Betina,V. (1989).** Mycotoxins: chemical, biological and environmental aspects.Bioactive Molecules, Vol. 9. Elsevier, Amsterdam.
- Bhargava, N.;Shukla, D.N.;Pandey, R.S. and Khati, D.V.S.(1982).** Foot-rot of wheat: a new disease.Indian-Phytopathology, 35(3): 543-544.
- Booth, C.(1977).** *Fusarium* laboratory guide to the identification of the major species. Common Wealth Mycological Institute, Kew Surey, England, 130-153.
- Centurion , M.A.P.C and Kimati, H.(1992).**Serological characterization of *Fusarium moniliforme* and *Fusarium moniliforme* var *subglutinans summa* , Phytopathologica ,18 (3-4):239-246 .
- Chakrabarty, S. and Chandra, A.L.(1979).**Antifungal activity of A-7, a new tetraene antibiotic.Indian Journal of Experimental Biology, 17 : 313-317.
- Chen, C.; Collins, D.J. and Morgan-Jones, G.(1996).** Fungi associated with root rot of winter wheat in Alabama. Journal of Phytopathology, 144(4):193-196.
- Ciccarese, F.; Frisullo, S.; Amenduni, M. and Cirulli, M. (1992).** Biological control of *Sclerotium rolfsii* root-rot of sugar beet with *Trichoderma harzianum*. In "Biological control of plant diseases:Pro-gress and Challenge for the Future"(Tjamos, E.C.; Papavizas, G.C. and Cook, R.J. Eds.), pp.243-248, Plenum Press, New York.
- Comporota, P.(1985).** Antagonism *in vitro* of *Trichoderma* spp. to *Rhizoctonia solani* Kuhn Agronomic,5:613-620 (C.F.Rev. Plant Pathology, (1985) 65 :581).
-

REFERENCES

- Cook, R.J.(1980).***Fusarium* foot rot of wheat and its control in the pacific Northwest. Plant Dis, 64, 1061-1066.
- Cook, R.J. and Veseth, R.J.(1991).**Wheat Health Management.APS Press, St. Paul, Minnesota.
- Dahiya, J.S.; Woods, D.L. and Tewari, J.P. (1988).** Control of *Rhizoctonia solani*, causal agent of brown girdling root-rot of rape seed, by *Pseudomonas fluorescens* . Botanical Bulletin of Academia Sinica, Taiwan, 29(3): 135-141.
- Dennis, C. and Webster, J. (1971).**Antagonistic properties of species groups of *Trichoderma*.I-Production of non-volatile antibiotics. Trans.Bri. Mycol. Soc. 57, 25-39.
- Dhingra, O.D. and Sinclair, J.B.(1995).**Basic plant pathology methods. Second Edition, CRC Press, Inc.pp.434.
- Ding, L.X.;Wang, X.;Wang, D.C. and Fang, S.K.(1999).**Studies on the agricultural antibiotic controlling tobacco brown spot. Journal of Jilin, Agricultural University, 21 (2) : 16-19.
- Doubly, J.A.; Flor, H.H. and Clagett, C.O. (1960).** Relation of antigens of *Melampsora lini* and *Linum usitatissimum* to resistance and susceptibility. Science, 131: 229.
- Duczek,L.J. (1994).** Relationship between a greenhouse and field assay for biological control of common root rot of spring wheat and barley. Canadian Plant Disease Survey, 74(2): 135-140.
- Duthie, J.A.; Hall, R. and Asselin, A.V. (1986).** *Fusarium* species from seed of winter wheat in Eastern Canada. Canadian Journal of Plant Pathology, 8 (3): 282- 288.
- Egorov, N.S.(1985).** Antibiotics a Scientific Approach. MIR Publishers Moscow, PP.439.
- Ehteshamul Haque, S.; Hashmi, R.Y. and Ghaffar, A. (1992).** Biological control of root-rot disease of lentil.Lens,19(2):43-45.
- Elad, Y., Chet, I. and Katan, J. (1980).** *Trichoderma harzianum*: Abio-control agent effective against *Sclerotium rolfsii* and *Rhizoctonia solani*. Phytopathology 70,119-121.
-

REFERENCES

- El-Eraky, A.M.;Saeed, F.A.;Mohamed, M.S. and Amein, A.M.(1993).** Seed and root exudates of wheat cultivars in relation to the association of pathogenic fungi. Assiut Journal of Agricultural Sciences,24(4): 185-205.
- El-Gammal, N.(2004).** Studies on *Fusarium* dry rot of potato. M. Sc., Faculty of Agriculture, Minufiya, University, pp. 84
- El-Kady, S. (1991).**Serological comparison of three races of *Fusarium oxysporum* f.sp. *Lycopersici*. J. Agric.Res.Tanta Univ., 17(2).
- El-Kady, S.;Ala El-Dein, O.; Hoy, J.W. and Schneider, R.W.(1993).** Serological relation-ships among nine *Pythium* species determined by crossed-immunoelectrophoresis.J.Agric.Res.Tanta univ.,19 (1).
- El-Kazzaz, M.K.;Hassan, M.A.;Ala El-Dein, O. and El-Shamy, M.M. (1997).**Serological comparison between two wheat stem rust races.Proceedings of the 9Th Conference of Microbiology Cairo, Egypt, March 25-27, 1997.
- El-Meleigi, M.A.(1988).** Fungal diseases of spring wheat in central Saudi Arabia. Crop- Protection, 4(3): 207-209.
- El-Meleigi, M.A.(1989).** Effect of soil *Pseudomonas* isolates applied to corn, sorghum and wheat seeds on seedling growth and corn yield. Canadian Journal of Plant Science,69 (1):101-108.
- El-Meleigi, M.A.;El-Rokaibah, A.A.;Ibrahim, G.H.;El-Shair, S.M. and Al-Omir, M.E.(1994).** Disease and pest outbreaks.Saudi Arabia. Survey of wheat diseases in central Saudi Arabia.Arab and Near East Plant Protection Newsletter. No.18, 30; 5.
- El-Nashar, F.K.;El-Mokadem, M.T. and Ammar, H.A.M.(2000).** Pathogens associated with root-rot disease of wheat in Egyptian dryland.Egyptian Journal of Agricultural Research, 78 (2) : 559-573.
- El-Nashar, F.K.;El-Mokadem, M.T.;Abd El-Moity, T.H.and Ammar, H.A.M.(2001).** Biological control of root-rot disease of wheat. Egyptian Journal of Agricultural Research, 79 (1): 1-19.
-

REFERENCES

- El-Shamy, M.M. and Amer, G.A.(2000).** Biological control for wheat loose smut using metabolites of certain fungal antagonists. Minufiya J. Agric. Res. Vol.25 (2):317-325.
- Evidente, A.; Conti, L.; Altomare, C.; Bottalico, A.; Sindona, G.; Segre, A.L. and Logrieco, A. (1994).** Fusapyrone and deoxyfusapyrone, two antifungal alpha-pyrones from *Fusarium semitectum*. Natural Toxins, 2 (1): 4-13.
- Fernandez, J.A.; Wofford, D.S. and Horton, J.L.(1985).** Augmentation of wheat common root rot by *Fusarium acuminatum*. Mycopathologia, 90 (3):177-179.
- Flor, H.(1971).** Current status of the gene-for-gene concept. Annu. Rev. Phytopath. 9: 275-296.
- Gesheva, V.; Ivanova, V. and Gesheva, R. (1994).** Biological characteristics and antibiotics production of *Streptomyces hygroscopicus* strains. Actinomycetes 5: 57-63.
- Ghisalberti, E.L.; Narbey, M.J.; Dewan, M.M. and Sivasithamparam, K.(1990).** Variability among strains of *Trichoderma harzianum* in their ability to reduce take-all and to produce pyrones. Plant and Soil, 121,287-291.
- Gilman, J.C.(1957).** A manual of soil fungi. Iowa State University Press, Ames. Iowa. USA, 450 pp.
- Godtfredsen, W.O. and Vangedal, S. (1965).** Trichodermin, a new sesquiterpene antibiotic. Acta Chemica Scandinavica 19, 1088-1102.
- Gupta, V.K.; Sunita Sen and Bhardwaj, L.N. (1992).** Management of *Fusarial* wilt of black locust. Nitrogen Fixing Tree Research Reports, 10: 191-192.
- Guranowska, M.R.; Wiatroszak, I. and Hornok, L.(1984).** Serological comparison of two races of *Fusarium oxysporum* f. sp. *Lupini*. Phytopathologische Zeitschrift, 110(3): 221-225.
- Guranowska, M.R. and Wolko, B.(1991)** Comparison of *Fusarium oxysporum* and *Fusarium oxysporum* var. *redolens* by analyzing the isozyme and serological patterns. Phytopathology, 132 : 287-293
-

REFERENCES

- Hammad, A.M. and El-Mohandes, M. (1999).** Controlling *Fusarium* wilt disease of cucumber plants via antagonistic microorganisms in free and immobilized states. *Microbiological Research*.154 (2): 113-117.
- Handelsman, J.; Raffel, S.; Mester, E.H.;Wunderlich, L. and Grau, C.R.(1990).** Biological control of damping-off alfalfa seedlings with *Bacillus cereus* UW85. *Appl. Environ. Microbiol.*,56: 713-718.
- Harding, H.(1973).**Fungi associated with sub-crown internodes of wheat (*Triticum aestivum*). *Can.J.Bot.* 51,2514-5216.
- Higginbotham, R.W.;Paulitz, T.C. and Kidwell, K.K.(2004).**Virulence of *Pythium* species isolated from wheat fields in Eastern Washington. *Plant Disease*, 88 (9): 1021–1026 .
- Hill, J.P.; Fernandez, J.A. and McShane, M.S. (1983).**Fungi associated with common root rot of winter wheat in Colorado and Wyoming *Plant Disease*, 67, 795-797.
- Hornok, L.(1980).** Serotaxonomy of *Fusarium* species of the sections Gibbosum and Discolor. *Trans. Briti. Mycol. Soc.* 74 (1): 73-78.
- Hussein, E.M.(1992).** Biochemical and serological studies for determining susceptibility of cotton cultivars to *Fusarium oxysporum* f.sp. *vasinfectum*. (In Russian). Ph .D. Thesis, All-Union Institute of Plant Protection, Leningrad, USSR.
- Hussein ,E.M.; Aly, A.A.; Ashour, A.Z.A. and Nasr, S.M.(1996)** Cluster analysis of serological protein patterns of three *Fusarium* species. *J. Agric .Sci . Mansoura Univ.*, 21: 3995- 4012 .
- Iannelli, D.;Capparelli, R.;Cristinzio, G.; Marziano, F.;Scala, F. and Noviella, C.(1982).** Serological differentiation among formae speciales and physiological races of *Fusarium oxysporum*. *Mycologia*, 74 (2): 313-319.
- Jack, A.L.;George, C.P. and Martha, D.H.(1993).** Biological control of damping-off of snapbeans caused by *Sclerotium rolfsii* in the green-house and field with formulations of *Gliocladium virens*. *Biological control* 3, 109-115.
- Jha, D.K. and Singh, D.K. (1997).** Biological control of chickpea wilt. *Indian Phytopathological Society Golden Jublee International Conference*, November 10-15 New Delhi, India.
-

REFERENCES

- Jharia, H.K. and Khare, M.N.(1986).**Biological control of *Rhizoctonia bataticola* causing diseases in soybean with *Bacillus subtilis* and *Bacillus* spp. abstracts of papers presented in seminar on management of soil borne diseases of crop plants, 8-10 January, 1986. Tamil Nadu Agricultural University, Coimbatore, p.32.
- Kappor, I.J. and Kar, B.(1989).**Antagonism of *Azotobacter* and *Bacillus* to *Fusarium oxysporum* f.sp.*Lycopersici*.Indian Phytopathology, 42: 400- 4004.
- Khalifa, E.Z. (1987).** Further studies on some soil-borne fungi affecting soybean and their control. Ph.D. thesis,Fac.of Agric.,Menoufiya Univ., Egypt, 148 pp.
- Kim, D.S.;Weller, D.M. and Cook, R.J.(1997).**Population dynamics of *Bacillus* sp. L324-92R12 and *Pseudomonas fluorescens* 2-79 RN10 in the Rhizosphere of Wheat .Phytopathology,87:559-564.
- Klaasen, J.A. and Marasas, W.F.(1998).**Comparative pathogenicity of four *Fusarium* species and *Gaeumannomyce sgraminis* var. *tritici* to wheat seedlings and mature plants. African Plant Protection, 4 (1) : 35-41.
- Korn Wendisch,F. and Kulzner,H.J.(1992).**The family Streptomycetaceae In: Balows A, Truper HG, Dworkin M, Harder W, Schleifer K-H (eds) The prokaryotes. A handbook on the biology of Bacteria: ecophysiology, isolation, identification, application, 2nd edn springer, Berlin Heidelberg New York, pp 921-995.
- Krasil'nikov, N.A.(1960).** Taxonomic principles in the Actinomycetes. Journal of Bacteriology. 79:65-74.
- Lacy, M.L. and Bridgman, G.H.(1962).** Phytopathology 52: 173.
- Laurell, C.B.(1965).** Antigen-Antibody Crossed-electrophoresis.Analytical Biochemistry, 10: 358-361.
- Lawn, D.A. and Sayre, K.D. (1992).**Soil-borne Pathogens on cereals in a highland location of Mexico.plant Dis.76,149-154.
- Lee, J.Y. and Hwang, B.K.(2002).**Diversity of antifungal Actinomycetes in various vegetative soils of korea. Can. J.Microbiol.48:407- 417.
-

REFERENCES

- Leur, J.A.G.van and Bailey, K.L.(2000).** The occurrence of barley root diseases in different agri-ecological zones of Syria. *Canadian Journal of Plant Pathology*,22 (1): 61-69
- Lewis, J.A.; Lumsden, R.D. and Locke, J.C. (1996).** Biocontrol of damping-off diseases caused by *Rhizoctonia solani* and *Pythium ultimum* with alginate prills of *Gliocladium virens*, *Trichoderma hamatum* and various food bases. *Biocontrol Science and Technology*. 6 (2): 163-173.
- Lewis, J.A. and Papavizas, G.C. (1987).** Application of *Trichoderma* and *Gliocladium* in alginate pellets for control of *Rhizoctonia* damping-off. *Plant Pathology*,36 (4): 438-446.
- Lewis, J.A.; Papavizas, G.C. and Lumsden, R.D. (1991).** Anew formulation system for the application of biocontrol fungi to soil. *Biocontrol Sci. Technol.* 1: 59-69.
- Leyns, F.; Lambert, B.; Joos, H. and Swings, J. (1990).** Antifungal Bacteria Data, 1990, 437-443.
- Liddell, C.M.(1985).** The comparative pathogenicity of *Fusarium graminearum* group 1, *Fusarium culmorum* and *Fusarium crookwellense* as crown, foot and root rot pathogens of wheat. *Australian Plant Pathology*,14 (2): 29-32.
- Lievens, K.H.;VanRijsbergen,R.;Leyns,F.R.;Lambert,B.;Tenning,P.; Swings,J. and Joos, H. (1989).** Dominant rhizosphere bacteria as a source for antifungal agents.*Pestic. Sci.*27,141-154.
- Liu, Z. and Sinclair,J.B.(1991).**Effect of seed coating with *Bacillus* spp. on *Rhizoctonia* damping-off, root and stem rot of soybean. *Biological and Cultural Tests for Control of Plant Diseases*,6: 62.
- Liu, Z. and Sinclair, J.B. (1992).** Population dynamics of *Bacillus megaterium* strain B153-2-2 in the rhizosphere of soybean. *Phytopathology*, 82:1297-1301.
- Liu, Z. and Sinclair, J.B. (1993).** Colonization of soybean roots by *Bacillus megaterium* B153-2-2. *Soil Biol. Biochem.*,25: 849-855.
- Lowry, O.H.; Rosebrough Nira, J.;Farr, A.L. and Randall Rose, J. (1951).**Protein measurment with the folin phenol reagent.*J. Biol.Chem.* 193: 265- 275.
-

REFERENCES

- Lumsden, R.D.; Locke, J.C.; Adkins, S.T.; Walter, J.F. and Ridout, C.J. (1992).** Isolation and localization of the antibiotic gliotoxin produced by *Gliocladium virens* from alginate prill in soil and soilless media. *Phytopathology*, 82 (2): 230-235.
- Mackinaite, R. (2001).** Root rot agents of cereal and their interrelation. *Beitrage zur Zuchtungsforchung Bundesanstalt fur Zuchtungsforchung an Kulturpflanzen*, 7 (1): 21-25.
- Mahafee, W.F. and Backman, P.A. (1993).** Effects of seed factors on spermosphere and rhizosphere colonization of cotton by *Bacillus subtilis* GB03. *Phytopathology*, 83:1120-1125.
- Mamluk, O.F.; Cetin, L.; Braun, H.J.; Bolat, N.; Bertschinger, L.; Makkouk, K. M.; Yildirim, A. F.; Saari, E. E.; Zencirci, N.; Albustan, S.; Ali, S.; Beniwal, S.P.S. and Dusunceli, F. (1997).** Current status of wheat and barley diseases in the Central Anatolian Plateau of Turkey. *Phytopathologia Mediterranea*. 36 (3): 167-181.
- Manka, M. and Fruzynska Jozwiak, D. (1996).** Biocontrol of greenhouse carnation *Fusarium* wilt with saprophytic forest soil fungi. *Folia Horticulturae*, 8 (1): 93-105.
- Maplestone, P.A. and Campbell, R. (1989).** Colonization of roots of wheat seedlings by bacilli proposed as biocontrol agents against take-all. *Soil Biol. Biochem.* 21: 543-550.
- Marziano, F.; Ianelli, D.; Cristinzio, G. and Noviello, C. (1981).** Serology in the systematics of the genus *Fusarium*. *Rivista di Pathologia Vegetale*, 17: 137-148.
- Massafumi, S.; Yoshiko, N.; Yukio, S.; Tamotsu, F.; Yasuhiro, I.; Hiroyasu, O.; Ryuji, Y. and Hitoshi, K. (2000).** Studies on Endophytic Actinomycetes (1) *Streptomyces* spp. isolated from *Rhododendron* and its antifungal activity. *Journal of General Plant Pathology*, 66 (4): 360-366.
- Mathivanan, N.; Srinivasan, K. and Chelliah, S. (1998).** Evaluation of different organic materials as carrier for formulated product of *Trichoderma*. *Journal of Biological Control*, 12 (1): 67-70.
- Mehiar, F.; El-Kady, S.; Gabr, M. and Hafez, Y.M. (2000).** Serological differences between virulent isolates and a virulent mutants of *Ralstonia (Pseudomonas) Solanacearum*. *J. Agric. Res., Tanta Univ.*, 26:122-130.
-

REFERENCES

- Milus, E.A. and Rothrock, C.S.(1993).** Rhizosphere colonization of wheat by selected soil bacteria over diverse environments. *Can.J.Microbiol*, 39: 335-341.
- Mishra, S.K.(1984).** Antigenic profile of some typical and septate phialide strains of *Aspergillus fumigatus*. *Sabouraudia*, 22 (2):91-100.
- Mierzwa ,Z .; Wiwart, M. and Fordonski, G.(1996) .** The production of antiseraum against *Fusarium avenaceum* conidial spores. *Biuletyn Instytutu Hodowlii Aklimatyzacji Roslin* No .197: 147-150
- Mohamed, Z.K. (1982).** Physiological and antagonistic activities of *Streptomyces* in rhizosphere of some plants. *Egyptian Journal of Phytopathology*, 14(1-2): 121-128 .
- Mohan, S.B. and Ride, J.P.(1982).** An immunoelectrophoretic approach to the identification of progressive and fluctuating isolates of the hop wilt fungus *Verticillium albo-atrum* .*J.Gen. Microbiol.* 128,255-265.
- Molan, Y.; Al-Doss, A.A. and El-Hussieni, S.(2001).** Evaluation of wheat genotypes for susceptibility to common root rot and foot rot diseases caused by *Bipolaris sorokiniana*, and *Fusarium graminearum*. *Assiut Journal of Agricultural Sciences*, 32 (5): 121-132.
- Morton, D.J. and Duke, P.D.(1967).** Serological differentiation of *Pythium aphanidermatum* from *Phytophthora parasitica* var. *nicotiana* and *P. parasitica*. *Nature*, 213-223.
- Ouchterlony, O. and Nilsson, A.L. (1978).** Immunodiffusion and immunoelectrophoresis, p.190. In "Handbook of Experimental Immunology" (Weir, D.M. ed.) 3rd Ed. Blackwell Scientific Publications, Oxford.
- Pelezar, M.J.; Chan, E.C.S. and Krieg, N.R.(1988).** The world of bacteria IV. Gram-positive filamentous bacteria of complex morphology. *Microbiology* McGraw Hill Book Company New York. PP, 320-330.
- Prasad, R.D. and Rangeswaran, R. (1999).** Granular formulation of *Trichoderma* and *Gliocladium* spp. in biocontrol of *Rhizoctonia solani* of chickpea. *Journal of Mycology and Plant Pathology*, 29(2): 222-226.
-

REFERENCES

- Raguchander, T.; Jayashree, K. and Samiyappan, R. (1997).** Management of *Fusarium* wilt of banana using antagonistic microorganisms. *Journal of Biological Control*, 11(1-2):101-105.
- Ramakrishnan, G.; Jeyarajan, R. and Dinakaran, D. (1994).** Talc-based formulation of *Trichoderma viride* for biocontrol of macrophomina phaseolina. *Biological control*, 8 (1): 41-44.
- Rewal, H.S. and Thooty, J.S. (1995).** Differential response of wheat varieties to systemic fungitoxics applied to *Ustilago tritici* (Pers) Rostr. *Indian J. Agric. Sci.*, 55: 548-549.
- Roberts, W.K. and Selitrennikoff, C.P. (1988).** Plant and bacterial chitinase differ in antifungal activity. *Journal of General Microbiology*, 134, 169-176.
- Rohacik, T. and Hudec, K. (2005).** Influence of agro-environmental factors on *Fusarium* infestation and population structure in wheat kernels. *Annals of Agricultural and Environmental Medicine*, 12 (1): 39-45.
- Rossi, V.; Cervi, C.; Chiusa, G. and Languasco, L. (1995).** Fungi associated with foot rots on winter wheat in Northwest Italy. *Journal of Phytopathology*, 143 (2):115-119.
- Rothrock, C.S. and Cunfer, B.M. (1986).** Absence of take-all decline in double-cropped fields. *Soil Biol. Biochem.* 18,113-114.
- Ryder, M.H.; Terrace, T.E.; Rovira, A.D.; Correll, R.L.; Yan, Z. and Tang, W. (1999).** Use of strains of *Bacillus* isolated in China to suppress take-all and *Rhizoctonia* root rot, and promote seedling growth of glass-house grown wheat in Australian soils. *Soil Biology and Biochemistry*, 31(1): 19-29.
- Sangle, U.R. and Bambawale, O.M. (2004).** New strains of *Trichoderma* spp. strongly antagonistic against *Fusarium oxysporum* f.sp. *sesami*. *Journal of Mycology and Plant Pathology*, 34 (1): 107-109.
- Singh, R.S. (1982).** *Plant pathogens "The fungi"* Oxford and IBH Publishing Co. New Delhi, Bombay, Calcutta, pp. 443.
- Song, Z. and Chen, H. (2002).** *In vitro* antagonism of *Trichoderma viride* against the soilborne fungal pathogens. *Journal of Fujian College of Forestry*, 22(3):219-222
-

REFERENCES

- Snedecor, J.W.(1965).**Statistical methods. Ames, Iowa, USA. Iowa State Univ. Press.
- Specht, L.P. and Rush, C.M.(1988).** Fungi associated with root and foot rot of winter wheat and populations of *Cochliobolus sativus* in the Texas Panhandle. Plant Disease, 72 (11): 959-963.
- Sreenivasaprasad, S. and Manibhushanrao, K. (1990).** Biocontrol potential of fungal antagonist *Gliocladium virens* and *Trichoderma longibrachiatum*. Zeitschrift fP flanzkrankheiten und-Pflanzenschutz, 97(6): 570-579.
- Strausbaugh, C.A.; Bradley, C.A.; Koehn, A.C. and Forster, R.L. (2004).**Survey of root diseases of wheat and barley in south eastern Idaho.Canadian Journal of Plant Pathology, 26 (2): 167-176.
- Torp, J. and Andersen, B. (1982).** Two-dimensional electrophoresis of proteins from cultures of *Erysiphe graminis* f.sp. *borderi*. Physiol. Plant Pathol. 21, 151-160.
- Toth, A.(1991).** The occurrence of *Fusarium* species in wheat samples from Pest county. Novenyvedelem,27 (2): 66-71.
- Turner, J.T. and Backman, P.A. (1991).**Factors relating to peanut yield increases after seed treatment with *Bacillus subtilis*. Plant Disease,75:347-353.
- Vafai, S.H.;Farrokhi-Nejad, R. and Darvishnia, M .(2001).** *Fusarium* species isolated from root and crown of wheat and barley in Khuzestan province.The Scientific Journal of Agriculture Vol.24 (1): 0254 – 3648.
- Vidhyasekaran, P. and Muthamilan, M. (1995).** Development of formulations of *Pseudomonas fluorescens* for control of chickpea wilt. Plant Dis.,79:782-786.
- Vidhyasekaran, P.;Sethuraman, K.; Rajappan, K. and Vasumathi, K. (1997).** Powder formulations of *Pseudomonas fluorescens* to control pigeonpea wilt. Biological control 8, 166-171.
- Wang, G.C.; Chen, H.K.; Xu, P.S. and Bao, J.Y.(1990).** Studies on the pathogens of the rice bakanae disease in Zhejiang. Acta Phytopathologica Sinica,20 (2): 93-97.
-

REFERENCES

- Waxman, S.A. (1959).** The actinomycetes vol I nature, occurrence and activities. The Williams and Wilkins Co., Baltimore.
- Wiese, M.V. (1987).** Compendium of Wheat Diseases. Second Edition. APS Press, St. Paul, Minnesota.
- Wilcoxson, R.D.; Kommedahl, T.; Ozmon, E.A. and Windels, C.E. (1988).** Occurrence of *Fusarium* species in scabby wheat from Minnesota and their pathogenicity to wheat. *Phytopathology*, 78(5): 586-589.
- Windels, C.E. and Wiersma, J.V. (1992).** Incidence of *Bipolaris* and *Fusarium* on subcrown internodes of spring barley and wheat grown in continuous conservation tillage. *Phytopathology*, 82 (6): 699-705.
- Wiwart, M. (1991).** The similarity of antigenic proteins of *Fusarium nivale* (Fr.) Ces isolates. *Mycotoxin Research*, 7(A(Suppl.) Part 1): 54-57.
- Wuike, R.V.; Shingare, V.S. and Dawane, P.N. (1998).** Biological control of wilt of sesame caused by *Fusarium oxysporum* f.sp. *sesame*. *Journal of Soils and Crops*, 8 (1): 103-104.
- Wyk, P.S. van; Los, O.; Pauer, G.D.C. and Marasas, W.F. (1987).** Geographic distribution and pathogenicity of *Fusarium* species associated with crown rot of wheat in The Orange Free State, South Africa. *Phytophylactica*, 19 (30): 271-274.
- Wyk, P.S. van; Los, O. and Marasas, W.F. (1988).** Pathogenicity of a new *Fusarium* sp. from crown rot of wheat in South Africa. *Phytophylactica*, 20 (1): 73-75.
- Zare, R. and Ershad, D. (1997).** *Fusarium* species isolated from cereals in Gorgan area. *Iranian Journal of Plant Pathology*, 33 (1/2): 1-14 (Pe), 1-4 (En).
-
-