

RÉFÉRENCES.

REFERENCES

- Admec, O., Matis, J. and Galvanek, M. (1962): Lancet, 1, 81, 1962. Cited by Sallam, 1966. A contribution to the utilisation of steroid sapogenins. Ph.D. Thesis, Faculty of Science Cairo University.
- Allam, A.M. and El-Kady, I.A. (1974): The 11 α - and 17 α -progesterone hydroxylases of Rhizopus nigricans, NRRL 1477. Acta. Microbiol. Pol. Ser. A Microbiol. Gen. 7(1): 41-44.
- American Cyanamide Co. (1962): Hydroxylation of steroids. Brit Patent, 889, 853, Feb. 21
- American Cyanamide Co. (1963): 17 α -Deoxy steroids. Brit. Patent, 935, 701, Sept.
- Asai, T., Aida, K., Ohki, E., Tanaka, T. and Hattori, M. (1958 a): Hydroxylation of steroid by microorganisms. I. Hydroxylation of progesterone by Syncephalastrum. Nippon Nogei Kagaku Kaishi, 32, 723-8.
- Asai, T., Tsuda, K., Aida, K., Ohki, E., Tanaka, T., Hattori, M. and Machida, H. (1958b): Microbiological hydroxylation of steroids. I. Microbiological hydroxylation of progesterone by Rhizopus. J. Gen. Appl. Microbiol. (Tokyo), 4, 63 - 6.
- Asai, T., Aida, K., Tanaka, T., Ohki, E., Matsuhisa, T., Takeda, Y. and Inui, T. (1959 a): Microbiological hydroxylation of steroids. IX - Hydroxylation of steroids by Syncephalastrum racemosum. J. Gen. Appl. Microbiol. (Tokyo), 5, 127 - 37.

- Asai, T., Aida, K., Tanaka, T., Ohki, E., Matsuhisa, T.
Takeda, Y. and Inui, T. (1959 b): Hydroxylation
of steroids by microorganisms. II-Hydroxylation
of steroids by Syncephalastrum racemosum, Nippon
Nogei Kagaku kaishi, 33 , 985 - 90.
- Bennet, R.D. and Heftmann, E. (1962): Thin - layer chroma-
tography of corticosteroids. J. Chromatog., 2,
348 - 52.
- Bernstein, S., Lenhard, R.H., Allen, W.S., Heller, R., Little,
R., Stolar, S.M., Feldman, L.I. and Blank, R.H. (1956):
16-Hydroxylated steroids. IV- The synthesis of the
16 α -hydroxy derivatives of 9 α -halo steroids. J.
Amer. Chem. Soc., 78 , 5693.
- Boedijn, K.B. (1958): Notes on the Mucorales of Indonesia.
Sydowia, 12 : 321 - 362.
- Brannon, D.R., Mortin, J., Oehlschlager, A.C., Durham, N.N.
and Zalkow, L.H. (1965): Transformation of proges-
terone and related steroids by Aspergillus tamarii.
J. Org. Chem., 30 (3), 760-2.
- Campillo, C.C. (1964): 11-Hydroxylation of steroids. U.S.
Patent, 3, 119, 748, Jan. 28.
- Capek, A. and Hanc, O. (1960): Microbiological transformation
of steroids. XII -Transformation of steroids by
different spp. and strains of Fusaria. Folia
Microbiol., 5 , 251.

- Capek, A. and Hanc, O. (1961): Microbiological transformation of steroids. XIV - Microbiol., 6, 237.
- Capek, A. and Hanc, O. (1962): Microbiological transformation. XVII - Transformation of progesterone by various species and strains of Penicillium. Folia Microbiol., 7, 121.
- Capek, A., Tadra, M. and Tuma, J. (1964): Microbiological transformation of steroids. XXIII - Separation of androstane 17-hydroxy Epimers. Folia Microbiol., 9, 380-82.
- Capek, A., Hanc, O., Kakac, B. and Tadra, M. (1966 a): Preparing 11 α -hydroxylated steroids. Czech. Patent, 120, 585, Nov. 15.
- Capek, A., Hanc, O. and Tadra, M. (1966 b): Microbiol transformation of steroids., Dr. Junk, W., Publishers, the Hague - the Netherlands.
- Carvajal, F. (1960): 6 β - Hydroxy steroids. German Patent, 1, 095, 278, Dec. 22.
- Charney, W., Herzog, H.L. and Sutter, D. (1962): 15 β -Hydroxy adrenocorticoids. U.S. Patent, 3, 014, 051.
- CIBA Ltd. (1959): Polyoxygenated steroids. Brit. Patent, 816, 922, July 22.
- Clements, F.E. and Shear, C.L. (1931): Genera Publishing Co. : New York.

- Colingsworth, D.R., Brunner, M.P. and Haines, W.J. (1952): A partial microbiological synthesis of adrenal cortex hormones. J. Amer. Soc., 74, 2381 - 2.
- Colingsworth, D.R., Karnemaat, J.N., Hanson, F.R., Brunner, M.P., Mann, K.M. and Haines, W.J. (1953): A partial microbiological synthesis of hydrocortisone. J. Biol. Chem., 203, 807.
- Djerassi, C. (1955): Cited by Peterson, 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.). Eds., Academic Press. London and New York, 708 PP.
- Dodson, R.M., Goldkamp, A.H. and Muir, R.D. (1957): Microbiological hydroxylation of steroids at positions C-1 and C-2. J. Amer. Chem. Soc., 79, 3921.
- Dodson, R.M. and Muir, R.D. (1958 a): Microbiological transformations. II- The microbiological aromatization of steroids. J. Amer. Chem. Soc. 80, 5004.
- Dodson, R.M. and Muir, R.D. (1958 b): Microbiological transformations. III- The hydroxylation of steroids at C-9. J. Amer. Chem. Soc., 80, 6148.
- Dodson, R.M., Nicholson, R.T. and Muir, R.D. (1959): Microbiological transformations. IV - The oxidation of dehydroisoandrosterone at C-7. J. Amer. Chem. Soc., 81, 6295.

- Dodson, R.M., Kraych, S., Nicholson, R.T. and Mizuba, S.
(1962): Microbiological transformations. IX-1B
Hydroxylation of androstendione. J. Org. Chem.,
27, 3159-64.
- Dodson, R.M., Langbein, G., Muir, R.D., Schubert, A., Siebert,
R., Tamm, Ch. and Weiss-Berg, E. (1965): Reactions
with microorganisms. XIV-Identify of the steroids des-
cribed as 6 β , 15 α -dihydroxyprogesterone; 11 α , 15 β -
dihydroxyprogesterone and 12 β , 15 β -dihydroxyprogesterone.
Helv. Chim. Acta, 48 (8), 1933 - 40.
- Donin, M.N., Fried, J. and Thoma, R.W. (1959): 11; 15-Dioxy-
genated progesterone. U.S. Patent, 2, 881, 189,
Apr. 7.
- Dorfmann, R.I. (1958): Biochemistry of steroid hormones. Ann.
Rev. Biochem., 26, 523.
- Dulaney, E.L. McAleer, W.J., Koslowski, M., Stapley, E. and
Jalom, J. (1955): Hydroxylation of progesterone and
11-desoxy -17-hydroxycorticosterone by Aspergillus and
Penicillium. Appl. Microbiol., 3, 336.
- Dulaney, E.L., McAleer, W.J. and Stoudt, T.H. (1960): 16-
Hydroxyprogesterone and derivatives. U.S. Patent, 2,
960, 434, Nov. 15.
- Dulaney, E.L. and McAleer, W.J. (1961): 11,17-Dihydroxysteroids.
U.S. Patent, 2, 970, 085, Jan. 31.

- Dumazert, C., Chiglione, C. and Pugnet, T. (1963): Thin-layer chromatography of some steroids. Ann. Pharm. Franc, 21, 227-32.
- El-Kady, I.A., Sallam, L.A.R. and El-Refai, A.H. (1972): 17α -Hydroxylation of progesterone by microorganisms, (1972). Pakistan Journal of Biochemistry, V 2, 39-43.
- El-Kady, I.A. and Allam, A.M. (1973): Hydroxylation of progesterone by extracts of R. nigricans NRRL. 1477. J. Gen. Microbiol. 77, 465-469.
- El-Refai, A.H., Sallam, L.A.R. and El-Kady, I.A. (1969): Microbiological transformation of progesterone, (1969), J. Gen. Appl. Microbiol. 15, 301-307.
- El-Refai, A.H., Sallam, L.A.R. and El-Kady; I.A. (1970 a): Studies on the mechanisms of progesterone transformation with Rhizopus nigricans. J. Gen. Appl. Microbiol. 16, 137 - 144.
- El-Refai, A.H., Sallam, L.A.R. and El-kady, I.A. (1970 b): Transformation of progesterone by R. nigricans REF 129 as influenced by modification of fermentation medium Bull. Chem. Soc. (Japan), 43, 3592 - 3599.
- El-Refai, A.H., Abdel Monem H., Lotfy A.R., Sallam, L.A.R. and Nadia Naim (1974): Transformation of cortisol with representative fungi. J. Gen. Appl. Microbiol. 20 (2): 111-115.

- Eppstein, S.H., Meister, P.D., Peterson, D.H., Murray, H.C.,
Leigh, H.M., Weintraub, A., Reineke, L.M. and Meeks,
R.C. (1958): Microbiological transformation of steroids. I
XV-Tertiary hydroxylation of steroids. J. Am. Chem. Soc.
80, 3382-3389.
- Feldman, L.I. and Allen, W.S. (1957): 11 β -Hydroxylation of
steroids. U.S. Patent, 2, 789, 940, Apr. 23.
- Feldman, L.I., Pruess, L.M. and Rigler, N.E. (1962): 1-Hydro-
xylated steroids. U.S. Patent, 3, 309, 937, June 19.
- Fried, J., Thoma, R.W., Gerke, J.R., Herz, J.E., Danin, M.N.
and Perlman, D. (1952): Oxidation of steroids by micro-
organisms. II-Hydroxylation in position 11 and synthesis
of cortisone from Reichstein's compounds S. J. Amer.
Chem. Soc. 74, 3962.
- Fried, J., Perlman, D., Klingsberg, A. and Bayan, A.P. (1955 a):
Cited by Fried et al., 1955. Recent. Prog. Hormone
Research, 11, 157.
- Fried, J., Thoma, R.W., Sabo, E.F. and Grabowicz, P. (1955 b):
Cited by Fried et al., 1955. Recent. Prog. Hormone
Research, 11, 157.
- Fried, J., Thoma, R.W., Danin, M.N. and Gerke, J.R. (1955 c):
Cited by Fried et al., 1955. Recent. Prog. Hormone
Research, 11, 157.
- Fried, J. and Borman, A. (1958): Synthetic derivatives of cortical
hormones. Vitamins and Hormones 16, 303.

- Fried, J., et al. (1958): Personal communication. Cited by Peterson, 1963. In Biochemistry of industrial microorganisms, 537-606 (Rainbow, C. and Rose, A.H.), Academic Press. London and New York), 708 PP.
- Fried, J. Thoma, R.W., Perlman, D. and Gerke, J.R. (1959 a): 15-Hydroxyprogesterone. U.S. Patent, 2,879, 280, Mar. 24.
- Fried, J., Perlman, D., Langlykke, A.F. and Titus, E.O. (1959 b): Microbiological oxidation of steroids. U.S. Patent, 2, 872, 381, Feb. 3. Continuation-in-part of U.S. Patent, 2, 709, 705.
- Ganch, O. and Chapek, A. (1960): Microbiological transformation of steroids. Pharmacotherapentica, 1950-59, collected papers. Aniversary Research Inst. Pharm. Biochem., 10 th, Prague, 1960, 157 - 71.
- Gilman, J.C. (1957): A manual of soil fungi, Iowa, State Univ. Press. Ames, Iowa, U.S.A.
- Goodman, J.J. and Smith, L.L. (1961): 16 α -Hydroxy steroids. XI-2B- and 16 α -hydroxylation of 8 α -fluorohydrocortisone by strains of Streptomyces roseochromogenes. Appl. Microbiol., 2, 372-5.
- Greenspan, G., Schaffner, C.P., Charney, W., Herzog, H.L. and Hershberg, E.B. (1959): Transformation of steroids by fungi. Introduction of 1 α and 2B -hydroxyl groups into Reichstein's compound S. J. Amer. Chem. Soc. 79, 3922.

- Gubler, A. and Tamm, Ch. (1958): Unwandlung von Cardenoliden durch Mikroorganismen: 1, 12 β -hydroxylierung von digitoxigenin. *Helv. Chim. Acta*, 41, 297-301.
- Gundu Rao, P. (1963): Microbiological transformation of steroids with Fusarium roseum. *Indian J. Pharm.*, 25, 131-5.
- Hanson, F.R., Mann, K.M., Nielson, E.D., Anderson, H.V., Brunner, M.P., Karnemaat, J.N., Colingsworth, D.R. and Haines, W.J. (1953): Microbiological transformations of steroids. VIII-Preparation of 17 α -hydroxycortico-sterone. *J. Amer. Chem. Soc.*, 75, 5369.
- Hasegawa, T. and Nishikawa, M. (1958 a) Oxidation of steroids. Japan Patent, 31, Jan. 10.
- Hasegawa, T. and Takahashi, T. (1958 b): *Bull. Agric. Chem. Soc. (Japan)*, 22, 212. Cited by Peterson, 1963. cf. Ref. 147.
- Hasegawa, T., Takahashi, T., Sakano, I., Nishikawa, M., Hagiwara, H. and Noguchi, S. (1958 c): Oxidation of steroids. Japan patent, 7372, Aug. 27.
- Hasegawa, T., Takahashi, K., Sakano, I., Nishikawa, M., Morita, K. and Noguchi, S. (1959 a): Oxidation of steroids-Japan Patent, 425, Feb. 5.
- Hasegawa, T., Takahashi, K., Itano, I., Nishikawa, M., Hagiwara, H. and Noguchi, S. (1959 b): Oxidation of steroids. Japan Patent, 529, Feb. 9.
- Hasegawa, T. and Takahashi, T. (1960): Microbiological transformation of steroids. V-Oxidation of steroids by Polyporus orientalis. *Takeda kenkyusho Nempo*, 12, 164-71.

- Hasegawa, T., Takahashi, K. and Uchibori, S. (1963): 19-Hydroxy steroids. Japan Patent, 2279, Mar. 18.
- Hayano, M. (1962): In "the oxygenases", 181, (Hayaishi, O., Ed., Acad. Press., New York and London, 588 PP.
- Hechter, O., Jacobson, R.P., Jeanloz, R., Levy, H., Marshall, C.W., Pincus, G. and Schenker, V. (1949): Biooxygenation of 11 desoxycorticosterone at C-11. J. Amer. Chem. Soc., 71, 3261.
- Helena, S. and Igor, B. (1958): Separation and identification of steroids produced by fermentative oxidation of progesterone. Fresenius Z. Anal. Chem., 243, 291-4.
- Hems. (1953): J. Pharm. Lond., 5, 409, 1953. A Contribution to the utilisation of steroid sapogenins. Ph.D. Thesis, Faculty of science Cairo University.
- Hench et al. (1949): Proc. Staff. Meeting May clinic, 24, 277, (1949). A contribution to the utilisation of steroid sapogenins. Ph. D. Thesis, Faculty of science Cairo University.
- Herzog, H.L., Gentles, M.J., Hershberg, E.B., Carvajal, F., Sutter, D., Charney, W. and Sheffner, C.P. (1957): Microbiological transformation of steroids. 2B - Hydroxylation. J. Amer. Chem. Soc., 79, 3921.
- Herzog, H.L., Gentles, M.J., Basch, A., Coscarelli, W., Zeitz, M.E.A. and Charney, W. (1960): Microbiological transformation of steroids. VIII-16B-Hydroxylation and other transformations of Wojnowicia graminis. J. Org. Chem., 25, 2177.

- Hromatka, O. and Ebner, H. (1951): Untersuchungen über die Essiggärung. II-Sauerstoffmangel der Fesselgärung
Intrestigations into vinegar production .
II-Oxygen deficiency of the fermentation generator.
- Iizuka, H., Natio, A. and Hattori, M. (1958): Microbiological hydroxylation of steroids. II-Microbiological hydroxylation of progesterone by Aspergillus. J.Gen. Appl. Microbiol. (Tokyo), 4, 67-78.
- Iizuka, H. and Naito, A. (1960 a): Microbiological hydroxylation of steroids. XII-Oxidation of progesterone and Reichstein's compound S by genus Pseudomonas and others. J. Gen. Appl. Microbiol. 6, 169.
- Iizuka, H. and Naito, A. (1960 b): Production of 6 β , 14 α - dihydroxyprogesterone and 6 β -hydroxy compound S by bacteria. Nippon Nogeikagaku kaishi, 34, 472-5.
- Ilavsky, J. and Herzog, H.L. (1961): 11 α -Hydroxylation of steroids by Beauveria. U.S. Patent, 3,013, 945, Dec.19.
- Ishu, H., Nozaki, Y., Okomura, T. and Sato, D. (1961): J. Agric. Chem. Soc. (Japan), 81, 1051, 1961. Cited by Peterson, 1963. of. Ref. 147.
- Kahbt, F.W., Meyster, Ch., Neher, R., Vischer, E. and Wettstein, A. (1952): Biologische hydroxylierungen von steroiden. Experientia, 8, 422.
- Karow, E.O. and Petsiavas, D.W. (1956): Effect of physical variables on microbiological transformation of steroids. Industr. Engng. Chem., 48, 2213.

Kemiske, L. Fabrik Ved A kongsted (1961): 11 α -Hydroxysteroids by microbiological oxidation. Dan. Patent, 91, 270, Oct. 2.

Kemiske, L. Fabrik Ved A Kongsted (1962): 11 α -Hydroxysteroids by microbiological oxidation. Dan. Patent, 94, 041, Nov. 5.

Kita, D.A. (1962): Oxygenation of steroid with Streptomyces halstedii. U.S. Patent, 2, 991, 230.

Kluger, B., Siebert, R. and Schubert, A. (1957): Microbial oxidation of steroids in the 15 α -position by Fusaria. Naturwissen Schäften, 44, 40.

Kogan, L.M., Elin, E.A., Barmenkov, A.S. and Torgov, I.V. (1964): Microbiological conversion of steroid. IV- Oxidation of pregnenolone with Rhizopus nigricans. Izv. Akad. Nauk, SSSR, Ser. Khim, (11).

Kogan, L.M., Voishvillo, N.E., Skryabin, G.K. and Torgov, I.V. (1965): Preparation of 17 α -hydroxyprogesterone. U.S.S.R. Patent, 170, 981.

Kogan, L.M., Voishvillo, N.E., Elin, E.A., Skryabin, G.K. and Torgov, I.V. (1966): Preparation of 16 α -hydroxy-steroids. U.S.S.R. Patent, 185, 344, Aug. 13.

Kogan, L.M., Volkova, I.M., Torgov, I.V. and Skryabin, G.K. (1968): 16 α -Hydroxysteroids. U.S.S.R. Patent, 231 717, (cl.A. 61 K), Nov. 28.

Kondo, E. and Tori, K. (1966): Dihydroxyandrostendione. Japan Patent, 8, 504, Oct. 24.

- Kramli, A. and Horva'th, J. (1948): Microbiological oxidation of sterols. Nature, London, 162 , 619.
- Kramli, A. and Horva'th, J. (1949): Microbiological oxidation of sterols. Nature, London, 163 , 219.
- Kurosawa, Y. and Sawai, M. (1963): Oxidation of steroids. Japan Patent, 25, 379.
- Laskin, A.I., Guiducci, M.A. and Fried, J. (1961): Synthesis of steroids with Kabatiella phoradendri. U.S. Patent, 2, 977, 286, Mar. 28.
- Laskin, A.I. and Fried, J. (1963): Microbial 6 β -hydroxylation of steroids containing aromatic A rings. U.S. Patent, 3, 115, 443.
- Leigh et al. (1956): Cited by Eppstein, et al., 1956. Microbiological transformation of steroid and their application to the synthesis of hormones. Vitamins and Hormones, 14, 359-432.
- Leudemann, G., Charney, W., Gentles, M.J., Marshall, H., Herzog, H.L. and Woyciesjes, A. (1960). Bact. Proc. A 21, 1960. Cited by Peterson, 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York , 708 PP.
- Leudemann, G., Charney, W., Woyciesjes, A., Peterson, E., Peckham, Gentles, M.J., Marshall, H. and Herzog, H.L. (1961): Microbiological transformation of Reichstein's compound S by Soenedesmus sp. Diketopiperazine metabolites from Soenedesmus sp. J. Org. Chem., 26, 4128.

- Levy, R.H. and Kushinsky, S. (1955): Arch. Biochem. Biophys., 55, 290, 1955. Cited by Peterson, 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York), 708 PP.
- Lisboa, B.P. (1964): Characterization of Δ^4 -3-oxo-C-21 steroids on thin-layer chromatograms "in situ" colour reactions. J. Chromatog., 16 (1), 136-51.
- Lisboa, B.P. (1965): Thin-layer chromatography of steroids. J. Pharm. Belg., 20, (11-12), 435-49.
- Mamoli, L. and Vercellone, A. (1937 a): Biochemische Umwandlung von Δ^4 -Androstendione in Δ^4 -Testosterone. Ein Beitrag zur Genese der Keimdrüsenhormone. Vorläufige Mitteilung. Ber., 70, 470.
- Mamoli, L. and Vercellone, A. (1937 b): Biochemische umwandlung von Δ^5 -Androstendione in Isoandrosterandiol und Δ^4 -Testosterone. Weiterer Beitrag zur Genese der Keimdrüsenhormone. Ber., 70, 2079.
- Mamoli, L. and Vercellone, A. (1937 c): Z. Physiol. Chem., 245, 39; 1937. Cited by Peterson, 1963. In Biochemistry of industrial microorganisms., 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York, 708 PP.
- Mann, K.M., Hanson, F.R., O'Connell, P.W., Anderson, H.V., Brunner, M.P. and Karnamaat, J.N. (1955): Studies of the microbiological of steroids by Cunninghamella blakesleeana H-334. I- The effect of alcohol and phenol, Appl. Microbiol., 3, 14.

- Marker, R.E., Wagner, R.B., Ulshafer, P.R., Wittbecker, E.L., Goldsmith, P.J. and Rouf, C.H. (1947): Steroidal sapogenins. J. Amer. Chem. Soc., 69, 2167, 1947., and previous communications.
- Mason, H.L., Meyers, C.S. and Kendall, E.C. (1936): Chemical studies of the suprarenal cortex. II- The identification of a substance which possesses the qualitative of cortin; its conversion into a diketone closely related to androstendione. J. Biol. Chem., 114, 613, 1936. J. Biol. Chem., 116, 276 .
- Masuo, E. and Kondo, J? (1963): Hydroxysteroids. Japan Patent, 16, 475, Aug. 30.
- McAleer, W.J. and Dulaney, E.L. (1956): Transformation of progesterone to 11-deoxycorticosterone by Wojnowicia graminis. Arch. Biophys., 62 , 109.
- McAleer, W.J. and Dulaney, E.L. (1958): Microbial polyhydroxylation of steroids. U.S. Patent, 2, 831, 798, Apr. 22.
- McAleer, W.J., Jacob, T.A., Turnbull, L.B., Schoenwaldt, E.F. and Stoudt, T.H. (1958): Hydroxylation of progesterone by Bacillus megaterium. Arch. Biochem Biophys., 73, 127.
- Meister, P.D., Eppstein, S.H., Peterson, D.H., Murray, H.C., Osborn, H.M.L., Weintraub, A., Reineke, L.M. and Meeks, R.C. (1953 a): Abstracts of the 123 rd Meeting of the Amer. Chem. Soc., Los Angeles P.5c. Cited by Peterson, 1963. In Biochemistry of industrial micro-organisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York , 708PP.

- Meister, P.D., Peterson, D.H., Murray, H.C., Eppstein, S.H.,
Reineke, L.M., Weintraub, A. and Leigh, H.M. (1953 b):
Microbiological transformations of steroids. II- The
preparation of 11 α -hydroxy-17 α -progesterone. J.
Amer. Chem. Soc., 75, 55.
- Meister, P.D., Reineke, L.M., Meeks, R.C., Murray, H.C.,
Eppstein, S.H., Osborn, H.M.L., Weintraub, A. and
Peterson, D.H. (1954 a): Microbiological transforma-
tions of steroids. XII-17 α -Hydroxylation. J. Amer.
Chem. Soc., 76, 4050.
- Meister, P.D., Reineke, L.M., Meeks, R.C., Murray, H.C.,
Osborn, H.M.L., Weintraub, A. and Peterron, D.H.
(1954 b): Cited by Eppstein et al., 1956. Microbiol-
ogical transformations of steroid and their application
to the synthesis of hormones. Vitamins and Hormones,
14, 359-432.
- Metz, H. (1961): Thin-layer chromatography for rapid assays of
enzymic steroid transformations. Naturwissenschaften,
48, 569.
- Meyer, A.S. (1955): 19-Hydroxylation of Δ^4 - androstene-3;17-
dione and dehydroepiandrosterone by Bovine adrenals.
Experientia, 11, 99.
- Meyster, Ch., Vischer, E. and Wettstein, A. (1954): Micro-
biologische hydroxylierungen von steroiden der 17 α -
und 21-stellung. Helv. Chim. Acta, 37, 1548.

- Meyster, Ch., Vischer, E. and Wettstein, A. (1955): Mikrobiologische hydroxylierungen von cortexon in der 7 α -; 15 α - oder 15 β - stellung. *Helv. Chim. Acta*, 38, 381.
- Minlon, H., Kin, Z.W., Wu, C.H. and Chow, Z.Z. (1959): Synthesis of hydroxortisone acetate. *Hua Hsueh Pao*, 25, 302 - 7.
- Murray, H.C. and Peterson, D.H. (1950): In "Perspectives and Horizons in Microbiology", Waksman, S.A. ed., P. 121. Rutgers University Press, New Brunswick, N.J.
- Murray, H.C. and Peterson, D.H. (1952): Oxygenation of steroids by Mucorales fungi. U.S. Patent, 2, 602, 769, July 8.
- Murray, H.C., and Reineke, L.M. (1962): 11 α , 17 α -Dihydroxyprogesterone. U.S. Patent, 3, 011, 951.
- Nawa, H., Uchibayashi, M., Kamiya, T., Yamano, Y., Hideo, A. and Abe, M. (1959): Microbiology. Microbiological transformation of cardiac Aglycone. *Nature, Lond.*, 184, 469.
- Nienstedt, W. (1964): A typical adsorption chromatography of steroids. *Biochem. J.*, 92 (1), 8p.
- Nishikawa, M. and Hagiwara, H. (1958 a): *Chem. Pharm. Bull.* 6, 226, cf. Ref. 147.
- Nishikawa, M. and Hagiwara, H. (1958 b): Steroids. VII- Microbiological oxidation of progesterone and 17 α -hydroxycorticosterone by Stemphylium botrysum. *J. Pharm. Soc., Japan*, 78, 1256.

- Nishikawa, M., Noguchi, S., Hasegawa, T., Takahashi, K. and Sakano, I. (1959): Oxidation of steroids Japan Patent, 426, Feb. 5.
- Nozaki, Y. (1961 a): Transformation of cardio-active steroids by microorganisms. Part I-Microbial dehydrogenation of 3-hydroxy group of 3-hydroxy group of digitalis cardiac aglycone. Agr. Biol. Chem. Japan, 25, 461.
- Nozaki, Y. (1961 b): Agr. Biol. Chem. Japan, 25 559. Cited by Peterson et al., 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds, Academic Press. London and New York, 708 pp.
- Nozaki, Y. and Okumura, T. (1961): Microbial 1 β -hydroxylation of digitoxigenin. Agr. Biol. Chem. Japan, 25, 515.
- O'Connell, P.W., Mann, K.M., Nielson, E.D. and Hanson, F.R. (1955): Studies of the microbiological oxidation of steroids by Cunninghamella blakesleeana H-334. II-Medium design. Appl. Microbiol., 3, 16.
- Okada, M., Yamada, A. and Ishidate, M. (1965): Microbiological transformation of dehydroepiandrosterone, progesterone, deoxycorticosterone and testosterone by Gibrella saubinett. Yakugaku Zasshi, 85 (9), 816-22.
- Okumura, T., Nozaki, Y. and Sato, D. (1964): Digitalis glucosides. XIX- Microbiological transformation of digitoxegenin derivatives by Absidia orchidis. Chem. Pharm. Bull. (Tokyo), 12 (10), 1143-51.

Perlman, D., Titus, E. and Fried, J. (1952): Microbiological hydroxylation of progesterone .J. Amer. Chem. Soc. 74, 2126.

Perlman, D., O'Brien, E., Bayan, A.P. and Green field, R.B. Jr. (1955): Antibiotic and vitamin B₁₂ production by a steroid oxidizing actinomycete. J. Bact. 69, 347.

Perlman, D. (1956): Abs. 130 th Meeting Amer. Chem. Soc. P.33
A. Cited by Peterson, 1963. of. Ref. 147.

Perlman, D., Weinstein, A.J. and Peterson, G.E. (1957); Canad. J. Microbiol., 3, 841, 1957. Cited by Peterson, 1963. cf. Ref. 147.

Peterson, D.H. and Murray, H.C. (1952): Microbial oxygenation of steroids at C-11. J. Amer. Chem. Soc., 74, 1871-72.

Peterson, D.H., Murray, H.C., Eppstein, S.H., Reineke, L.M., Weintraub, A., Meister, P.D. and Leigh, H.M. (1952): Microbiological transformation of steroids. I-Intruduction of oxygen at C-11 of progesterone. J. Amer. Soc., 74, 5933.

Peterson, D.H., Eppstein, S.H., Meister, P.D., Murray, H.C., Leigh, H.M., Weintraub, A. and Reineke, L.M. (1953 a): Microbiological transformations of steroids. IX-Degradation of C-21 steroids to C-19 ketones and to testosterone. J. Amer. Chem. Soc., 75, 5768.

- Peterson, D.H., Eppstein, S.H., Meister, P.D., Magerlein, B.J., Murray, H.C., Leigh, H.M., Weintraub, A. and Reineke, L.M. (1953 b): The 11-epimer of compound F and other new oxygenated derivatives of compound S. A new route to cortisone. J. Amer. Chem. Soc., 75, 412.
- Peterson, D.H., Murray, H.C. and Reineke, L.M. (1956): Microbiological transformation of steroids and their application to the synthesis of hormones. Vitamins and Hormones, 14, 359-432.
- Peterson, D.H. (1963): In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A. H.), Eds. Academic Press. London and New York), 708 PP.
- Pfizer, C. and Co. (1957 a): 11B-Hydroxylation of steroids. Brit. Patent, 769, 998, Mar. 13.
- Pfizer, C. and Co. (1957 b): Steroids. Brit. Patent, 783, 048, Sept. 18.
- Povah, A.H.W. (1917): A critical study Mucor. (Bull. Torrey Bot. Cl. 44, 241-259, 287-313).
- Principe, P.A. and Diassi, P.A. (1963): 9 α -Hydroxylation of steroids. U.S. Patent, 3, 080,298, Mar. 5.
- Prochazka, Z. and Sorm, F. (1965): Microbial hydroxylation of 14B, 18-cycloprogesterone. Collection Czech. Chem. Commun, 30 (6), 1874-9.

- Raspé, G. and Kieslich, K. (1961): *Naturwissen*, 48, 479.
In *Biochemistry of industrial microorganisms*,
537-606, (Rainbow, C. and Rose, A.H), Eds., Academic
Press. London and New York, 708 PP.
- Reichstein (1936): *Upper Bestandteile der Nebennieren-Rinde*.
VI. -Trennungsmethoden, Sowie Isolierung der Subs-
tanzen F. a Hund. J. Helf. Chim Acta, 19, 1107.
- Repole, R., Gasbarro, R., Maglione, S. and Rossi, P. (1965):
Detection of androgens by the use of sodium fluoresce-
in on thin-layer chromatography plates. *Rass.Med*.
Sper., 12 (3), 133-8.
- Reulas, J.P., Iriarte, J., Kincl, F. and Djerassi, C.J. (1958):
Steroids. XCVIII-Synthesis of some 10 β -hydroxy-19-
norsteroids. *J. Org. Chem.* 23, 1744.
- Roland, J.F. and Weiner, B.A. (1955): Bio-oxygenation of
progesterone by Mushrooms. *Science*, 121, 803.
- Rsenkranz, and Sondheimer, (1953): *Fortschr. Chem. Organ*.
Naturstoffe, 10, 274, 1953. Cited by Sallam, 1966
of. Ref. 161.
- Routin, J.B. and Shull, G.M. (1959): 11 β -Hydroxylation of
steroids by Epicoccum. U.S. Patent, 2,875, 134, Feb. 24.
- Rubin, B.A., Compillo, C.O., Hendricks, G., Cordova, F. and
Zaffaroni, A. (1956): *Bact. Proc.*, 33, 1956. Cited
by Peterson, 1963. In *Biochemistry of industrial
microorganisms*, 537-606, (Rainbow, C. and Rose, A.H.),
Eds., Academic Press. London and New York, 708 PP.

- Sallam, L.A.R. (1966): A contribution to the utilisation of steroid sapogenins. Ph.D. Thesis, Faculty of Science Cairo University.
- Sallam, L.A.R., El-Kady, I.A. and El-Refai, A.H. (1974): Some Culture conditions influencing the 17 α -hydroxylation of progesterone with Trichothecium roseum, Microbiol. Espan, 27, 299-308.
- Sardinas, J.L. (1962): Hydroxylating steroid. U.S. Patent, 3,032,475, May 1.
- Sarett, L.H. (1946): The structure of some derivatives of 3 α -hydroxy $\Delta^9,11$ -cholonic acid. J. Biol. Chem., 162, 591.
- Sato, Y., Tanaka, T., Kato, M. and Tsuda, K. (1963): Microbiological hydroxylation of steroids. XX-Hydroxylation of progesterone with Syncephalastrum racemosum, Chem. and Pharm. Bull., 11 (1), 1579, 85.
- Sato, D., Masuo, E. and Nozaki, Y. (1968): Production of steroids by microorganisms. Japan Patent, 68, 29, 583 (cl. 16 E 311), Dec. 18.
- Schubert, A., Langbein, G. and Siebert, R. (1957): Mikrobielle Hydroxylierung von Steroiden in 9- und 14-Stellung. Chem. Ber., 91, 2549.
- Schubert, A., Onken, D., Siebert, R. and Heller, K. (1958): Mikrobielle Hydroxylierung von Steroiden in 9- und 14-Stellung. Chem. Ber., 91, 2549.

- Schubert, A. and Siebert, R. (1960): Steroids hydroxylated in the 14 α -position. German (East) Patent, 19, 658, Aug. 11.
- Schubert, A., Siebert, R. and Langbein, G. (1960 a): Hydroxylated steroids. Ger (East) Patent, 19, 318, July 28.
- Schubert, A., Siebert, R. and Langbein, G. (1960 b): Dihydroxy- Δ^4 -oxo-steroids. Ger(East) Patent, 19, 651, Aug. 11.
- Searle, G.D. and Co. (1961): 9-Hydroxysteroids by microbiological oxidation. Brit. Patent, 862, 701, Mar. 15.
- Sebek, O.K. and Michaels, R.M. (1957): Interconversion of 17-keto and 11 β - hydroxyl groups in steroids by Protozoa. Nature, Lond., 179, 210
- Sebek, O.K. (1963): 9 α -Hydroxylation of steroids with Ascophyta. U.S. Patent, 3, 11, 116, 220.
- Shiraska, M., Tsuruta, M. and Nakamura, M. (1958): Microbiological hydroxylation of Reichstein's compound S, 17 α -hydroxydeoxycorticosterone by Sclerotinia sp. 2 β -Hydroxylation. Bull. Agri. Chem. Soc., Japan, 22, 273-4.
- Shirasaka, M. and Tsuruta, M. (1960): 11 α -Hydroxylation of steroids by Streptomyces spp. Nature, 185, 845-6.
- Shirasaka, M., Tsuruta, M., Natio, A., Sugawara, S. and Nakamura, M. (1960): Transformation of steroids by microorganisms. Takamine Kenkyusho Nempo, 11, 52-6.

- Shirasaka, M. and Tsuruta, M. (1961 a): Microbiological transformation of steroids. IV-11-Hydroxylation of steroid by Gloesporium kaki and Glomerella lagenarium. Chem. Pharm. Bull. (Tokyo), 2, 159-63.
- Shirasaka, M. and Tsuruta, M. (1961 b): Microbiological of steroids. V-Hydroxylation of steroid by Sclerotium hydroxphilum. Chem. and Pharm. Bull. (Tokyo), 2, 195-203.
- Shirasaka, M. and Tsuruta, M. (1961 c): VII-Hydroxylation of steroids by Gibberella saubinetii, 6B- and 15 α -hydroxylation. Chem. and Pharm. Bull. (Tokyo), 2, 238-45.
- Shirasaka, M. and Ozaki, M. (1961 d): Oxidation of steroids. Hakko Kyokaishi, 19, 389-93.
- Shirasaka, M., Ozaki, M. and Sugawara, S. (1961 e): Microbiological transformation of steroid compounds. J. Gen. Appl. Microbiol. (Tokyo), 7(1), 341-52.
- Shull, G.M. and Kita, D.A. (1955): Microbiological conversion of steroids. I-Introduction of the 11 β -hydroxyl group into C-21 steroids. J. Amer. Chem. Soc. 77, 763.
- Singh, K., Sehgal, S.N. and Verzina, (1967): Transformation of steroids by Mucor griseo-ovatus Con. J. Microbiol. 13(9), 1271-82.
- Smith, L.L. and Foell, T. (1962): Thin-layer chromatography of steroids on starch-bound silica gel chromatoplates. J. Chromatog., 2, 339-344.

- Smith, L.L., Marx, M., Mendelsohn, H., Foell, T. and Goodman, J. (1962): 16 α -Hydroxy steroid. XII-21-Amino derivatives of 9 α -fluorohydrocortisone. J. Amer. Chem. Soc., 84, 1265-70.
- Smith, L.L., Greenspan, G. and Rees, R. (1966): Microbiological hydroxylation of nor steroids. U.S. Patent, 3,294,646, Dec. 27.
- Soffer, J., Dorfman, R.I. and Gabrilove, J.L. (1961): In "The Human Adrenal Gland". Lea and Febiger, Philadelphia. Cited by Peterson 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York, 708 PP.
- Starka, L. and Malikova. (1961) J. Endocrinol., 22, 215, 1961. Cited by Sallam, 1966. A contribution to the utilisation of steroid sapogenins. Ph.D. Thesis, Faculty of Science Cairo University.
- Stimel, B.F., Bucknell, T.E. and Notchen, V. (1960): Federation Proc., 19, 171, 1960. Cited by Peterson, 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York, 708 PP.
- Stone, P., Hayano, M., Dorfman, R.I., Hechter, O., Robinson, C.R. and Djerassi, C. (1955): Hydroxylation of desoxycorticosterone with Neurospora crassa. J. Amer. Chem. Soc. 77, 3926.

Sweat, M.L. and Bryson, M.J. (1962): Separation of two Components of the steroid 11 β - hydroxylating system. Arch. Biochem. Biophys., 96 , 186.

Takahashi, R. (1961): Cited by Peterson, 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York , 708 PP.

Tanabe, K., Takasaki, R. and Shirasaka, M. (1959): Chem. Pharm. Bull., 7, 804-811. Cited by Peterson, 1963. In Biochemistry of industrial microorganisms, 537-606, (Rainbow, C. and Rose, A.H.), Eds., Academic Press. London and New York , 708 PP.

Thoma, R.W., Herzog, J.E., Fried, J., Perlman, D. and Gerke, J.R. (1955): Cited in Fried, J. et al., Recent Progr. Hormone Research-, 11 , 163.

Thoma, R.W., Fried, J., Bonanna, S. and Grabowicz, P. (1957): Oxidation of steroids by microorganisms. IV-16 α - Hydroxylation of 9 α -fluorohydrocortisone and 9 α - fluoroprednisolone by Streptomyces roseochromogenus. J. Amer. Chem. Soc., 79, 4818.

Thoma, R.W. and Fried, J. (1960): Microbial 7 α -hydroxylation of androstan and pregnane steroids. U.S. Patent, 2, 960, 436.

Timofeeva, A.G., Gusakova, E.G. and Shpingis, A.A. (1961):

Comparative study of steroid transformation by some molds of the imperfecti group. Izvest. Akad. Nauk, S.S.S.R., Ser Biol., 26, 574-81.

Tauda, K., Asai, T., Ohki, E., Tanaka, A. and Hattori, M. (1958):

Microbiological hydroxylation of steroids. IV-7 α , 15 β -Dihydroxylation of progesterone by Syncephalastrum. Chem. Pharm. Bull. (Tokyo), 6, 387-91.

Tsuda, K., Asai, T., Sato, Y., Tanaka, T., Matsuhisa, T. and Hasegawa, H. (1960): Microbiological hydroxylation of

steroids. X-7 and 15 β -dihydroxypregn-4-ene-3,20-dione. Chem. Pharm. Bull. (Tokyo), 8, 626-8.

Tsuda, K., Asai, T., Sato, Y., Tanaka, T. and Hasegawa, H. (1961 a): Microbiological hydroxylation of steroids.

XI-Hydroxylation of progesterone in the 7 α -, 14 α -, 15 α - or 15 β - position by Helminthosporium sativum. Chem. Pharm. Bull. (Tokyo), 9, 735-9.

Tsuda, K., Asai, T., Iizuka, H., Tanaka, T., Nakamura, M.,

Okazaki, H., Shirasaka, M. and Natio, A. (1961 b): Prednisolone. U.S. Patent, 2,993,839, July 25.

Tsuda, K., Matsui, M., Nakamura, M., Shirasaka, A. and Tsuruta, M. (1961 c): Oxidation of steroids by microorganisms.

Japan Patent, 5723, May 25.

Tsuda, K., Matsui, M., Nakamura, R., Shirasaka, A. and

Tsuruta, M. (1961 d): Oxidation of steroids by microorganisms. Japan Patent, 7, 024.

Tsuda, K., Tomoi, I., Iizuka, H. Tanaka, T., Ohki, E., Natio, A., Matsui, M., Nakamura, R. and Shirasaka, A. (1961 e): Oxidation of steroids by microorganisms. Japan Patent, 175, Jan. 17.

Tsuda, K., Hayashi, R., Takasaki, R., Shirasaka, A. and Tsuruta, M. (1963 a): Oxidation of steroid. Japan Patent, 5680, May 11.

Tsuda, K., Okazaki, H., Shirasaka, R. and Tsuruta, M. (1963 b): Oxidation of steroid. Japan Patent, 5680, May 11.

Tsuda, K., Okazaki, H., Shirasaka, R. and Tsuruta, M. (1963 c): Oxidation of steroid. Japan Patent, 8123, June 5.

Tumova, E., Siblikova, Z.O. and Hanc, O. (1955): Biochemika Oxydace Pregnanovych Steroidu, 4, 65. Cited by Sallam 1966. A contribution to the utilisation of steroid sapogenins. Ph. D. Thesis, Faculty of Science Cairo University.

Vischer, E. and Wettstein, A. (1953): Mikrobiologische Reaktionen. Seitenketten-Abbau und Dehydrierung bei Steroiden; Experientia, 2, 371.

Vondrova, O. and Hanc, O. (1960): 16 α -Hydroxylation of progesterone by a strain of Actinomyces globosus. Folia Microbiol. (Prague), 5, 247-50.

Vondrova, O. and Capek, A. (1963): Microbiological transformation of steroids. XX-Transformation reactions as a diagnostic feature in classification of Actinomyces. Folia Microbiol. (Prague), 8, 117.

- Waldi, D. (1955): Thin-layer chromatography, A Laboratory handbook, (Egon Stahl, Acad. Press, Inc., New York, London, P. 249).
- Walpole, (1941): Cited by Vogel, A.J. (1953). Text book of quantitative inorganic analysis second ed. Longmans, Green and Coy. (London-New York-Toronto).
- Weintraub, A. and Meister, P.D., (1959): Oxygenation in the 6- and 11-position of steroids by fungi. U.S. Patent, 2,879, 206, Mar. 24.
- Weisz, E., Wix, G and Bodanszky, M. (1956): Die Bildung von 11-Epicorticosterone auf mikrobiologischem Wege. Naturwissenschaften, 43, 39.
- Wintersteiner, O. and Pfiffner, J.J. (1936): Chemical studies on the adrenal cortex. III-Isolation of two new physiologically inactive compounds. J. Biol. Chem., 116, 291.
- Wix, G., Natonek, M. and Kovacs, M. (1959): Acta microbiol. Acad. Sci. Hungary, 6, 197, 1959. Cited by Peterson 1963 .cf. Ref. 147.
- Wix, G., Weisz, E., Albrecht, K., Varga, B., Kasszan, B. and Doczy, B. (1963): Hydrocortisone and prednisolone. Hung. Patent, 150, 245, May 31.
- Zaffaroni, A., Burton, R. and Kutman, E.H. (1950): Adrenal cortical hormones: Analysis by paper partition chromatography and occurrence in the urine of normal persons. Science, 111, 6-8.

- Zaffaroni, A., Campillo, C.C., Cordova, F. and Rosenkronz, G. (1955): Steroids. LXVI-Microbiological hydroxylation of steroids in position 21. *Experientia*, 11 (6), 219.
- Zaffaroni, A. and Rubin, B.A. (1957): 21-Hydroxylation of cyclopentanophenanthrene derivatives by Aspergillus niger. U.S. Patent, 2, 812, 285, Nov. 5.
- Zyoha, H. (1935): Pilze II, Mucorineae. Verlag Von Gbruder Bormtraeger, Leipzig (Neu Druck, Verlag Von. J. GRAMER, WEINHEIM, 1963).