
Rescue pct versus a conservative approach for failed fibrinolysis in patients with stemt

Mohamed Mahrous Ali Mohamed

Alpert JS, Thygesen K, Antman E, Bassand JP (2000): Myocardial infarction redefined—a consensus document of The Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. *J Am Coll Cardiol*; 36:959. Anderson HR, Nielsen TT, Rasmussen K, et al (2003): A comparison of coronary angioplasty with fibrinolytic therapy in acute myocardial infarction. *N Engl J Med*; 349:733. Anderson HV, Smalling RW, Henry TD. (2007): Drug-eluting stents for acute myocardial infarction. *J Am Coll Cardiol*; 49:1931. Anderson J, Adams C, Antman E, et al. (2007): ACC/AHA -2007 guidelines for the management of patients with unstable angina/non-ST elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to revise the 2002 Guidelines for the Management of Patients with Unstable Angina/Non-ST-Elevation Myocardial Infarction): developed in collaboration with the American College of Emergency Physicians, American College of Physicians, Society for Academic Emergency Medicine, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*; 50:el. Available at: [www.acc.org/qualityandscience/ clinical/ statements. htm](http://www.acc.org/qualityandscience/clinical/statements.htm). Antman EM, Anbe DT, Armstrong PW, et al, (1999): Abciximab facilitates the rate and extent of thrombolysis: Result of thrombolysis in myocardial infarction (TIMI) 14 trial. *Circulation*; 99: 2720-25. Antman EM, Anbe DT, Armstrong PW, et al. (2004): ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction; a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1999 Guidelines for the Management of Patients with Acute Myocardial Infarction). *J Am Coll Cardiol*; 44: 671-719. Antman EM, Anbe DT, Armstrong PW, et al. (2006): ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction. Available at: [www.acc.org/ qualityandscience/ clinical/ stateinents. htm](http://www.acc.org/qualityandscience/clinical/statements.htm). Antman EM, Hand M, Armstrong PW, et al. (2007): focused update of the ACC/AHA 2004 Guidelines for the Management of Patients With ST- Elevation Myocardial Infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Group to Provide New Evidence and Update the ACC/AHA 2004 Guidelines for the Management of Patients With ST-Elevation Myocardial Infarction). *T Am Coll Cardiol* 2008;51:210-47. Antman EM, Hand M, Armstrong PW, et al. (2008): focused update of the ACC/AHA 2004

Guidelines for the Management of Patients With ST-Elevation Myocardial Infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Group to Review New Evidence and Update the ACC/AHA 2004 Guidelines for the Management of Patients With ST-Elevation -Myocardial Infarction). *J Am Coll Cardiol* 2008;51: -XXX.

Armstrong PW, Fu Y, Chang W C, et al. (1998): for the GUSTO-IIb Investigators. Acute coronary syndromes in the GUSTO-IIb trial: Prognostic insights and impact of recurrent ischemia. *Circulation*; 98:1860-66.

Belenkie I, Traboulsi M, Hall CA, et al. (1992): Rescue angioplasty during myocardial infarction has a beneficial effect on mortality: a tenable hypothesis. *Can J Cardioi*; 8: 357-62.

Boersma E, Maas ACP, Deckers JW, et al. (1996): Early thrombolytic treatment in acute myocardial infarction: Reappraisal of the golden hour. *Lancet*; 348:771-78.

Brass LM, Lichtman JH, Wang Y, et al. (2000): Intracranial hemorrhage associated with thrombolytic therapy for elderly patients with acute myocardial infarction. Results from the Cooperative Cardiovascular Project. *Stroke*; 31:1802.

Brodie BR, Stuckey TD, Wall TC, et al. (1998): Importance of time to reperfusion for 30 day and late survival and recovery of left ventricular function after primary angioplasty for acute myocardial infarction. *J Am Coll Cardiol* ;32:1312-19.

Burjonrappa Sukesh C, Paul D, Varosy, et al, (2011): Survival of patients undergoing rescue percutaneous coronary intervention *J Am Coll cardiol* 4; 42-50.

Califf RM, O'Neill W, Stack Steward RS et al, (1988): Failure of simple clinical measurements to predict perfusion status after intravenous thrombolysis. *Ann Intern Med*; 108:658-62.

Cannon CP, Bahit MC, Haugland JM, et al. (2002): Underutilization of evidence-based medications in acute ST elevation myocardial infarction. Results of the Thrombolysis in Myocardial Infarction (TIMI) 9 Registry. *Cric Pathways Cardiol*; 1: 44-50.

CZitrom D, Karila-Cohen D, Brochet E, et al. (1999): Acute assessment of micro vascular perfusion patterns by myocardial contrast echocardiography during myocardial infarction: relation to timing and extent of functional recovery. *Heart*; 81:12-16.

Davies CH, Ormerod OJ. (1998): Failed coronary thrombolysis. *Lancet* 351:1191-96.

De Luca G, Suryapranata H, van't Hof AW, et al. (2004)a: Prognostic assessment of patients with acute myocardial infarction treated with primary angioplasty: implications for early discharge. *Circulation*; 109:2737.

Eagle KA, Goodman SG, Avezum A, et al, (2002): Practice variation and missed opportunities for reperfusion in ST- segment-elevation myocardial infarction: findings from the Global Registry of acute coronary Events (GRACE). *Lancet*: 359: 373-77.

Ellis S.G., E.R. Da Silva and G. Heyndrickx, et al. (1994): Randomized comparison of rescue angioplasty with conservative management of patients with early failure of thrombolysis for acute anterior myocardial infarction. *Circulation*, 90, 2280-2284.

Fesmire FM, Percy RF, Bardoner JB, et al. (1998): Usefulness of automated serial 12-lead ECG monitoring during the initial emergency department evaluation of patients with chest pain. *Ann Emerg Med*; 31:3-6.

Fibrinolytic Therapy Trialists' (FTT) Collaborative Group. (1994): Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet*; 343:311-17.

Gerschlick AH, Stephens-Lloyd A, Hughes S, et al, (2005): Rescue angioplasty after failed thrombolytic therapy for acute myocardial infarction. *N Engl J Mea*;353:2758-68.

Gersh BJ, Stonwe GW, White HD and Holmes DR Jr.

(2005): Pharmacological facilitation of primary percutaneous coronary intervention for acute myocardial infarction: is the slope of the curve the shape of the future? JAMA; 293: 979.

Gibson CM, Canon CP, Greene RM, Sequeria RF, Margoriln -^ RD, Leya F et al.(1997): Rescue angioplasty in thrombolysis in myocardial infarction (ITVffiV) trial. Am J Cardiol ;80:21-26.

Gibson CM, Karha J, Murphy SA, et al. (2003): Early and long-term clinical outcomes associated with reinfarction following fibrinolytic administration in the Thrombolysis in Myocardial Infarction trials. J Am Coll Cardiol; 42:2979-68.

GISSI (1986): Effectiveness of intravenous thrombolytic treatment in acute myocardial infarction. Results of the Gruppo Italiano per lo Studio della Streptochinasi nell'Infarto Miocardico. Lancet; 1:397.

Go AS, Barren HV, Rundle AC, et al. (1998): Bundle-branch block and in-hospital mortality in acute myocardial infarction. National Registry of Myocardial Infarction 2 Investigators. Ann Intern Med; 129:690.

Goldman LE, Eisenbeng MJ. (2000): Identification and management of patients with failed thrombolysis after acute myocardial infarction. Ann intern Med; 132: 556-65.

Goodman SG, Steg PG, Eagle KA, et al. (2006): The diagnostic and prognostic impact of the redefinition of acute myocardial infarction: lessons from the Global Registry of Acute Coronary Events (GRACE). Am Heart J; 151:654-60.

Gore JM, Granger CB, Simoons ML, et al. (1998): for the GUSTO-I Investigators. Stroke after thrombolysis: mortality and functional outcomes in the GUSTO-I trial. Circulation; 92:2811.

Hamm CW, Goldmann B, Heeshen C, et al. (2002): Emergency room triage of patients with acute chest pain by means of rapid testing for cardiac troponin T or troponinI. N Engl J Med; 337:1648.

Hudson MP, Granger CB, Topol EJ, et al. (2001): Early reinfarction after fibrinolysis: experience from the global utilization of streptokinase and tissue plasminogen activator (alteplase) for occluded coronary arteries (GUSTO I) and global use of strategies to open occluded coronary arteries (GUSTO HD trials Circulation; 104:1229.

ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. (1988): Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. Lancet; 2:349.

Jaffe AS, Ravkilde J, Roberts R, et al. (2000): It's time for a change to a troponin standard. Circulation; 102:1216.

Juliard JM, Himbert D, Cnstofuu P, et al. (1999): A matched comparison of prehospital thrombolysis and stand by rescue angioplasty with primary angioplasty. Am J cardiol ; 83: 305-10.

Kastrati A, Mehilli J, Nekolla S, et al. (2004): A randomized trial comparing myocardial salvage achieved by coronary stenting versus balloon angioplasty in patients with acute myocardial infarction considered ineligible for reperfusion therapy. J Am Coll Cardiol; 43:734.

Kaul U, Sapra R, Singh B, et al, (2001): Abciximab during rescue angioplasty after failed thrombolysis. Asian cardiovasc thorac Ann; 9: 31-35.

Keeley EC, Boura JA, Grines CL. (2006): Comparison of primary and facilitated percutaneous coronary interventions for ST-elevation myocardial infarction: quantitative review of randomised trials. Lancet; 367:579-88.

Keeley EC, Boura JA, Grines CL.(2003): Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. Lancet; 361:13-20.

Klootwijk P, Langer A, Meij S, Browne KF. (1996): Noninvasive prediction of reperfusion and coronary artery patency by continuous ST segment monitoring in GUSTO I trial. Eur Heart ;17:689-98.

Kudenchuk PJ, Maynard C, Cobb LA, et al. (1998): for the MITI

Investigators. Utility of the prehospital electrocardiogram in diagnosing acute coronary syndromes; The Myocardial infarction Triage and Intervention Project. *J Am Coll Cardiol*; 32:17. Massel D. (2005): Primary angioplasty in acute myocardial infarction: hypothetical estimate of superiority over aspirin or untreated controls. *Am J Med*; 118:113. Matezky S, Freimark D, Chouraqui P, et al. (1998): The distinction between coronary and myocardial reperfusion after thrombolytic therapy by clinical markers of reperfusion. *J Am Coll Cardiol* 32:1326-30. Mazzota G, Protosido SK, Casati E. (1998): Myocardial scintigraphy in acute myocardial infarction treated with systemic thrombolysis: how far are we from obtaining reliable information for rescue PTCA? *Int J Cardiol* ;29(Suppl I): S69-73. Mehta RH, Harjai KJ, Cox D, et al. (2003): Clinical and angiographic correlates and outcomes of suboptimal coronary flow inpatients with acute myocardial infarction undergoing primary percutaneous coronary intervention. *J Am Coll Cardiol*; 42:1739. Meier MA, Al-Badr WH, Cooper JV, et al. (2002): The new definition of myocardial infarction: Diagnostic and prognostic implications in patients with acute coronary syndromes. *Arch Intern Med*; 162:1585. Montori VM, Devereaux PJ, Adhikari NK, et al. (2005): Randomized trials stopped early for benefit: a systematic review. *JAMA*; 294: 2203-9. Mounsey JP, Skinner IS, Hawkins T, et al. (1995): Rescue thrombolysis; alteplase as adjuvant treatment after streptokinase in acute myocardial infarction. *Br Heart*; 74, 348-53. Nallamothu BK, Wang Y, Magid DJ, et al. (2006): Relation between hospital specialization with primary percutaneous coronary intervention and clinical outcomes in ST-segment elevation myocardial infarction: National Registry of Myocardial Infarction-4 analysis. *Circulation*; 113:222-9. Ndrepepa G, Mehilli J, Schulz S, et al. (2008): Prognostic significance of epicardial blood flow before and after percutaneous coronary intervention in patients with acute coronary syndromes. *J Am Coll Cardiol* ;52:512-17. Newby LK, Roe MT, Chen AY, et al. (2006): Frequency and clinical implications of discordant creatine kinase-MB and troponin measurements in acute coronary syndromes. *J Am Coll Cardiol*; 47:312. Nordmann A, Bucher H, Hengstler P, et al. (2005): Primary stenting versus primary balloon angioplasty for treating acute myocardial infarction. *Cochrane Database Syst Rev*; CD005313. Ohman EM, Califf RM, Topol EJ, et al. (1990): Consequences of reocclusion after successful reperfusion therapy in acute myocardial infarction: the TAMI Study Group. *Circulation*; 82:781. Ortolan P, Marzocchi A, Marrozzini C, et al. (2006): Clinical impact of direct referral to primary percutaneous coronary intervention following pre-hospital diagnosis of ST-elevation myocardial infarction. *Eur Heart J*; 27:1550. Pinto DS, Kirtane AJ, Nallamothu BK, et al. (2006): Hospital delays in perfusion in ST-elevation myocardial infarction: implications when selecting a reperfusion strategy. *Circulation*; 114:2019. Pope JH, Ruthazer R, Beshansky JR, et al. (1998): Clinical features of emergency department patients presenting with symptoms suggestive of acute cardiac ischemia: a multicenter study. *J Thromb Thrombolysis*; 6:63. Prendergast BD, Shandall A, Buchalter MB. (1997): What do we do when thrombolysis fails? A United Kingdom survey. *Int J Cardiol*; 61: 39-42. Rathore SS, Weinfurt KP, Gersh BJ, et al. (2001): Treatment of patients with myocardial infarction who present with a paced rhythm. *Ann Intern Med*; 134:644. Ross AM, Coyne KS, Moreyra E, et al. (1998): For the GUSTO-I Angiographic Investigators. Extended mortality benefit of early postinfarction

reperfusion. *Circulation*; 97:1549. Ross AM, Lundergan CF, Rohrbeck SC, Boyle DH, Brand M, Buller CH et al. (1998): Rescue angioplasty after failed thrombolysis: technical and clinical outcomes in a large thrombolysis trial. *J Am Coll Cardiol*; 31:1511-7. Sarallo FM, Americo L, Di pasquala P, et al (2000): Efficacy of rescue thrombolysis in patients with acute myocardial infarction preliminary finding. *Cardiovasc drugs ther*; 14: 83-9. Sarda L, Colin P, Boccara F, et al. (2001): Myocarditis in patients with clinical presentation of myocardial infarction and normal coronary angiograms. *J Am Coll Cardiol*; 37:786. Schomig A, Ndrepepa G, Mehilli J, et al. (2003): Therapy-dependent influence of time-to-treatment interval on myocardial salvage in patients with acute myocardial infarction treated with coronary artery stenting or thrombolysis. *Circulation*; 108:1084-8. Schomig A, Ndrepepa J, Mehilli J, et al. (2004): A randomized trial of coronary stenting versus balloon angioplasty as a rescue intervention after failed thrombolysis in patients with acute myocardial infarction. *J Am Coll Cardiol*; 44:2073. Schomig A, Mehilli J, Antoniucci D, et al (2005): Mechanical reperfusion in patients with acute myocardial infarction presenting more than 12 hours from symptom onset. *JAMA*; 293: 2865-2872. Sgarbossa BB, Pinski SL, Barbagelatu A, (1996): Electrocardiographic diagnosis of evolving acute myocardial infarction in the presence of left bundle branch block. *N Engl J med*; 334: 481-486. Shavelle DM, Salami A, Abdelkarim M, et al. (2006): Rescue percutaneous coronary intervention for failed thrombolysis. *Cathet Cardiovasc Int*; 67:214-20. Shlipak MG, Go AS, Frederick PD, et al. (2000): Treatment and outcomes of left bundle-branch block patients with myocardial infarction who present without chest pain. National Registry of Myocardial Infarction 2 Investigators. *J Am Coll Cardiol*; 36:706. Steg PG, Francois L, Jung B, et al. (2007): Long-term clinical outcomes after rescue angioplasty are not different from those of successful thrombolysis for acute myocardial infarction. *Eur Heart J*; 26:1831-7. Stewart JT, French JK, Theroux P, et al. (1998): Early noninvasive identification of failed reperfusion after intravenous thrombolytic therapy in acute myocardial infarction. *J Am Coll Cardiol*; 31:1499-505. Stone GW, Brodie BR, Griffin JJ, et al. (1999): For the PAMI Stent Pilot Trial Investigators. Clinical and angiographic follow-up after primary stenting in acute myocardial infarction: The Primary Angioplasty in Myocardial Infarction (PAMI) stent pilot trial. *Circulation*; 99:1548. Stone GW, Grinesel, Cox DA, et al, (2002): Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med*; 346: 957-10. Sutton AG, Campbell PG, Graham R, et al. (2004): A randomized trial of rescue angioplasty versus a conservative approach for failed fibrinolysis in ST segment elevation myocardial infarction: the Middlesbrough Early Revascularization to Limit Infarction (MERLIN) trial. *J Am Cardiol*; 44:287-96. Taher T, Fu Y, Wagner GS, et al. (2004): Aborted myocardial infarction in patients with ST-segment elevation: insights from the Assessment of the Safety and Efficacy of a New Thrombolytic Regimen-3 Trial Electrocardiographic Substudy. *J Am Coll Cardiol*; 44:38. Testa L, Vangaol WJ, G.G.L Biondi-Zocni, Abbate A, Agostoni P, Bhindi R, and Banming A., P., (2008): Repeat thrombolysis or conservative therapy Vs. rescue PCI for failed thrombolysis: Systematic review and meta-analysis. *Oxford Journals Medicin*; 101,387-395. The GUSTO Angiographic Investigators. (1994): The effects of tissue plasminogen activator, streptokinase, or both on coronary artery patency,

ventricular function, and survival after acute myocardial infarction. N Engl J Med 1993; 329:1615-22 [Erratum, N Engl J Med; 330:516.]. The GUSTO Investigators. (1993): An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. N. Engl J Med ;239:673-82. The GUSTO Investigators. (1993): The effects of tissue plasminogen activator, streptokinase, or both on coronary-artery patency, ventricular function, and survival after acute myocardial infarction. N Engl J Med; 329:1615-22. The TIMI Research Group. (1988): Immediate versus delayed catheterization and angioplasty following thrombolytic therapy for acute myocardial infarction: TIMI II A results. JAMA; 260: 2849-58. Thygesen K, Alpert JS, White HD, et al. (2007): Universal definition of myocardial infarction: Kristian Thygesen, Joseph S. Alpert and Harvey D. White on behalf of the Joint ESC/ACCF/AHA/WHF Task Force for the Redefinition of Myocardial Infarction. Eur Heart J; 28:2525. Van de Merkhof LF, Zijlstra F, Olsson H, et al. (1999): Abciximab in the treatment of acute myocardial infarction eligible for primary percutaneous coronary angioplasty. Results of the glycoprotein receptor antagonist patency evaluation (GRAPE) study. J Am Coll Cardiol; 33:1528-32. Van de Werf F, Ardissino D, Betriu A, et al. (2003): Management of acute myocardial infarction in patients presenting with ST-segment elevation. The Task Force on the Management of Acute Myocardial Infarction of the European Society of Cardiology. Eur Heart J; 24:28-66. Van de Werf F, Bax J, Betriu A, et al. (2008): Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. Eur Heart J ;29:2909-45. Verheugt FW, Gersh BJ, Armstrong PW. (2006): Aborted myocardial infarction: a new target for reperfusion therapy. Eur Heart J; 27:901.